

\* Lower Triangular Matrix:- If all elements above the principal diagonals are zeroes.

$$\begin{bmatrix} 1 & 0 & 0 \\ 2 & 3 & 0 \\ 1 & 2 & 5 \end{bmatrix} \quad \Delta = 15$$

\* Upper Triangular Matrix:- If all elements below the principal diagonal are zeroes, then it is said to be an upper triangular matrix.

$$\begin{bmatrix} 1 & 2 & 3 \\ 0 & -4 & 2 \\ 0 & 0 & 7 \end{bmatrix} \quad \Delta = -28$$

NOTE:-

If a matrix is either lower triangular or upper triangular then the determinant is the product of the principle diagonal elements.

Q. Find the determinant of Matrix

$$\begin{vmatrix} 1 & a & a^2 \\ 1 & b & b^2 \\ 1 & c & c^2 \end{vmatrix}$$

$$\rightarrow \begin{vmatrix} \textcircled{1} & a & a^2 \\ \textcircled{1} & b & b^2 \\ 1 & c & c^2 \end{vmatrix} \quad \begin{array}{l} R_1 \rightarrow R_1 - R_2 \\ R_2 \rightarrow R_2 - R_3 \end{array}$$

$$= \begin{vmatrix} 0 & a-b & a^2-b^2 \\ 0 & b-c & b^2-c^2 \\ 1 & c & c^2 \end{vmatrix}$$

$$= (a-b)(b-c) \begin{vmatrix} 0 & 1 & a+b \\ 0 & 1 & b+c \\ 1 & c & c^2 \end{vmatrix} = (a-b)(b-c) [1 \cdot (b+c - a) - 1 \cdot (c^2 - a^2)]$$

$$= (a-b)(b-c)(c-a)$$

\*

$$\boxed{|A| = |A^T|}$$

$$\begin{vmatrix} 1 & 1 & 1 \\ a & b & c \\ a^2 & b^2 & c^2 \end{vmatrix} = \begin{vmatrix} 1 & a & a^2 \\ 1 & b & b^2 \\ 1 & c & c^2 \end{vmatrix}$$

$$|A| = |A^T| = (a-b)(b-c)(c-a)$$

Imp Result...

Q.  $\begin{vmatrix} 1 & 1 & 1 \\ 1 & 1+a & 1 \\ 1 & 1 & 1+b \end{vmatrix} \quad R_2 - R_1, R_3 - R_1$

$$= \begin{vmatrix} 1 & 1 & 1 \\ 0 & a & 0 \\ 0 & 0 & b \end{vmatrix} = \underline{\underline{ab}}$$

Q.  $\begin{vmatrix} 1 & 1 & 1 \\ 1 & 5 & 1 \\ 1 & 1 & 6 \end{vmatrix} = ?$

- A 20    B 30    C 40    D 0

Imp Result

Q.  $\begin{vmatrix} 1 & 1 & 1 & 1 \\ 1 & 1+a & 1 & 1 \\ 1 & 1 & 1+b & 1 \\ 1 & 1 & 1 & 1+c \end{vmatrix} = abc$

Q. 
$$\begin{vmatrix} 1+a & 1 & 1 \\ 1 & 1+b & 1 \\ 1 & 1 & 1+c \end{vmatrix}$$

Imp  
Result

$$= abc \begin{vmatrix} 1+\frac{1}{a} & \frac{1}{a} & \frac{1}{a} \\ \frac{1}{b} & 1+\frac{1}{b} & \frac{1}{b} \\ \frac{1}{c} & \frac{1}{c} & 1+\frac{1}{c} \end{vmatrix} \quad R_1 \rightarrow R_1 + (R_2 + R_3)$$

$$= abc \left( 1 + \frac{1}{a} + \frac{1}{b} + \frac{1}{c} \right) \begin{vmatrix} 1 & 1 & 1 \\ \frac{1}{b} & 1+\frac{1}{b} & \frac{1}{b} \\ \frac{1}{c} & \frac{1}{c} & 1+\frac{1}{c} \end{vmatrix} \quad C_2 - C_1, C_3 - C_1$$

$$= abc \left( 1 + \frac{1}{a} + \frac{1}{b} + \frac{1}{c} \right) \begin{vmatrix} 1 & 0 & 0 \\ \frac{1}{b} & 1 & 0 \\ \frac{1}{c} & 0 & 1 \end{vmatrix}$$

$$= abc \left( 1 + \frac{1}{a} + \frac{1}{b} + \frac{1}{c} \right)$$

Q. 
$$\begin{vmatrix} 1+a & 1 & 1 & 1 \\ 1 & 1+b & 1 & 1 \\ 1 & 1 & 1+c & 1 \\ 1 & 1 & 1 & 1+d \end{vmatrix}$$

Imp  
Result

$$= abcd \left( 1 + \frac{1}{a} + \frac{1}{b} + \frac{1}{c} + \frac{1}{d} \right)$$

Q. 
$$\begin{vmatrix} 2 & 1 & 1 & 1 \\ 1 & 2 & 1 & 1 \\ 1 & 1 & 2 & 1 \\ 1 & 1 & 1 & 2 \end{vmatrix} = ?$$

(A) 4 (B) 5 (C) 6 (D) 7

Q. 
$$\begin{vmatrix} 0 & 1 & 2 \\ 1 & 2 & 3 \\ 3 & 1 & 1 \end{vmatrix} = 0(1-9) - 1(1-9) + 2(1-6) = 8-10 = -2$$

\* SHORTCUT to determinant of Matrix:-

✓ [Applicable only for 3x3 Matrix]

Q. 
$$\begin{bmatrix} 0 & 1 & 2 \\ 1 & 2 & 3 \\ 3 & 1 & 1 \end{bmatrix} \begin{matrix} 0 & 1 \\ 1 & 2 \\ 3 & 1 \end{matrix}$$

$$0 + 9 + 2 - 12 - 0 - 1 = -2$$

Q. 
$$\begin{bmatrix} 1 & 2 & 2 \\ 2 & 1 & 2 \\ 2 & 2 & 1 \end{bmatrix} \begin{matrix} 1 & 2 \\ 2 & 1 \\ 2 & 2 \end{matrix}$$

$$\Delta = 1 + 8 + 8 - 4 - 4 - 4$$

$$\Delta = 17 - 12 = 5$$

Q. 
$$\begin{bmatrix} 1 & 2 & -2 \\ -1 & 3 & 0 \\ 0 & -2 & 1 \end{bmatrix} \begin{matrix} 1 & 2 \\ -1 & 3 \\ 0 & -2 \end{matrix}$$

$$\Delta = 3 + 0 - 4 - 0 - 0 + 2$$

$$\Delta = 1$$

$$Q. \begin{bmatrix} 1 & 2 & 5 \\ 3 & 1 & 4 \\ 1 & 1 & 2 \end{bmatrix}$$

$$\Delta = 2 + 8 + 15 - 5 - 4 - 12$$

$$\Delta = 25 - 21$$

$$\Delta = 4$$

### # INVERSE OF MATRIX!-

$$A^{-1} = \frac{\text{adj } A}{\Delta}$$

$$\text{if } A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$$

$$A^{-1} = \frac{1}{ad-bc} \begin{bmatrix} d & -b \\ -c & a \end{bmatrix}$$

Principle  
 → Interchange diagonal element  
 → & sign change in second diagonal element.

$$Q. A = \begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix}$$

$$A^{-1} = \frac{1}{4-6} \begin{bmatrix} 4 & -2 \\ -3 & 1 \end{bmatrix}$$

$$Q. A = \begin{bmatrix} 5 & 4 \\ 1 & 2 \end{bmatrix}$$

$$A^{-1} = \frac{1}{10-4} \begin{bmatrix} 2 & -4 \\ -1 & 5 \end{bmatrix}$$

$$Q. A = \begin{bmatrix} \cos\theta & -\sin\theta \\ \sin\theta & \cos\theta \end{bmatrix}$$

$$A^{-1} = \frac{1}{\cos^2\theta + \sin^2\theta} \begin{bmatrix} \cos\theta & \sin\theta \\ -\sin\theta & \cos\theta \end{bmatrix}$$

$$= \begin{bmatrix} \cos\theta & \sin\theta \\ -\sin\theta & \cos\theta \end{bmatrix}$$

# For 3x3 Inverse matrix :-

$$A^{-1} = \frac{\text{adj } A}{\Delta}$$

adj A = Transpose of cofactors Matrix

Cofactor of elements =  $(-1)^{i+j}$  minor

↳ Just to check sign for even or odd (row+column) no

minor of an Element = determinant of sq. sub matrix in which the row & column of the particular element lines to be deleted.

ex:-

$$\begin{bmatrix} 0 & 1 & 2 \\ \textcircled{1} & 2 & 3 \\ 3 & 1 & 1 \end{bmatrix}$$


$$\text{minor of } 1 \rightarrow \begin{vmatrix} 1 & 2 \\ 1 & 1 \end{vmatrix} = 1 - 2 = -1$$

$$\text{Cofactor of } 1 = (-1)^{2+1} (-1) = \underline{\underline{1}}$$

\* Shortcut to find Cofactor of 3x3 Matrix:-

Q. 
$$\begin{bmatrix} 0 & 1 & 2 \\ 1 & 2 & 3 \\ 3 & 1 & 1 \end{bmatrix}$$


$$\Delta = 0 + 9 + 2 - 12 - 0 - 1 = -2$$



$$A^{-1} = \frac{1}{-2} \begin{bmatrix} -1 & 1 & -1 \\ 8 & -6 & 2 \\ 5 & 3 & -1 \end{bmatrix}$$

Q. 
$$\begin{bmatrix} 1 & 2 & 2 \\ 2 & 1 & 2 \\ 2 & 2 & 1 \end{bmatrix}$$

$$\Delta = 1 + 8 + 8 - 4 - 4 - 4 = 17 - 4 = 13$$



$$A^{-1} = \frac{1}{13} \begin{bmatrix} -3 & 2 & 2 \\ 2 & -3 & 2 \\ 2 & 2 & -3 \end{bmatrix}$$

Q.

$$\begin{bmatrix} 1 & 2 & -2 \\ -1 & 3 & 0 \\ 0 & -2 & 1 \end{bmatrix}$$

$$\Delta = 0 \times 0 \times (-2) - 0 \times 3 + 0 - 0 - 0 + 2 = 4.$$

$$\begin{array}{c|ccccc} 1 & 2 & -2 & 1 & 2 \\ -1 & 3 & 0 & -1 & 3 \\ 0 & -2 & 1 & 0 & -2 \\ \hline 1 & 2 & -2 & 1 & 2 \\ -1 & 3 & 0 & -1 & 3 \end{array}$$

$$A^{-1} = \begin{bmatrix} 3 & 2 & 6 \\ 1 & 1 & 2 \\ 2 & 2 & 5 \end{bmatrix}$$

Q.

$$\begin{bmatrix} 1 & 2 & 5 \\ 3 & 1 & 4 \\ 1 & 1 & 2 \end{bmatrix}$$

$$\Delta = 2 + 8 + 15 - 5 - 4 - 12 = 25 - 21 = 4$$

$$\begin{array}{c|ccccc} 1 & 2 & 5 & 1 & 2 \\ 3 & 1 & 4 & 3 & 1 \\ 1 & 1 & 2 & 1 & 1 \\ \hline 1 & 2 & 5 & 1 & 2 \\ 3 & 1 & 4 & 3 & 1 \end{array}$$

$$A^{-1} = \frac{1}{4} \begin{bmatrix} -2 & 1 & 3 \\ -2 & -3 & 11 \\ 2 & 1 & -5 \end{bmatrix}$$

$$\frac{Q}{\left[ \begin{array}{ccc} \cos \theta & -\sin \theta & 0 \\ \sin \theta & \cos \theta & 0 \\ 0 & 0 & 1 \end{array} \right]}$$

$$Q \left[ \begin{array}{ccc} \cos \theta & -\sin \theta & 0 \\ \sin \theta & \cos \theta & 0 \\ 0 & 0 & 1 \end{array} \right]$$

$$\Delta = \cos^2 \theta + \sin^2 \theta = 1$$

$$A^{-1} = \begin{array}{c|cc} \cos \theta & -\sin \theta & 0 \\ \sin \theta & \cos \theta & 0 \\ 0 & 0 & 1 \end{array} \begin{array}{c} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \\ 0 & 0 \\ \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{array}$$

$$Q \left[ \begin{array}{ccc} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 1 \end{array} \right]$$

$$\begin{array}{c|ccc} 0 & 1 & 0 & 0 & 1 \\ \hline 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 1 & 0 & 0 \\ 0 & 1 & 0 & 0 & 1 \\ 1 & 0 & 0 & 1 & 0 \end{array}$$

$$A^{-1} = \frac{1}{-1} \begin{bmatrix} 0 & -1 & 0 \\ -1 & 0 & 0 \\ 0 & 0 & -1 \end{bmatrix}$$

$$= \begin{bmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

## # RANK OF MATRIX:-

If all minors of order  $(r+1)$  are zeroes, but there is at least one non zero minor of order  $r$ . if exists, It is called the rank of matrix and is denoted by  $\boxed{p(A) = r}$

### \* Properties of Rank:-

i) If  $A$  is a null matrix or zero matrix then rank of  $A = 0$

ex:-  $\begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}$ ,  $\begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}$ ,  $\begin{bmatrix} 0 & 0 \\ 0 & 0 \\ 0 & 0 \end{bmatrix}$ ,  $\begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}$  etc.

ii) If  $A$  is a non zero matrix, then rank of  $A \geq 1$

iii) If 'I' be the unit matrix or identity matrix of order  $n \times n$ , then rank of  $I_n = n$

ex:-  $I_2 = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$  rank = 2

$I_3 = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$  rank = 3

$I_4 = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$  rank = 4.

iv) If  $A$  is a matrix of order  $m \times n$ , then rank of  $A \leq \min\{m, n\}$

ex:-  $A = \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \end{bmatrix}$   
2x3

rank  $\leq 2$  ✓

\* If all determinants of  $2 \times 2 = 0$   
rank  $< 2$

\* If at least one determinant of  $2 \times 2$   
is non zero, then  
rank = 2.

Q. Find rank of  $\begin{bmatrix} 1 & 2 & 3 & 4 \\ -2 & 0 & 5 & 7 \end{bmatrix}_{2 \times 4}$

$$\rho(A) \leq 2$$

By checking determinant of any <sup>one of</sup>  $2 \times 2 \neq 0$  here

$$\rho(A) = 2$$

\* \*

Determinant checking

1<sup>st</sup> check = 0

2<sup>nd</sup> check = 0

3<sup>rd</sup> check = 0

4<sup>th</sup> check (no need)

of course there must be rows identical with some common factors.

$\therefore$  rank will  $< \min\{m, n\}$   
 $\neq \min\{m, n\}$

Q.  $\begin{bmatrix} 1 & 2 & 3 \\ 1 & 4 & 2 \\ 2 & 6 & 5 \end{bmatrix}$

\* purpose  $\rightarrow$  make at least one row = 0

$$\begin{bmatrix} 1 & 2 & 3 \\ 1 & 4 & 2 \\ 2 & 6 & 5 \end{bmatrix}$$

$$\rho(A) \leq 3$$

$$R_3 - (R_1 + R_2) \rightarrow \neq 0 \Rightarrow \rho(A) = 2.$$

$$\begin{bmatrix} 1 & 2 & 3 \\ 1 & 4 & 2 \\ 0 & 0 & 0 \end{bmatrix}$$

Q.  $\begin{bmatrix} 1 & 1 & 1 \\ 1 & 1 & 1 \\ 1 & 1 & 1 \end{bmatrix}$   $\neq 0 \Rightarrow P(A) = 2.$   
 $P(A) \leq 3$   
 Can be made 0  
 $3 \times 3$

Q.  $\begin{bmatrix} 1 & 1 & 0 \\ 0 & 1 & 0 \\ 2 & 2 & 0 \end{bmatrix}$   $\neq 0 \Rightarrow P(A) = 2.$   
 $P(A) \leq 3$   
 $3 \times 3$

Q.  $\begin{bmatrix} 1 & 2 & 3 \\ -1 & 2 & 2 \\ 0 & 5 & 6 \end{bmatrix}$   $P(A) \leq 3$   
 $P(A) = 3.$   
 $3 \times 3$

Q.  $\begin{bmatrix} 2 & 3 & 4 & -1 \\ 5 & 2 & 0 & -1 \\ -4 & 5 & 12 & -1 \end{bmatrix}$   $P(A) \leq 3.$   
 $R_3 - 3R_1$   
 $3 \times 4$

$\begin{bmatrix} 2 & 3 & 4 & -1 \\ 5 & 2 & 0 & -1 \\ -10 & -4 & 0 & 2 \end{bmatrix}$   
 $= 0 \Rightarrow P(A) \leq 3$   
 $P(A) = 2.$

Q.  $\begin{bmatrix} 1 & -1 & 3 & 6 \\ 1 & 8 & -3 & -4 \\ 5 & 3 & 3 & 11 \end{bmatrix}$   $P(A) \leq 3.$   
 $R_2 + R_1$   
 $R_3 - R_1$   
 $3 \times 4$

$\begin{bmatrix} 1 & -1 & 3 & 6 \\ 2 & 7 & 0 & 2 \\ 4 & 4 & 0 & 5 \end{bmatrix}$   
 $\neq 0 \Rightarrow P(A) = 3$

Q. 
$$\begin{bmatrix} 1 & 3 & 4 & 3 \\ 3 & 9 & 12 & 3 \\ \hline 1 & 3 & 4 & 1 \end{bmatrix}$$

$P(A) = 2$

$\Rightarrow P(A) \leq 3$   
 $P(A) < 3 \quad \left\{ \begin{array}{l} R_2, R_1 \text{ identical} \end{array} \right\}$

Q. 
$$\begin{bmatrix} 1 & 2 & 3 & 0 \\ 2 & 4 & 3 & 2 \\ 3 & 2 & 1 & 3 \\ 6 & 8 & 7 & 5 \end{bmatrix}$$

4x4

$P(A) \leq 4$   
 $R_4 - (R_1 + R_2 + R_3)$

$$\begin{bmatrix} 1 & 2 & 3 & 0 \\ 2 & 4 & 3 & 2 \\ 3 & 2 & 1 & 3 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

$P(A) < 4$   
 $R_2 - 2R_1$

$$\begin{bmatrix} 1 & 2 & 3 & 0 \\ 0 & 0 & -3 & 2 \\ 3 & 2 & 1 & 3 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

$\neq 0 \Rightarrow P(A) = 3.$

Q. 
$$\begin{bmatrix} 2 & 3 & -1 & -1 \\ 1 & -1 & -2 & -4 \\ 3 & 1 & 3 & -2 \\ 6 & 3 & 0 & -7 \end{bmatrix}$$

$P(A) \leq 4$

$R_{12}$

$$\begin{bmatrix} 1 & -1 & -2 & 4 \\ 2 & 3 & -1 & -1 \\ 3 & 1 & 3 & -2 \\ 6 & 3 & 0 & -7 \end{bmatrix}$$

$R_2 - 2R_1, R_3 - 3R_1$   
 $R_4 - (R_1 + R_2 + R_3)$

$$= \left[ \begin{array}{ccc|c} 1 & -1 & -2 & -4 \\ 0 & 5 & 3 & 7 \\ 0 & 4 & 9 & 10 \\ 0 & 0 & 0 & 0 \end{array} \right]$$

$$r(A) < 4$$

$$\neq 0 \Rightarrow r(A) = 3$$

$$0. \left[ \begin{array}{cccc} 6 & 1 & 3 & 8 \\ 4 & 2 & 6 & -1 \\ 10 & 3 & 9 & 7 \\ 16 & 4 & 12 & 15 \end{array} \right]$$

$$r(A) \leq 4$$

$$R_4 - (R_1 + R_3)$$

$$R_3 - (R_1 + R_2)$$

$$\left[ \begin{array}{cccc} 6 & 1 & 3 & 8 \\ 4 & 2 & 6 & -1 \\ \hline 0 & 0 & 0 & 0 \\ \hline 0 & 0 & 0 & 0 \end{array} \right]$$

$$r(A) < 4$$

$$r(A) < 3$$

$$r(A) = 2$$

# The consistency & Inconsistency of the system of eq<sup>n</sup>:-

$$a_1x + b_1y + c_1z = d_1$$

$$a_2x + b_2y + c_2z = d_2$$

$$a_3x + b_3y + c_3z = d_3$$

$$A = \begin{bmatrix} a_1 & b_1 & c_1 \\ a_2 & b_2 & c_2 \\ a_3 & b_3 & c_3 \end{bmatrix}$$

Coefficient Matrix

$$B = \begin{bmatrix} d_1 \\ d_2 \\ d_3 \end{bmatrix}$$

Consto Matrix.

$$X = \begin{bmatrix} x \\ y \\ z \end{bmatrix}$$

Variable Matrix.

$$AB = \begin{bmatrix} a_1 & b_1 & c_1 & d_1 \\ a_2 & b_2 & c_2 & d_2 \\ a_3 & b_3 & c_3 & d_3 \end{bmatrix}$$

Augmented Matrix.

Eq<sup>n</sup> form  $\rightarrow$   $AX = B$

\* Any system is said to be a consistent system, if it has a solution.

\* Inconsistent system has no solution.

$\rightarrow$  If  $P(AB) = P(A) = \text{No. of Unknowns}$ , then the system is said to be consistent and having unique solution.

$\rightarrow$  If  $P(AB) = P(A) < \text{No. of Unknowns}$ , then the system is said to be consistent and having more than one solution, or  $\infty$  no. of solutions.

\* Remember  $\rightarrow$  either 1 sol<sup>n</sup> or  $\infty$  sol<sup>n</sup> exist for system  
To confuse, in exam option will be 2 sol<sup>n</sup> or 3 sol<sup>n</sup> ~~(X)~~

$\rightarrow$  If  $P(AB) \neq P(A)$ , then the system is said to be inconsistent and have no solution.

$$\begin{aligned}
 6. \quad & x + y + z = 3 \\
 & x + 2y + 3z = 4 \\
 & x + 4y + 9z = 6
 \end{aligned}$$

Ⓐ 0   Ⓑ 1   Ⓒ ~~2~~   Ⓓ ∞

NOTE!-

- If  $AB$  is rectangle matrix, then last column must be excluded.
- If  $AB$  is a square matrix, then last column must be included.
- If last column is excluded, then it represents both the matrix's rank.
- If last column included, then it represents only  $AB$  matrix rank.

Now,

$$[AB] = \left[ \begin{array}{ccc|c} 1 & 1 & 1 & 3 \\ 1 & 2 & 3 & 4 \\ 1 & 4 & 9 & 6 \end{array} \right]_{3 \times 4} \quad R_2 - R_1, R_3 - R_1$$

$$= \left[ \begin{array}{ccc|c} 1 & 1 & 1 & 3 \\ \hline 0 & 1 & 2 & 1 \\ 0 & 3 & 8 & 3 \end{array} \right]$$

→  $\neq 0 \Rightarrow P(AB) = P(A) = 3 = 3$  (no. of unknown)  
 $\therefore$  System has unique sol<sup>n</sup>.

Q. Check the consistency of system -

$$x - 2y + 3z = 2$$

$$2x - 3z = 3$$

$$x + y + z = 0$$

$$\rightarrow \left[ \begin{array}{ccc|c} 1 & -2 & 3 & 2 \\ 2 & 0 & -3 & 3 \\ 1 & 1 & 1 & 0 \end{array} \right]_{3 \times 4} \quad R_1 + 2R_3$$

$$\left[ \begin{array}{ccc|c} 3 & 0 & 5 & 2 \\ 2 & 0 & -3 & 3 \\ \hline 1 & 1 & 1 & 0 \end{array} \right]$$

$\neq 0 \Rightarrow \rho(AB) = \rho(A) = 3 = \text{no. of unknowns } \&$

$\therefore$  System having consistent and having unique sol<sup>n</sup>.

Q.  $3x + 3y + 2z = 1$

$$x + 2y = 4$$

$$10y + 3z = -2$$

$$2x - 3y - z = 5$$

$$[AB] = \left[ \begin{array}{ccc|c} 1 & 2 & 0 & 4 \\ 3 & 3 & 2 & 1 \\ 0 & 10 & 3 & -2 \\ 2 & -3 & -1 & 5 \end{array} \right]_{4 \times 4} \quad R_2 - 3R_1, R_4 - 2R_1$$

$$= \left[ \begin{array}{ccc|c} 1 & 2 & 0 & 4 \\ 0 & -3 & 2 & -11 \\ 0 & 10 & 3 & -2 \\ 0 & -7 & -1 & 3 \end{array} \right] \quad R_2 + 2R_4, R_3 + 3R_4$$

Note:-

If 0 present  $\neq$  make adjacent '0'

If no '0' present, then check if  $\pm$  or  $-1 \rightarrow$  make adjacent of  $1$  or  $-1 = 0$

$$\begin{bmatrix} 1 & 2 & 0 & 4 \\ 0 & -17 & 0 & -17 \\ 0 & -11 & 0 & -11 \\ 0 & -7 & -1 & -3 \end{bmatrix}$$

take common ←

take common ←

then  $R_3 - R_2$  and interchange  $R_4$  &  $R_3$

$$\left[ \begin{array}{ccc|c} 1 & 2 & 0 & 4 \\ 0 & 1 & 0 & 1 \\ 0 & -7 & -1 & -3 \\ \hline 0 & 0 & 0 & 0 \end{array} \right]$$

$\neq 0 \Rightarrow \rho(A) = \rho(AB) = 3 = \text{no. of unknowns}$

$\therefore$  System is consistent & having unique solution.

Q.

$$4x - 2y + 6z = 8$$

$$x + y - 3z = -1$$

$$15x - 3y + 9z = 21$$

$$[AB] = \begin{bmatrix} 1 & 1 & -3 & -1 \\ 2 & -1 & 3 & 4 \\ 5 & -1 & 3 & 7 \end{bmatrix}_{3 \times 4}$$

$R_2 + R_1$   
 $R_3 + R_1$

$$\left[ \begin{array}{ccc|c} 1 & 1 & -3 & -1 \\ \hline 3 & 0 & 0 & 3 \\ \hline 6 & 0 & 0 & 6 \end{array} \right]$$

$\therefore \rho(AB) = \rho(A) = 2 < 3$ .

System - Consistent  
But  $\infty$  Solution.

$\Rightarrow 0 \Rightarrow$

We can write system again as —

$$\begin{cases} x + y - 3z = -1 \\ 3x = 3 \\ \Rightarrow x = 1. \end{cases}$$

$$\begin{aligned} 1 + y - 3z &= -1 \\ \Rightarrow y - 3z &= -2 \quad \text{--- (1)} \end{aligned}$$

Since, we are discussing the consistency of only real system of equations, therefore solutions always real numbers.

Let  $z = k$  where  $k \in \mathbb{R}$

$$\begin{aligned} y - 3k &= -2 \\ \Rightarrow \begin{cases} y = 3k - 2 \\ z = k \\ x = 1. \end{cases} \end{aligned}$$

For different values of  $k$ , we will have diff. solutions and these solutions are infinite.

a.

$$\begin{aligned} 2x - y + z &= 4 \\ 3x - y + z &= 6 \\ 4x - y + 2z &= 7 \\ -x + y - z &= 9 \end{aligned}$$

check the consistency of system.

$$\rightarrow [AB] = \begin{bmatrix} -1 & 1 & -1 & 9 \\ 2 & -1 & 1 & 4 \\ 3 & -1 & 1 & 6 \\ 4 & -1 & 2 & 7 \end{bmatrix}_{4 \times 4}$$

$$\begin{aligned} R_2 + R_1 \\ R_3 + R_1 \\ R_4 + R_1 \end{aligned}$$

$$\equiv \begin{bmatrix} -1 & 1 & -1 & 9 \\ 1 & 0 & 0 & 13 \\ 2 & 0 & 0 & 15 \\ 3 & 0 & 1 & 16 \end{bmatrix}$$

$C_{12}$

$$\left[ \begin{array}{c|ccc} 1 & -1 & -1 & 9 \\ \hline 0 & 1 & 0 & 13 \\ 0 & 2 & 0 & 15 \\ 0 & 3 & 1 & 16 \end{array} \right]$$

$\rightarrow \neq 0 \Rightarrow \rho(AB) = 4$

But  $\rho(A) = 3$

$\therefore \rho(AB) \neq \rho(A)$

Hence, System is inconsistent and having no solution.

Q. for what values of  $\lambda$  and  $\mu$ , does the system of equations

$$x + y + z = 6$$

$$x + 2y + 3z = 10$$

$$x + 2y + \lambda z = \mu$$

has —

- (A) no solution
- (B) Unique sol<sup>n</sup>
- (C) More than one solution

$\rightarrow$

$$[AB] = \left[ \begin{array}{ccc|c} 1 & 1 & 1 & 6 \\ 1 & 2 & 3 & 10 \\ 1 & 2 & \lambda & \mu \end{array} \right]_{3 \times 4}$$

$R_3 - R_2$

$$\left[ \begin{array}{ccc|c} 1 & 1 & 1 & 6 \\ 1 & 2 & 3 & 10 \\ \hline 0 & 0 & \lambda - 3 & \mu - 10 \end{array} \right]$$

$\rightarrow \neq 0 \Rightarrow \rho(AB) = \rho(A) = 3$  if  $\lambda - 3 \neq 0$   
 $< 3$  if  $\lambda - 3 = 0$

Case - I

Case - II

Case - III

Q.

$$2x + 3y + 5z = 9$$

$$7x + 3y - 2z = 8$$

$$2x + 3y + \lambda z = 4$$

}

for what values of  $\lambda$  &  $\mu$  system has —

(A) no sol<sup>n</sup>

(B) unique sol<sup>n</sup>

(C) more than one sol<sup>n</sup>

## # Eigen Values and Eigen Vectors :-

# Characteristic equation :- Let  $A$  be the sq. matrix of order  $n \times n$  and  $I$  be the unit matrix of order  $n \times n$  then  $|A - \lambda I| = 0$  is called the characteristic eq<sup>n</sup> where  $\lambda$  is a parameter.

The roots of characteristic eq<sup>n</sup> are called characteristic roots / latent / eigen / proper values.

$X = \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix}$  or  $[x_1 \ x_2 \ \dots \ x_n]^T$  which satisfies the matrix eq<sup>n</sup>  $[A - \lambda I]X = 0$  is called the corresponding eigen vector of the matrix.

Note:-

i) The sum of eigen values of any matrix is equal to sum of the elements of its principal diagonals.

Ex:-

$$\begin{bmatrix} 1 & 1 & 3 \\ 1 & 5 & 1 \\ 3 & 1 & 1 \end{bmatrix}$$

Ⓐ 1   Ⓑ 3   Ⓒ 5   Ⓓ 7

$$\text{Trace of } A = 1 + 5 + 1 = 7$$

ii) The product of eigen values of any matrix is equal to its determinant.

Ex:- The characteristic roots of matrix  $\begin{bmatrix} 5 & 4 \\ 1 & 2 \end{bmatrix}$

Ⓐ 2, 5, 4

Ⓑ 3, 4

Ⓒ 1, 6

Ⓓ 2, 3

$\left. \begin{array}{l} \rightarrow \text{Sum} = \text{sum of principle diagonal} \\ \rightarrow 6 + 1 = 7 \\ \rightarrow \text{Product} = \text{determinant value} \end{array} \right\}$

iii) The eigen values of a symmetric matrix ( $A^T = A$ ) are purely real.

Ex:-  $A = \begin{bmatrix} 3 & 2 \\ 2 & 3 \end{bmatrix} = A^T$

∴ Eigen values are 1 & 5 which are purely real

Cross check  $\rightarrow 1 + 5 = 6$

$1 \times 5 = 5$  Equal to determinant value

equal to sum of principle diagonal

iv) The eigen value of skew symmetric matrix are ( $A^T = -A$ ) either purely imaginary or zeroes

Ex:-

$$A = \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix}$$

Eigen Values?

Ⓐ 1, -1  
X

Ⓑ i, -i  
✓

Ⓒ -1, 0  
X

Ⓓ 0, 1  
X

$$A^T = \begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix} = - \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix} = -A$$

v) If a matrix is either lower triangular or upper triangular, then principle diagonal elements are called eigen values.

\* If Given Matrix only is L.T.M or U.T.M (without operation)

Ex:-

$$\begin{bmatrix} 1 & 2 & 3 \\ 0 & -4 & 2 \\ 0 & 0 & 7 \end{bmatrix}$$

$$\begin{bmatrix} 1 & 0 & 0 \\ 2 & 3 & 0 \\ 4 & 5 & 6 \end{bmatrix}$$

only original

↓  
Then only follow this rule.

$$\lambda = 1, -4, 7$$

$$\lambda = 1, 3, 6$$

Don't Use this Property to find eigen value when L.T.M or U.T.M is made by some operation.

vii) If  $\lambda$  is an eigen value of A

then  $\lambda^2$  \_\_\_\_\_  $A^2$

$\lambda^3$  \_\_\_\_\_  $A^3$

$\lambda^n$  \_\_\_\_\_  $A^n$

$\frac{1}{\lambda}$  \_\_\_\_\_  $A^{-1}$

Ex:- if  $B = \begin{bmatrix} 3 & 2 \\ 2 & 3 \end{bmatrix}$  and  $\lambda = 1, 5$

then for  $B^2$  what will be eigen value → After getting result  $\lambda$ , verify it by

sum & product rule.

$$A = \begin{bmatrix} 1 & 2 \\ 2 & 1 \end{bmatrix}$$

then eigen values of  $A^3$  are -

a) 5, 4

b) 9, 1

c) 16, -1

d) 27, -1

Q. Find the eigen values and corresponding eigen vectors of

$$\text{Matrix } A = \begin{bmatrix} 5 & 4 \\ 1 & 2 \end{bmatrix}$$

$$\rightarrow |A - \lambda I| = 0$$

$$\left| \begin{bmatrix} 5 & 4 \\ 1 & 2 \end{bmatrix} - \begin{bmatrix} \lambda & 0 \\ 0 & \lambda \end{bmatrix} \right| = 0$$

$$\begin{vmatrix} 5-\lambda & 4 \\ 1 & 2-\lambda \end{vmatrix} = 0$$

$$\Rightarrow 10 - 7\lambda + \lambda^2 - 4 = 0$$

$$\Rightarrow \lambda^2 - 7\lambda + 6 = 0$$

$$\Rightarrow (\lambda - 1)(\lambda - 6) = 0$$

$$\lambda = 1, 6$$

Eigen Vector —

$$[A - \lambda I] X = 0$$

Subtract  $\lambda$  from matrix and multiply ~~vector~~  $X$  Matrix

$$\text{for } \lambda = 1 \quad \begin{bmatrix} 4 & 4 \\ 1 & 1 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = 0$$

$$\Rightarrow 4x_1 + 4x_2 = 0$$

$$x_1 + x_2 = 0$$

$$\Rightarrow x_1 = -x_2$$

$$\Rightarrow \frac{x_1}{1} = \frac{x_2}{-1}$$

$$\text{eigen vector} = \begin{bmatrix} 1 \\ -1 \end{bmatrix} \text{ or } \begin{bmatrix} 1 & -1 \end{bmatrix}^T$$

for  $\lambda = 6$  —

$$\begin{bmatrix} -1 & 4 \\ 1 & -4 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = 0$$

$$-x_1 + 4x_2 = 0$$

$$\frac{x_1}{4} = \frac{x_2}{1}$$

$$\text{eigen vector} = \begin{bmatrix} 4 \\ 1 \end{bmatrix} \text{ or } \begin{bmatrix} 4 & 1 \end{bmatrix}^T$$

Q. find the ch. roots of matrix  
or Eigen Value.

$$\begin{bmatrix} 8 & -6 & 2 \\ -6 & 7 & -4 \\ 2 & -4 & 3 \end{bmatrix}$$

1st Step

- (a) 3, 7, 8 ✓  
 (b) 2, 2, 14 ✓  
 (c) 0, 3, 15 ✓  
 (d) 1, 4, 9 ✗
- By sum of diagonal rate = sum of eigen value.

2nd step

check determinant.

$$\Delta = 0.$$

So, one root must be 0 so that to make product rule of eigen values correct.

Q. find the latent root of matrix -

$$\begin{bmatrix} 1 & 0 & -1 \\ 1 & 2 & 1 \\ 2 & 2 & 3 \end{bmatrix}$$

- (a) 1, 2, 3 ✓ ✓  
 (b) 0, 2, 4 ✓ ✗  
 (c) 1, 1, 4 ✓ ✗  
 (d) 2, 2, 2 ✓ ✗

2nd step

$$4 + 2 = 6.$$

Q. find Eigen Values and Corresponding eigen vectors of

$$\begin{bmatrix} 1 & 2 & 3 \\ 0 & -4 & 2 \\ 0 & 0 & 7 \end{bmatrix}$$

$$\lambda = 1, -4, 7 \quad \left\{ \because \text{It is U.T.M} \right\}$$

for  $\lambda = 1$

$$\begin{bmatrix} 0 & 2 & 3 \\ 0 & -5 & 2 \\ 0 & 0 & 6 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = 0.$$

$$2x_2 + 3x_3 = 0.$$

$$-5x_2 + 2x_3 = 0.$$

$$6x_3 = 0 \Rightarrow x_3 = 0 \Rightarrow x_2 = 0 \rightarrow \text{not able to identify any info abt } x_1.$$

Remember

\* for any eigen values if  $\lambda$  eigen vectors = 0 then third must be only non zero.

$$\therefore \text{eigen vector set} = \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}$$

for  $\lambda = -4$

$$\begin{bmatrix} 5 & 2 & 3 \\ 0 & 0 & 2 \\ 0 & 0 & 11 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = 0.$$

$$5x_1 + 2x_2 + 3x_3 = 0. \Rightarrow 5x_1 + 2x_2 = 0.$$

$$2x_3 = 0 \Rightarrow x_3 = 0 \Rightarrow 5x_1 = -2x_2$$

$$11x_3 = 0. \Rightarrow x_3 = 0. \Rightarrow \frac{x_1}{2} = \frac{x_2}{-5}$$

Eigen Vector:  $\begin{bmatrix} 2 \\ -5 \\ 0 \end{bmatrix}$

for  $\lambda = 7$

$$\begin{bmatrix} -6 & 2 & 3 \\ 0 & -11 & 2 \\ 0 & 0 & 0 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = 0.$$

$$-6x_1 + 2x_2 + 3x_3 = 0.$$

$$-11x_2 + 2x_3 = 0$$

$$\Rightarrow \frac{x_2}{2} = \frac{x_3}{11}$$

$$\Rightarrow 6x_1 = 3x_3 \Rightarrow x_1 = \frac{3x_3}{6}$$

eigen vector:  $\begin{bmatrix} \frac{37}{6} \\ 2 \\ 11 \end{bmatrix} \propto \begin{bmatrix} 37 \\ 12 \\ 66 \end{bmatrix}$

Q. for the matrix  $P = \begin{bmatrix} 9 & -2 & 2 \\ 0 & -2 & 1 \\ 0 & 0 & 1 \end{bmatrix}$  one of the eigen value is

-2 . which of the following is an eigen vector .

- (A)  $\begin{bmatrix} 3 \\ -2 \\ 1 \end{bmatrix}$     (B)  $\begin{bmatrix} -3 \\ 2 \\ -1 \end{bmatrix}$     (C)  $\begin{bmatrix} 1 \\ -2 \\ 3 \end{bmatrix}$     (D)  $\begin{bmatrix} 2 \\ 5 \\ 0 \end{bmatrix}$

→ for  $\lambda = -2$   $\begin{bmatrix} 5 & -2 & 2 \\ 0 & 0 & 1 \\ 0 & 0 & 3 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = 0$

$5x_1 - 2x_2 + 2x_3 = 0 \Rightarrow 5x_1 - 2x_2 = 0$

$\frac{x_1}{2} = \frac{x_2}{5}$

$\left. \begin{matrix} x_3 = 0 \\ 3x_3 = 0 \end{matrix} \right\} x_3 = 0$

eigen vector =  $\begin{bmatrix} 2 \\ 5 \\ 0 \end{bmatrix}$

Q. for the matrix  $P = \begin{bmatrix} 1 & 1 & 0 \\ 0 & 2 & 2 \\ 0 & 0 & 3 \end{bmatrix}$  which one of the following is an eigen vector .

- a)  $[1 \ -1 \ 1]^T$   
 b)  $[1 \ 2 \ 1]^T$   
 c)  $[2 \ -1 \ 1]^T$   
 d)  $[1 \ 1 \ -1]^T$

Give 1st pref to least eigen value .

If option not found check only by subtracting  $-\lambda$  in 1st row check eigen value .

Again 2nd pref to 2nd least eigen value .  
 Again if option not found .  
 same subtract  $\lambda$  from 2nd row check eigen value .

Again 3rd pref to 3rd least eigen value .  
 but do not subtract to 3rd row subtract from either 1st or 2nd row (no effect) .  
 check eigen value .

## # Cayley - Hamilton Theorem :-

Every sq. matrix satisfies its own characteristic eqn.

Ex:-  $A = \begin{bmatrix} 5 & 4 \\ 1 & 2 \end{bmatrix}$

$$\begin{vmatrix} 5-\lambda & 4 \\ 1 & 2-\lambda \end{vmatrix} = 0.$$

$$\Rightarrow 10 - 7\lambda + \lambda^2 - 4 = 0$$

$$\Rightarrow \lambda^2 - 7\lambda + 6 = 0$$

$$\Rightarrow \lambda = \dots$$

Means —

$$A^2 - 7A + 6I = 0 \rightarrow \text{A/c to Cayley Hamilton Theorem}$$

$$\Rightarrow 6I = 7A - A^2$$

$$\Rightarrow 6IA^{-1} = 7AA^{-1} - A^2A^{-1}$$

$$\Rightarrow 6A^{-1} = 7I - A$$

$$= \begin{bmatrix} 7 & 0 \\ 0 & 7 \end{bmatrix} - \begin{bmatrix} 5 & 4 \\ 1 & 2 \end{bmatrix}$$

$$= \frac{1}{6} \begin{bmatrix} 2 & -4 \\ -1 & 5 \end{bmatrix}$$

↓  
For Inverse finding  
without knowing  
adj of A & determinant.

Q. Let M be 3x3 matrix with ch. eqn  $x^3 + 1 = 0$  then the inverse of M = —

→ A/c to Cayley Hamilton theorem —

$$M^3 + I = 0$$

$$I = -M^3$$

$$IM^{-1} = -M^3M^{-1}$$

$$\boxed{M^{-1} = -M^2}$$

Q. Let P be a 3x3 matrix with ch. eq<sup>n</sup>  $\lambda^3 + \lambda^2 + 2\lambda + 1 = 0$   
 then inverse of P = —

- (A)  $P^2 + P + I$   
 (B)  $P^2 + P + 2I$   
 (C)  $-(P^2 + P + 2I)$   
 (D)  $-(P^2 + P + I)$

→  $P^3 + P^2 + 2P + I = 0$

⇒  $I = -(P^3 + P^2 + 2P)$

⇒  $P^{-1} = -(P^2 + P + 2I)$

Q. find the value of  $\lambda$  for which the following system of eq<sup>s</sup>

~~$\lambda x + 3y + 5z = 0$~~   
 $\lambda x + 3y + 5z = 0$   
 $2x - 4\lambda y + \lambda z = 0$   
 $-4x + 18y + 7z = 0$

Note:-

$AX = 0$  is called Homogeneous.

$X = 0$  will always be 0 solution.

$\begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix}$

↓  
 This is called Trivial equation.

Note:-

for the system of Eq<sup>n</sup>  $AX = 0$

$X = 0$  will always a solution

and this sol<sup>n</sup> is called a trivial sol<sup>n</sup>.

for a non trivial solution determinant of coefficient matrix = 0 (always)

~~$\begin{bmatrix} \lambda & 3 & 5 & \lambda & 3 \\ 2 & -4\lambda & \lambda & 2 & -4\lambda \\ -4 & 18 & 7 & -4 & 18 \end{bmatrix}$~~

$-28\lambda^2 - 12\lambda + 180 - 80\lambda - 18\lambda^2 - 42 = 0$

⇒  $-46\lambda^2 - 92\lambda + 138 = 0$

⇒  $\lambda^2 + 2\lambda - 3 = 0$

$\lambda = -3$  or  $1$ .

Q. Rank of  $\begin{bmatrix} 4 & 2 & 1 & 3 \\ 6 & 3 & 4 & 7 \\ 2 & 1 & 0 & 1 \end{bmatrix}$

(A) 1 (B) 2 (C) 3 (D) 4

→  $\begin{bmatrix} 4 & 2 & 1 & 3 \\ 6 & 3 & 4 & 7 \\ 2 & 1 & 0 & 1 \end{bmatrix}$   
 $3 \times 4$

$\rho(A) \leq 3$

$R_2 - 4R_1$

$\begin{bmatrix} 4 & 2 & | & 1 & 3 \\ -10 & -5 & | & 0 & -5 \\ 2 & 1 & | & 0 & 1 \end{bmatrix}$

$\rightarrow = 0 \Rightarrow \rho(A) < 3$

$\rho(A) = 2$

ADD PHOTO AT & BOOK CENTRE  
 ADD NEAR MAE EASY LANE NO. 4  
 WESTEND MANS, SAIDULAJAB  
 NEW DELHI-110030  
 MOB: 9560127702

Q. Rank and nullity of matrix

$\begin{bmatrix} 6 & 1 & 8 & 3 \\ 2 & 3 & 0 & 2 \\ 4 & -1 & -8 & -3 \end{bmatrix}$

\*\*\*

Nullity of A = no. of columns - Rank

$\begin{bmatrix} 10 & 0 & | & 0 & 0 \\ 2 & 3 & | & 0 & 2 \\ 4 & -1 & | & -8 & -3 \end{bmatrix}$

$\rho(A) \leq 3$

$\rightarrow \neq 0 \Rightarrow \rho(A) = 3$

$\therefore$  nullity of A =  $4 - 3$

= 1 Ans



Q. The eigen vector pair of matrix  $\begin{bmatrix} 3 & 4 \\ 4 & -3 \end{bmatrix}$

(a)  $\begin{bmatrix} 2 \\ 1 \end{bmatrix} \begin{bmatrix} 1 \\ 2 \end{bmatrix}$

(b)  $\begin{bmatrix} 2 \\ 1 \end{bmatrix} \begin{bmatrix} 1 \\ -2 \end{bmatrix}$

(c)  $\begin{bmatrix} -2 \\ 1 \end{bmatrix} \begin{bmatrix} 1 \\ 2 \end{bmatrix}$

(d)  $\begin{bmatrix} -2 \\ 1 \end{bmatrix} \begin{bmatrix} 1 \\ -2 \end{bmatrix}$

$\rightarrow \begin{bmatrix} 3 & 4 \\ 4 & -3 \end{bmatrix} - \lambda \Rightarrow -9 - 3\lambda + 3\lambda + \lambda^2 - 16 = 0$

$\Rightarrow \lambda^2 - 25 = 0$

$\lambda = 5, -5$

for  $\lambda = 5$

$\begin{bmatrix} -2 & 4 \\ 4 & -8 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = 0$

$-2x_1 + 4x_2 = 0$

$4x_1 - 8x_2 = 0$

$\frac{x_1}{2} = \frac{x_2}{1}$

$\begin{bmatrix} 2 \\ 1 \end{bmatrix}$

for  $\lambda = -5$

$\begin{bmatrix} 8 & 4 \\ 4 & 2 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = 0$

$8x_1 + 4x_2 = 0$

$\frac{x_1}{1} = \frac{x_2}{-2}$

$\begin{bmatrix} 1 \\ -2 \end{bmatrix}$

Q. How many of the following matrices have an eigen value = 1

$\begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}$      $\begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix}$      $\begin{bmatrix} 1 & -1 \\ 1 & 1 \end{bmatrix}$      $\begin{bmatrix} -1 & 0 \\ 1 & -1 \end{bmatrix}$   
 $\lambda = 1, 0$      $\lambda = 0, 0$      $(1-\lambda)^2 + 1 = 0$      $(-1-\lambda)^2 = 0$   
 (A) 1    (B) 2    (C) 3    (D) 4

Q. If the following represents eq<sup>n</sup> of line then line passes through the point

$$\begin{bmatrix} x & 2 & 4 \\ y & 8 & 0 \\ 1 & 1 & 1 \end{bmatrix} = 0$$

- a) (0, 0)
- b) (3, 4)
- c) (4, 3)
- d) (4, 4)

→

~~$$\begin{bmatrix} x & 2 & 4 & x & 2 \\ y & 8 & 0 & y & 8 \\ 1 & 1 & 1 & 1 & 0 \end{bmatrix}$$~~

$$8x + 4y - 32 - 2y = 0$$

$$\Rightarrow 8x + 2y = 32$$

~~$$\begin{aligned} -32 + 0 + 2y - 8x - 0 + 0 &= 0 \\ \Rightarrow -8x + 2y + 32 &= 0 \\ \Rightarrow 8x + 2y &= 32 \end{aligned}$$~~

- Q. A is  $3 \times 4$  real matrix and  $Ax = B$  is an inconsistent system of eq<sup>s</sup> then the highest possible rank of A —  
 @ 1  2  3  4 .

$$A = 3 \times 4$$

$$[AB] = 3 \times 5 \Rightarrow \rho(A) \leq 3$$

for inconsistent system  $\rho(A) \neq \rho(AB)$   $\left\{ \begin{array}{l} \because \rho(A) \text{ can't be} \\ \text{greater than} \\ \rho(AB) \end{array} \right.$

$$\therefore \rho(A) < 3$$

$$\rho(A) = 2$$

- Q. Given an orthogonal matrix  $\begin{bmatrix} 1 & 1 & 1 & 1 \\ 1 & 1 & -1 & -1 \\ 1 & -1 & 0 & 0 \\ 0 & 0 & 1 & -1 \end{bmatrix}$  then  $(AA^T)^{-1} = \underline{\underline{I}}$

NOTE:-

A matrix is said to be orthogonal matrix if  $AA^T = I$ .

Q.  $A = \begin{bmatrix} 2 & -0.1 \\ 0 & 3 \end{bmatrix}$  .  $A^{-1} = \begin{bmatrix} \frac{1}{2} & a \\ 0 & b \end{bmatrix}$

then  $a+b = \underline{\hspace{2cm}}$

$$A^{-1} = \frac{1}{6} \begin{bmatrix} 3 & 0.1 \\ 0 & 2 \end{bmatrix}$$

$$\therefore a = \frac{0.1}{6}$$

$$b = \frac{2}{6}$$

$$\therefore a+b = \frac{0.1}{6} = \frac{21}{60} = \frac{7}{20} \underline{\underline{A}} \underline{\underline{B}}$$

Q. Let  $A = \begin{bmatrix} 2 & 0 & 0 \\ -1 & -2 & 0 \\ 4 & 2 & -1 \end{bmatrix}$  and  $B = 2A^2$  then  $|B| =$  \_\_\_\_\_

- (A) 16    (B) 32    (C) 64    (D) 128

$$B = 2A^2$$

$$|B| = |2A^2| \rightarrow \text{order} = \underline{\underline{3 \times 3}}$$
$$= 2^3 |A^2|$$


$$= 8(16)$$

$$= 128$$

# DIFFERENTIAL CALCULUS

Limit:- A no  $l$  is said to be the limit of the fn  $f(x)$  as  $x \rightarrow a$  if  $\forall \epsilon > 0$  (however small)  $\exists$  a  $\delta > 0$  such that  $|f(x) - l| < \epsilon$   
 $\forall |x - a| < \delta$

$$\lim_{x \rightarrow a} f(x) = l$$



$$\therefore l - \epsilon < f(x) < l + \epsilon$$
$$\forall a - \delta < x < a + \delta$$

\* Left limit:-  $\lim_{x \rightarrow a^-} f(x) = \lim_{h \rightarrow 0} f(a-h), (a-\delta < x < a)$

\* Right limit:-  $\lim_{x \rightarrow a^+} f(x) = \lim_{h \rightarrow 0} f(a+h), (a < x < a+\delta)$

\* Existence of Limit:- The limit of a function exists if both the left and right limits are existed and are equal.

ex:-  $\lim_{x \rightarrow a} \frac{1}{x-a}$

$$\therefore \lim_{x \rightarrow a^-} f(x) = \lim_{h \rightarrow 0} f(a-h) = \frac{1}{a-h-a} = -\frac{1}{h}$$

$$\lim_{x \rightarrow a^+} f(x) = \lim_{h \rightarrow 0} f(a+h) = \frac{1}{a+h-a} = \frac{1}{h}$$

$\therefore$  LH limit  $\neq$  RH limit

$\therefore$  Limit not exists

# # Formulae:-

$$\lim_{x \rightarrow a} \frac{x^n - a^n}{x - a} = n a^{n-1}$$

$$\lim_{x \rightarrow a} \frac{x^m - a^m}{x^n - a^n} = \frac{m}{n} a^{m-n}$$

$$\lim_{x \rightarrow 0} \frac{e^x - 1}{x} = 1$$

$$\lim_{x \rightarrow 0} \frac{e^{mx} - 1}{x} = m$$

$$\lim_{x \rightarrow 0} \frac{a^x - 1}{x} = \log a$$

$$\lim_{x \rightarrow 0} \frac{a^{bx} - b^x}{x} = \log\left(\frac{a}{b}\right)$$

$$\bullet \lim_{x \rightarrow 0} (1+x)^{1/x} = e$$

$$\bullet \lim_{x \rightarrow 0} (1+ax)^{1/x} = e^a$$

$$\lim_{x \rightarrow 0} (1+ax)^{b/x} = e^{ab}$$

$$\dots \lim_{x \rightarrow 0} (1+ax)^{1/bx} = e^{a/b}$$

$$\lim_{x \rightarrow \infty} \left(1 + \frac{1}{x}\right)^x = e$$

$$\lim_{x \rightarrow \infty} \left(1 - \frac{1}{x}\right)^x = e^{-1}$$

$$\lim_{x \rightarrow 0} \frac{\sin x}{x} = 1$$

$$\lim_{x \rightarrow 0} \frac{\tan x}{x} = 1$$

$$\lim_{x \rightarrow 0} \frac{\sin^{-1} x}{x} = 1$$

$$\lim_{x \rightarrow 0} \frac{\tan^{-1} x}{x} = 1$$

$$\lim_{x \rightarrow 0} \frac{1 - \cos mx}{1 - \cos nx} = \frac{m^2}{n^2}$$

$$\lim_{x \rightarrow 0} \frac{\cos ax - \cos bx}{x^2} = \frac{b^2 - a^2}{2}$$

$\frac{0}{0}, \frac{\infty}{\infty}, \frac{\infty}{0}, 0 \times \infty, 1^\infty$  etc are indeterminate forms.

If  $\frac{0}{0}, \frac{\infty}{\infty}$  then apply

L.H. rule

$$\lim_{x \rightarrow a} \frac{f(x)}{g(x)} = \lim_{x \rightarrow a} \frac{f'(x)}{g'(x)}$$

$$= \lim_{x \rightarrow a} \frac{f''(x)}{g''(x)} = \dots$$

$$\lim_{x \rightarrow a} [f(x)]^{g(x)} = e^{\lim_{x \rightarrow a} g(x)[f(x)-1]}$$

$$\lim_{x \rightarrow \infty} \frac{f(x)}{g(x)}, \text{ if } f(x) \text{ \& } g(x) \text{ are algebraic fn.}$$

Case-1 if degree of  $f(x) > g(x)$  degree then result =  $\infty$

ex:-  $\lim_{x \rightarrow \infty} \frac{2x^3 + 3x + 5}{5x^2 + 6x + 7} = \infty$

Case-2 if degree of  $f(x) < g(x)$  degree then result = 0

ex:-  $\lim_{x \rightarrow \infty} \frac{2x^2 + 3x + 4}{x^3 + 8} = 0$

Case-3 If  $f(x)$  and  $g(x)$  have equal degrees then -

result =  $\frac{\text{co. eff of } N^r}{\text{co. eff of } D^r}$

ex:-  $\lim_{x \rightarrow \infty} \frac{2x^2 + 3x + 4}{5x^2 + 5x + 8} = \frac{2}{5}$

\*  $|x| = x$  if  $x > 0$

\*  $|x| = -x$  if  $x < 0$

\*  $[x]$  = greatest integer not greater than  $x$ .

\* Examples :-

①  $\lim_{x \rightarrow 0} \frac{1 - \cos x}{x^2} = \left(\frac{1}{2}\right)$

②  $\lim_{x \rightarrow 0} \frac{1 - \cos x}{x \sin x} = \frac{1 - \cos x}{x^2} = \frac{\sin x}{2x} = \frac{\cos x}{2} = \left(\frac{1}{2}\right)$

In Trigonometric  
fn, avoid  
LH rule  
Try alternatives

③  $\lim_{x \rightarrow 0} \frac{\tan x - \sin x}{\sin x \cdot \sin^3 x} = \frac{1 - \cos x}{\cos x (1 + \cos x)} = \frac{1}{1 \cdot (1+1)} = \left(\frac{1}{2}\right)$

④  $\lim_{x \rightarrow 1} \frac{\cos\left(\frac{\pi x}{2}\right)}{1 - \sqrt{x}} = \frac{f'(x) \sin\left(\frac{\pi x}{2}\right)}{f'\left(\frac{1}{2\sqrt{x}}\right)} = \frac{\pi/2}{1/2} = \left(\pi\right)$

⑤  $\lim_{x \rightarrow 0} \frac{\frac{2}{8} \cos x \left( \sin^8\left(\frac{\pi}{6} + x\right) - \sin^8\frac{\pi}{6} \right)}{x}$

$= \frac{2}{8} \left[ \frac{8 \sin^7\left(\frac{\pi}{6} + x\right) \cos\left(\frac{\pi}{6} + x\right) - 0}{1} \right]$

$= \frac{2}{8} \left[ 8 \times \frac{1}{2} \cdot \frac{\sqrt{3}}{2} \right]$

$= \left(\sqrt{3}\right)$  Ans

⑥  $\lim_{x \rightarrow 0} \frac{\sqrt{1-x^2} - \sqrt{1+x^2}}{x^2} = -1$

⑦  $\lim_{x \rightarrow 0} \frac{\sqrt[3]{1+x} - \sqrt[3]{1-x}}{x} = \frac{2}{3}$

⑧  $\lim_{x \rightarrow 0} \frac{\log(1+x)}{x} = 1$

$$(9) \quad \lim_{x \rightarrow 0} (1 + \sin x)^{\operatorname{cosec} x} = e^{\lim_{x \rightarrow 0} \operatorname{cosec} x [1 + \sin x - 1]} = 1$$

$$(10) \quad \lim_{x \rightarrow 0} (1 + 2x)^{1/3x} = e^{\lim_{x \rightarrow 0} \frac{1}{3x} [1 + 2x - 1]} = e^{\frac{2}{3}}$$

$$(11) \quad \lim_{n \rightarrow \infty} \left(1 - \frac{1}{n}\right)^{2n} = e^{\lim_{n \rightarrow \infty} 2n \left[1 - \frac{1}{n} - 1\right]} = e^{-2}$$

$$(12) \quad \lim_{x \rightarrow \infty} \left(\frac{x-1}{x-2}\right)^x = e^{\lim_{x \rightarrow \infty} x \left[\frac{x-1-x+1}{x-2}\right]} = e$$

$$(13) \quad \lim_{n \rightarrow \infty} \frac{1 + 2 + 3 + \dots + n}{n^2} = \frac{n(n+1)}{2n^2} = \frac{1}{2}$$

$$(14) \quad \lim_{n \rightarrow \infty} \frac{1 + 4 + 9 + \dots + n^2}{n^3} = \frac{n(n+1)(2n+1)}{6n^3} = \frac{2n^2 + 3n + 1}{6n^2} = \frac{2}{6} = \frac{1}{3}$$

$$(15) \quad \lim_{n \rightarrow \infty} \frac{n [1^3 + 2^3 + \dots + n^3]^2}{(1^2 + 2^2 + \dots + n^2)^3} = \frac{n \left[ \frac{n^2(n+1)}{4} \right]^2 \times 6^3}{\left[ \frac{n^3(n+1)}{6} \right]^3} = \frac{6^3 \frac{n^2(n+1)}{4^2}}{\frac{n^3(n+1)}{4^2 \times 8}} = \frac{6^3}{4^2 \times 8} = \frac{27}{16}$$

Trick for Partial fraction

for two terms

अस) भाग - दोन भाग

$$\lim_{n \rightarrow \infty} \left[ \frac{1}{1 \cdot 2} + \frac{1}{2 \cdot 3} + \dots + \frac{1}{n(n+1)} \right] = \sum \frac{1}{n(n+1)} = \sum \frac{1}{n} - \frac{1}{n+1} = 1 - \frac{1}{2} + \frac{1}{2} - \frac{1}{3} + \frac{1}{3} - \frac{1}{4} + \dots - \frac{1}{n+1} = 1$$

$$(17) \quad \lim_{x \rightarrow 0} \frac{1 - \cos 2x}{1 - \cos 3x} = \frac{2 \sin 2x}{3 \sin 3x} = \frac{4 \cos 2x}{9 \cos 3x} = \frac{4}{9}$$

$$(18) \quad \lim_{x \rightarrow 0} \frac{\cos 2x - \cos 3x}{x^2} = \frac{-2 \sin 2x + 3 \sin 3x}{2x} = \frac{-4 \cos 2x + 9 \cos 3x}{2} = \frac{5}{2}$$

## # Continuity of function:-

A function  $f$  is said to be continuous at  $x=a$ , if  $\lim_{x \rightarrow a} f(x) = f(a)$ , otherwise it is said to be a discontinuous function.

### \* Types of Discontinuous fn:-

#### i) Discontinuity of 1<sup>st</sup> Type (Jumped Discontinuity)

A fn.  $f$  is said to have discontinuity of 1<sup>st</sup> type if

$$\lim_{x \rightarrow a^-} f(x) \neq \lim_{x \rightarrow a^+} f(x)$$

#### ii) Discontinuity of 2<sup>nd</sup> Type

A fn.  $f$  is said to have discontinuity of 2<sup>nd</sup> type if either the left limit or the right limit are both does not exist.

#### iii) Discontinuity of 3<sup>rd</sup> Type (Removal Discontinuity)

A fn.  $f$  is said to have discontinuity of 3<sup>rd</sup> type

$$\lim_{x \rightarrow a^-} f(x) = \lim_{x \rightarrow a^+} f(x) \neq f(a)$$

### \* Certain standard functions:-

#### \* Standard Continuous function:-

①  $x^n$  is a continuous fn  $\forall x$  when  $n > 0$   
( $x, x^2, x^3, \dots, x^{7/3}, x^{3/2}, \dots$ )

②  $x^n$  is a continuous fn  $\forall x$  except  $x=0$  when  $n < 0$   
( $x^{-2} = \frac{1}{x^2}, x^{-3} = \frac{1}{x^3}, \dots$ )

③  $|x|$  is a continuous fn  $\forall x$ .

iv)  $\log x$  is a continuous fn  $\forall x > 0$ .

v) Every exponential function ( $e^x$  or  $a^x$ ) is continuous.

vi)  $\sin x$  and  $\cos x$  are continuous fn  $\forall x$

vii)  $\tan x$  and  $\sec x$  are continuous fn  $\forall x$  except  $x = (2n+1)\frac{\pi}{2}$

viii)  $\cot x$ ,  $\operatorname{cosec} x$  are continuous fn  $\forall x$  except  $x = n\pi$

Q.  $f(x) = \frac{1}{1+2^{1/x}}$  at  $x=0$

$$\lim_{h \rightarrow 0^-} f(0-h) = \frac{1}{1+2^{\frac{1}{0-h}}} = 1$$

$$\lim_{h \rightarrow 0^+} f(0+h) = \frac{1}{1+2^{\frac{1}{0+h}}} = 0$$

$\therefore$  L.H limit  $\neq$  R.H limit

$\therefore$  discontinuous at  $x=0$ .

Q  $f(x) = \frac{x(e^{1/x} - 1)}{e^{1/x} + 1}$ ,  $x \neq 0$

$$= 0, \quad x = 0.$$

$$\begin{aligned} \lim_{h \rightarrow 0^-} f(0-h) &= \frac{-h(e^{-\frac{1}{h}} - 1)}{e^{-\frac{1}{h}} + 1} \\ &= \frac{-hx - 1}{1} = 0 \end{aligned}$$

$$\lim_{h \rightarrow 0^+} f(0+h) = \frac{h(e^{\frac{1}{h}} - 1)}{e^{\frac{1}{h}} + 1} = \frac{h(1 - \frac{1}{e^{1/h}})}{(1 + \frac{1}{e^{1/h}})} = h = 0$$

$$f(0) = 0$$

$\therefore$  L.H limit = R.H limit =  $f(0) = 0$   $\therefore$  Continuous at  $x=0$

Q. A fn  $f(x)$  is defined by —

$$f(x) = \begin{cases} 0, & x \leq 0 \\ \frac{1}{2} - x, & 0 < x < \frac{1}{2} \\ \frac{1}{2}, & x = \frac{1}{2} \\ \frac{3}{2} - x, & \frac{1}{2} < x < 1 \\ 1, & x \geq 1 \end{cases}$$

Which of the following is true —

- (a)  $f$  is continuous at  $x=0$  . . . x
- (b)  $f$  is discontinuous at  $x=\frac{1}{2}$  . . . v
- (c) Continuous at  $x=1$  . . . x
- (d) All are true.


Q.

25.9.14.

Q. A fn  $f(x)$  is given by -

$$f(x) = \begin{cases} 0, & x \leq 0 \\ 5x-4, & 0 < x \leq 1 \\ 4x^2-3x, & 1 \leq x \leq 2 \\ 3x+4, & x > 2 \end{cases}$$

Check continuity @  $x = 0, 1, 2$  -

-  At  $x=0$  → discontinuous  
 At  $x=1$  → continuous.  
 At  $x=2$  → continuous.

### DIFFERENTIABILITY

- A fn  $f$  is said to be differentiable at  $x=a$  if both ~~the~~ limit

$$L.H.D = \lim_{h \rightarrow 0} \frac{f(a-h) - f(a)}{-h}$$

$$R.H.D = \lim_{h \rightarrow 0} \frac{f(a+h) - f(a)}{h}$$

exists and are equal. They are equal to  $f'(a)$ .

$$\begin{aligned} d(x^n) &= nx^{n-1} \\ d\sqrt{x} &= \frac{1}{2\sqrt{x}} \\ d\left(\frac{1}{\sqrt{x}}\right) &= \frac{-1}{2x\sqrt{x}} \\ d\left(\frac{1}{x}\right) &= \frac{-1}{x^2} \\ d\left(\frac{1}{x^2}\right) &= \frac{-2}{x^3} \\ d(e^{ax}) &= ae^{ax} \\ d(a^x) &= a^x \log a \\ d(\sin^{-1}x) &= \frac{1}{\sqrt{1-x^2}} \end{aligned}$$

$$\begin{aligned} d(\sin^{-1}xz) &= \frac{1}{\sqrt{1-x^2}} \\ d(\tan^{-1}x) &= \frac{1}{1+x^2} \\ *** \frac{d}{dx}(x^x) &= x^x(1+\log x) \\ *** \frac{d}{dx}(x^{\frac{1}{x}}) &= x^{\frac{1}{x}} \cdot \frac{1}{x^2}(1-\log x) \\ \frac{d}{dx}(u \pm v) &= \frac{du}{dx} \pm \frac{dv}{dx} \\ \frac{d}{dx}(u \cdot v) &= u \frac{dv}{dx} + v \frac{du}{dx} \\ \frac{d}{dx}\left(\frac{u}{v}\right) &= \frac{v \frac{du}{dx} - u \frac{dv}{dx}}{v^2} \end{aligned}$$

$$\begin{aligned} \text{If } y &= \sin^{-1}\left(\frac{2x}{1+x^2}\right) \\ &= \cos^{-1}\left(\frac{1-x^2}{1+x^2}\right) \\ &= \tan^{-1}\left(\frac{2x}{1-x^2}\right) \end{aligned}$$

$$\text{then } \frac{dy}{dx} = \frac{2}{1+x^2}$$

Q. If  $y = x^{x^{\dots}}$  then  $\frac{dy}{dx} = ?$

$$y = x^y$$

$$\log y = y \log x$$

$$\frac{dy}{y} = y \left(\frac{1}{x} dx\right) + \log x dy$$

$$dy \left(\frac{1}{y} - \log x\right) = \frac{y}{x} dx$$

$$\Rightarrow \frac{dy}{dx} = \frac{(y/x)}{(1/y - \log x)} = \frac{y^2}{x - xy \log x}$$

Q.  $x^y = e^{(x-y)}$  in terms of  $x$   $\frac{dy}{dx} = ?$

$$y \log x = x - y$$

$$y(\log x + 1) = x$$

$$dy = \frac{(\log x + 1) - x\left(\frac{1}{x}\right)}{(\log x + 1)^2} dx$$

$$\frac{dy}{dx} = \frac{\log x}{(\log x + 1)^2}$$

2 marks

Q. If  $x^m \cdot y^n = a^{m+n}$  then  $\frac{dy}{dx}$

$$m \log x + n \log y = m+n \log a$$

$$\frac{m}{x} dx + \frac{n}{y} dy = 0$$

$$\star \frac{dx}{dy} = x$$

$$\Rightarrow \frac{dy}{dx} = -\frac{m}{n} \frac{y}{x} = -\frac{m}{n} \cdot \frac{y}{x}$$

Q.  $x^m \cdot y^n = (x+y)^{m+n} \Rightarrow \frac{dy}{dx} = ?$

$$m \log x + n \log y = m+n \log(x+y)$$

$$\frac{m}{x} dx + \frac{n}{y} dy = \frac{m+n}{x+y} (dx+dy)$$

$$\Rightarrow \left( \frac{m}{x} - \frac{m+n}{x+y} \right) dx = \left( \frac{m+n}{x+y} - \frac{n}{y} \right) dy$$

Imp Result.

$$\Rightarrow \boxed{\frac{dy}{dx} = \frac{y}{x}}$$

Q. If  $y = \sqrt{x + \sqrt{x + \sqrt{x + \dots}}}$  then  $\frac{dy}{dx} = ?$

$$y = \sqrt{x+y}$$

$$\Rightarrow y^2 = x+y$$

$$\Rightarrow 2y dy = dx + dy$$

$$\Rightarrow dy(2y-1) = dx$$

$$\Rightarrow \frac{dy}{dx} = \frac{1}{2y-1}$$

Imp Result

$$\text{If } y = \sqrt{f(x) + \sqrt{f(x) + \sqrt{f(x) + \dots}}} \quad \text{then } \frac{dy}{dx} = \frac{f'(x)}{2y-1}$$

Q.  $y = \sqrt{\tan x + \sqrt{\tan x + \sqrt{\tan x + \dots}}}$  then  $\frac{dy}{dx} = ?$   $\frac{\sec^2 x}{2y-1}$

Q.  $y = \sqrt{\log x + \sqrt{\log x + \sqrt{\log x + \dots}}}$  then  $\frac{dy}{dx} = ?$   $\frac{1/x}{2y-1}$

Q.  $y = \sqrt{\sin x + \sqrt{\sin x + \sqrt{\sin x + \dots}}}$  then  $\frac{dy}{dx} = ?$   $\frac{\cos x}{2y-1}$

Q.  $y = \frac{\sin^{-1} x}{\sqrt{1-x^2}}$  then  $(1-x^2)y_1 - xy = \underline{\hspace{2cm}}$

- Ⓐ 0   Ⓑ 1   Ⓒ y   Ⓓ -y

to save time  
uv form  
is easy than u/v

$$(y\sqrt{1-x^2})^2 = (\sin^{-1} x)^2$$

$$\Rightarrow y^2(1-x^2) = (\sin^{-1} x)^2$$

$$\Rightarrow 2y \cdot \frac{dy}{dx} \cdot (-2x) = 2 \sin^{-1} x \cdot \frac{1}{\sqrt{1-x^2}} dx$$

$$\Rightarrow (1-x^2) 2y dy + y^2(-2x dx) = 2 \sin^{-1} x \cdot \frac{1}{\sqrt{1-x^2}} dx$$

$$\Rightarrow (1-x^2) 2y dy = dx \left( \frac{2 \sin^{-1} x}{\sqrt{1-x^2}} + 2y^2 x \right)$$

$$\Rightarrow 2y [(1-x^2)y_1 - xy] = 2y$$

$$\Rightarrow (1-x^2)y_1 - xy = 1$$

Q.  $y = a \cos(\log x) + b \sin(\log x)$  then  $x^2 y_2 + x y_1 = \underline{\hspace{2cm}}$

$$y_1 = \frac{-a \sin(\log x) + b \cos(\log x)}{x}$$

$$x y_1 = -a \sin(\log x) + b \cos(\log x)$$

$$x y_2 + y_1 \cdot 1 = \frac{-a \cos(\log x) - b \sin(\log x)}{x}$$

$$x^2 y_2 + x y_1 = y$$

Q. If  $y = \tan^{-1} \left[ \frac{(3-x)\sqrt{x}}{1-3x} \right] \Rightarrow \frac{dy}{dx} \Big|_{x=1} = ?$

General differentiation  
 Conserves time  
 ∴ replacement  
 substitute.

Let  $x = \tan^2 \theta \Rightarrow \theta = \tan^{-1} \sqrt{x}$ .

$y = \tan^{-1} \left( \frac{(3 - \tan^2 \theta) \tan \theta}{1 - 3 \tan^2 \theta} \right)$

$= \tan^{-1} \left( \frac{3 - \tan^3 \theta}{1 - 3 \tan^2 \theta} \right)$

$= \tan^{-1} (\tan 3\theta)$

$= 3\theta$

$y = 3 \tan^{-1} \sqrt{x}$

$\Rightarrow \frac{dy}{dx} = 3 \cdot \frac{1}{1+(\sqrt{x})^2} \cdot \frac{1}{2\sqrt{x}}$

$\Rightarrow \frac{dy}{dx} \Big|_{x=1} = \frac{3}{1+1} \cdot \frac{1}{2} = \frac{3}{4}$  Ans

Q.  $y = \tan^{-1} \left( \frac{\cos x}{1 + \sin x} \right) \Rightarrow \frac{dy}{dx} = ?$

$1 + \sin x$   
 we don't have formula  
 But  $1 + \cos x$   
 we have formula.  
 That's why this  
 approach

$= \tan^{-1} \left( \frac{\sin(90-x)}{1 + \cos(90-x)} \right)$

$= \tan^{-1} \left( \frac{\sin(90-x)}{2 \cos^2(45-\frac{x}{2})} \right)$

$= \tan^{-1} \left( \frac{2 \sin(45-\frac{x}{2}) \cdot \cos(45-\frac{x}{2})}{2 \cos^2(45-\frac{x}{2})} \right)$

$= 45 - \frac{x}{2}$

$y = \frac{\pi}{4} - \frac{x}{2}$

$\Rightarrow \frac{dy}{dx} = -\frac{1}{2}$  Ans

Q. If  $x = a \cos^3 \theta$ ,  $y = a \sin^3 \theta \Rightarrow \frac{dy}{dx} = ?$

$$\begin{aligned} dx &= -3a \cos^2 \theta \times \cancel{\cos \theta} \times \sin \theta \\ dy &= 3a \sin^2 \theta \times \cancel{\sin \theta} \times \cos \theta \\ \Rightarrow \frac{dy}{dx} &= \frac{3a \sin^2 \theta \times \cos \theta}{-3a \cos^2 \theta \times \sin \theta} = -\tan^2 \theta \end{aligned}$$

$$\frac{dy}{dx} = \frac{3a \sin^2 \theta \times \cancel{\cos \theta}}{-3a \cos^2 \theta \times \cancel{\sin \theta}} = -\tan \theta$$

Q.  $x = 3 \cos \theta - \cos^3 \theta$

$y = 3 \sin \theta - \sin^3 \theta$

$$dy = 3 \cos \theta - 3 \sin^2 \theta \cos \theta = 3 \cos \theta (1 - \sin^2 \theta)$$

$$dx = -3 \sin \theta + 3 \cos^2 \theta \cdot \sin \theta = -3 \sin \theta (1 - \cos^2 \theta)$$

$$\therefore \frac{dy}{dx} = \frac{3 \cos \theta \cdot \cos^2 \theta}{-3 \sin \theta \cdot \sin^2 \theta} = -\cot^3 \theta \quad \underline{\text{Ans}}$$

Q.  $x = a(\theta + \sin \theta)$

$y = a(1 - \cos \theta)$

$$\frac{dy}{dx} = \frac{a(0 + \cos \theta)}{a(1 + \cos \theta)} = \frac{a \sin \theta/2 \cdot \cancel{\cos \theta/2}}{a \cos^2 \theta/2} = \tan \frac{\theta}{2} \quad \underline{\text{Ans}}$$

Q.  $x = a[\theta \sin \theta + \cos \theta]$

$y = a[\sin \theta - \theta \cos \theta]$

$$\frac{dy}{dx} = \tan \theta \quad \underline{\text{Ans}}$$


## # MEAN VALUE THEOREM:-

### (1). Rolle's mean value theorem:-

Let  $f(x)$  be a function defined such that -

- i)  $f(x)$  is continuous in  $[a, b]$
- ii)  $f(x)$  is differentiable in  $(a, b)$
- iii)  $f(a) = f(b)$

then  $\exists$  at least one value  $c \in (a, b)$  such that  $f'(c) = 0$



Proof:-

$$f'(c) = \text{slope of tangent line..}$$

$$= \text{slope of } \overline{AB} \quad \left\{ \because \text{two parallel line have equal slopes} \right\}$$

$$= \frac{y_2 - y_1}{x_2 - x_1}$$

$$= \frac{f(b) - f(a)}{b - a}$$

$$= \frac{0}{b - a}$$

$$\left\{ \because f(a) = f(b) \right\}$$

$$\boxed{f'(c) = 0}$$

Q. Find  $c$  for the Rolle's mean value theorem for -

$$f(x) = x^3 - 4x \text{ in } [-2, 2]. \quad \textcircled{a} \frac{2}{\sqrt{3}} \quad \textcircled{b} -\frac{2}{\sqrt{3}} \quad \textcircled{c} \pm \frac{2}{\sqrt{3}} \quad \textcircled{d} \text{none}$$

$$\rightarrow f(-2) = -8 + 8 = 0$$

$$f(2) = 8 - 8 = 0$$

$$\therefore f'(x) = 3x^2 - 4$$

$$f'(c) = 3c^2 - 4 = 0$$

$$\Rightarrow c^2 = \pm \sqrt{\frac{4}{3}} = \pm \frac{2}{\sqrt{3}} \quad \underline{\text{Ans}}$$

But Problem is not over.

Cross-verify the points in case of  $\pm \sqrt{\quad}$

$\rightarrow$  are points inside interval or not?

Here  $\pm \frac{2}{\sqrt{3}}$  both are inside so ans is  $\pm \frac{2}{\sqrt{3}} \quad \underline{\text{Ans}}$

Q.  $f(x) = \frac{\sin x}{e^x}$  in  $[0, \pi]$ .

$$f(0) = \frac{0}{1} = 0$$

$$f(\pi) = \frac{0}{e^\pi} = 0$$

$$\therefore f'(x) = \frac{e^x \cos x - \sin x e^x}{e^{2x}}$$

$$f'(c) = e^c \cos c - \sin c \cdot e^c = 0$$

$$\Rightarrow e^c \cos c = e^c \sin c$$

$$\Rightarrow \tan c = 1$$

$$\therefore c = \frac{\pi}{4} \quad \underline{\text{Ans}}$$

$c \in [0, \pi]$  must satisfy this

Q.  $f(x) = \frac{x^2 + ab}{2(a+b)}$ ,  $[a, b]$ ,  $a > 0$ ,  $b > 0$ .

- (a)  $-\sqrt{ab}$     (b)  $\sqrt{ab}$     (c)  $\pm\sqrt{ab}$     (d) none.

$$f'(x) = x(a+b)[2x] - [x^2 + ab][a+b] = 0$$

$$\Rightarrow 2x^2(a+b) = (x^2 + ab)(a+b)$$

$$\Rightarrow x^2 = ab$$

$$\Rightarrow x = \pm\sqrt{ab}$$

But  $x = -\sqrt{ab}$  should not  $\in [a, b]$   
When  $a > 0$ ,  $b > 0$

$$\therefore x = \sqrt{ab}$$

$$\Rightarrow \boxed{c = \sqrt{ab}}$$


# Lagrange's mean value theorem:-

Let  $f(x)$  be a function defined such that -

- i)  $f(x)$  is continuous in  $[a, b]$
- ii)  $f(x)$  is differentiable in  $(a, b)$
- iii)  $f(a) \neq f(b)$

Then  $\exists$  at least one value  $c \in (a, b)$  such that

$$\boxed{f'(c) = \frac{f(b) - f(a)}{b - a}}$$



Q.  $f(x) = \log x$  in  $[1, e]$ .

$f(e) = \log e$

$f(1) = \log 1 = 0.$

$\therefore f'(c) = \frac{\log e - \log 1}{e-1} = \frac{\log e}{e-1}$

$\Rightarrow \frac{1}{c} = \frac{\log e}{e-1}$

$\Rightarrow c = \frac{e-1}{\log e}$  Ans

$\Rightarrow c = e-1 \in (1, e) \quad \left\{ \because e \approx 2.71 \right\}$

Q.  $f(x) = x^3 - 6x^2 + 11x - 6$  in  $[0, 4]$ .

$3x^2 - 12x + 11 = \frac{4^3 - 6(4)^2 + 11(4) - 6 - (-6)}{4-0}$

$\Rightarrow 3x^2 - 12x + 11 = \frac{64 - 96 + 44}{4} = \frac{108 - 96}{4} = \frac{12 \cdot 3}{4} = 3$

$\Rightarrow 3x^2 - 12x + 9 = 0$

$\Rightarrow 3c^2 - 12c + 9 = 0$

$\Rightarrow c = \frac{+12 \pm \sqrt{144 - 96}}{6}$

$= \frac{12 \pm \sqrt{48}}{6}$

$= 2 \pm \frac{4\sqrt{3}}{3}$

$= 2 \pm \frac{2}{\sqrt{3}} \in (0, 4)$

$\therefore c = 2 \pm \frac{2}{\sqrt{3}}$  Ans

## # CAUCHY'S MEAN VALUE THEOREM :-

Let  $f(x)$  and  $g(x)$  be two functions defined such that

i)  $f$  and  $g$  are continuous in the closed interval  $[a, b]$

ii)  $f$  and  $g$  are differentiable in  $(a, b)$

iii)  $g'(x) \neq 0 \forall x \in (a, b)$  then there exists at least one value  $c \in (a, b)$  such that -

$$\boxed{\frac{f'(c)}{g'(c)} = \frac{f(b) - f(a)}{g(b) - g(a)}}$$

Q.  $f(x) = \sqrt{x}$ ,  $g(x) = \frac{1}{\sqrt{x}}$  in  $[a, b]$

A)  $\frac{ab}{2}$

B)  $\frac{a+b}{2}$

C)  $\sqrt{ab}$

D)  $\frac{2ab}{a+b}$

$$\frac{\frac{1}{2}x^{-\frac{1}{2}}}{-\frac{1}{2}x^{-\frac{3}{2}}} = \frac{\sqrt{b} - \sqrt{a}}{\frac{1}{\sqrt{b}} - \frac{1}{\sqrt{a}}}$$

$$\Rightarrow -x^{-\frac{1}{2} + \frac{3}{2}} = \frac{\sqrt{b} - \sqrt{a}}{\frac{\sqrt{a} - \sqrt{b}}{\sqrt{ab}}}$$

$$\Rightarrow \cancel{-} c = \cancel{-} \sqrt{ab}$$

$$\therefore \boxed{c = \sqrt{ab}} \quad \text{Ans}$$

Which  $\in (a, b)$

Q.  $f(x) = e^x$ ,  $g(x) = e^{-x}$  in  $[a, b]$

$$\frac{e^c}{-e^{-c}} = \frac{e^b - e^a}{e^{-b} - e^{-a}}$$

$$\Rightarrow -e^{2c} = \frac{e^b - e^a}{\frac{e^a - e^b}{e^{b+a}}}$$

$$\Rightarrow e^{2c} = e^{b+a}$$

$$\Rightarrow c = \frac{b+a}{2} \quad \underline{\underline{\text{Ans}}}$$

$$0. \quad f(x) = \frac{1}{x^2} \quad g(x) = \frac{1}{x} \quad \text{in } [a, b]$$

$$\rightarrow \frac{f(2c^{-3})}{f(c^{-2})} = \frac{\frac{1}{b^2} - \frac{1}{a^2}}{\frac{1}{b} - \frac{1}{a}}$$

$$\Rightarrow 2c^{-1} = \frac{b^{\cancel{a+b}} \frac{a^2}{\cancel{b^2}}}{\cancel{a+b}} \cdot \frac{\cancel{ab}}{a^2 b^{\cancel{2}}}$$

$$\Rightarrow c = \frac{2ab.}{a+b.}$$

$$0.. \quad f(x) = \sin x, \quad g(x) = \cos x \quad \text{in } [0, \frac{\pi}{2}]$$

$$\Rightarrow \frac{\cos x}{-\sin x} = \frac{1-0}{0-1}$$

$$\Rightarrow -\cot x = -1$$

$$\Rightarrow \tan x = 1.$$

$$\therefore c = \frac{\pi}{4}$$

#

## # Taylor's mean value Theorem:-

Let  $f(x)$  be a function defined such that -

i)  $f(x)$  is continuous in  $[a, a+h]$

ii)  $f^{(n-1)}(x)$  is differentiable in  $[a, a+h]$

then there exists at least one value  $\theta \in (0, 1)$  such that -

$$f(a+h) = f(a) + hf'(a) + \frac{h^2}{2!} f''(a) + \frac{h^3}{3!} f'''(a) + \dots + \frac{h^{n-1}}{(n-1)!} f^{(n-1)}(a) + \frac{h^n}{n!} f^{(n)}(a)$$

$$\text{Let } a+h = x \Rightarrow h = x-a.$$

\*\*\*

$$f(x) = f(a) + (x-a)f'(a) + \frac{(x-a)^2}{2!} f''(a) + \dots$$

## # Maclaurin's formula:- $a=0$

$$f(x) = f(0) + xf'(0) + \frac{x^2}{2!} f''(0) + \frac{x^3}{3!} f'''(0) + \dots$$

①  $f(x) = \sin x$  about  $x=0$

$$f(0) = 0$$

$$f'(0) = \cos 0 = 1$$

$$f''(0) = -\sin 0 = 0$$

$$f'''(0) = -\cos 0 = -1$$

\*\*\*

$$f(x) = \sin x = x - \frac{x^3}{3!} + \frac{x^5}{5!} - \frac{x^7}{7!} + \dots$$

$$\cos x = 1 - \frac{x^2}{2!} + \frac{x^4}{4!} - \frac{x^6}{6!} + \dots$$

Imp Result.

Q.  $f(x) = e^x$  about  $x=0$

$f(0) = 1$

$f'(0) = e^0 = 1$

$f''(0) = e^0 = 1$

$f'''(0) = e^0 = 1$

Imp Result

$$f(x) = e^x = 1 + x + \frac{x^2}{2!} + \frac{x^3}{3!} + \frac{x^4}{4!} + \dots$$

Q.  $f(x) = \log(1+x)$  about  $x=0$ .

$f(0) = 0$

$f'(0) = \frac{1}{1+x} = 1$

$f''(0) = -(1+x)^{-2} = -1$

$f'''(0) = +2(1+x)^{-3} = 2$

$f^{(4)}(0) = -6(1+x)^{-4} = -6$

$\therefore f(x) = \log(1+x) = x + \frac{x^2}{2!} + \frac{x^3}{3!} + \frac{x^4}{4!} (-6)$

Imp Result

$$\log(1+x) = x - \frac{x^2}{2} + \frac{x^3}{3} - \frac{x^4}{4} + \dots$$

Q.  $f(x) = \frac{x}{1+x}$  about  $x=0$ .

$f(x) = \frac{x}{1+x} = 1 - \frac{1}{1+x} = 0$

$f'(x) = \frac{1}{(1+x)^2} = 1$

$f''(x) = \frac{-2}{(1+x)^3} = -2$

$f'''(x) = \frac{6}{(1+x)^4} = 6$

$$\therefore \frac{x}{1+x} = 0 + x(1) + \frac{x^2(-2)}{2!} + \frac{x^3(6)}{3!} + \dots$$

Imp Result

$$\frac{x}{1+x} = x - x^2 + x^3 - x^4 + \dots$$

Alternate method

$$f(x) = \frac{x}{1+x}$$

$$= x(1+x)^{-1}$$

$$= x(1 - x + x^2 - x^3 + \dots) \quad \left\{ \text{Binomial Expansion} \right\}$$

$$= x - x^2 + x^3 - x^4 + \dots$$

Q. Find the coeff. of  $x^2$  in the expansion of  $f(x) = \cos^2 x$  about  $x=0$ .

$$\rightarrow f(x) = \cos^2 x$$

$$= \cos x \cdot \cos x$$

$$= \left( 1 - \frac{x^2}{2!} + \frac{x^4}{4!} - \dots \right) \left( 1 - \frac{x^2}{2!} + \frac{x^4}{4!} - \dots \right)$$

$$= -\frac{x^2}{2!} - \frac{x^2}{2!}$$

$$= -\frac{2x^2}{2!}$$

$$= -x^2 = -1$$

Alternate method

$\therefore$  coeff. of  $x^2$  asked.

$$\hookrightarrow = \frac{f''(0)}{2!}$$

$$\therefore f(x) = \cos^2 x =$$

$$f'(x) = 2 \cos x (-\sin x) = -\sin 2x$$

$$f''(x) = -2 \cos 2x$$

$$\therefore \frac{f''(0)}{2!} = \frac{-2}{2!} = -1$$

Q. find the coeff of  $(x-2)^4$  in the expansion of  $e^x$  about  $x=2$

$$\rightarrow \text{coeff of } (x-2)^4 = \frac{f^{(4)}(a)}{4!}$$

$$f(x) = e^x$$

$$f^{(4)}(x) = e^{x^2} = e^2$$

$$\therefore \frac{f^{(4)}(a)}{4!} = \frac{e^2}{4!} \quad \underline{\text{Ans}}$$

Q. find expansion of  $x^2$  about  $x=1$

(A)  $1-x+x^2$

(B)  $1+x^2$

(C)  $x^2$

(D)  $1+x+\frac{x^2}{2}+\frac{x^3}{3}$

$\left. \begin{array}{l} \because f(x) = x^2 \text{ is pure algebraic fn} \\ \therefore \text{expansion not possible} \end{array} \right\}$

Q.  $f(x) = \log\left(\frac{1+x}{1-x}\right)$  about  $x=0$ .

$$f(x) = \log(1+x) - \log(1-x)$$

$$\log(1+x) = x - \frac{x^2}{2} + \frac{x^3}{3} - \frac{x^4}{4} + \dots$$

$$\log(1-x) = -x - \frac{x^2}{2} - \frac{x^3}{3} - \frac{x^4}{4} + \dots$$

$$\log\left(\frac{1+x}{1-x}\right) = 2\left(x + \frac{x^3}{3} + \frac{x^5}{5} + \dots\right)$$

## # PARTIAL AND TOTAL DERIVATIVES :-

O.D	P.D
$y = f(x)$	$z = f(x, y)$
$z = g(y)$	$u = f(x, y, z)$

Q.  $z = x^2 - xy + y^2$

$$\frac{\partial z}{\partial x} = 2x - y$$

$$\frac{\partial^2 z}{\partial y \partial x} = -1$$

$$\frac{\partial z}{\partial y} = -x + 2y$$

$$\frac{\partial^2 z}{\partial x \partial y} = -1$$

$$\frac{\partial^2 z}{\partial x \partial y} = \frac{\partial^2 z}{\partial y \partial x}$$

### \* HOMOGENEOUS FUNCTION :-

A function  $f(x, y)$  is said to be homogeneous of degree  $n$  in  $x$  and  $y$  if —

$$f(kx, ky) = k^n f(x, y)$$

or else check overall degree of each term must be same.

ex:-  $f(x, y) = x^2 - x^1 y^1 + y^2 \rightarrow$  Homogeneous (Directly by degree)

$$f(kx, ky) = k^2 x^2 - k^2 x y + k^2 y^2$$

$$= k^2 (x^2 - xy + y^2)$$

Note:-

The product of two homogeneous fn is again a homogeneous function....

ex:-  $f(x, y) = (x^3 + y^3)(x^2 + y^2)$

$$n = 3 + 2 = 5$$

If a fn is in rational form, if both numerator and denominator are homogeneous function, therefore the given function is also homogeneous.

ex:-  $f(x,y) = \frac{x^3+y^3}{x^2-y^2}$

$n = 3-2 = 1$  ✓

$f(x,y) = \frac{x^{1/4}+y^{1/4}}{x^{1/5}-y^{1/5}}$

$n = \frac{1}{4} - \frac{1}{5} = \frac{1}{20}$  ✓

### # Euler's Theorem:-

If  $z$  is a homogeneous fn of degree  $n$  in  $x$  and  $y$  then  $x \frac{\partial z}{\partial x} + y \frac{\partial z}{\partial y} = n \cdot z$

Note:-

This formula is applicable directly if  $z$  is an algebraic fn

ex:- Q.  $z = (x^2+y^2)^{1/3}$  then  $x \frac{\partial z}{\partial x} + y \frac{\partial z}{\partial y} = \frac{2}{3} z$

Q.  $z = \frac{xy}{x+y} \Rightarrow x z_x + y z_y = 1 \cdot z$

Q. If  $z = a_0 x^n + a_1 x^{n-1} y + a_2 x^{n-2} y^2 + \dots + a_n y^n$  then  $x \frac{\partial z}{\partial x} + y \frac{\partial z}{\partial y} = n z$

Q. If  $z = \log(x^2+y^2)$  then  $x \frac{\partial z}{\partial x} + y \frac{\partial z}{\partial y} = \underline{\quad}$

Note:-

If  $z$  is not algebraic and let  $\phi(z)$  is algebraic and homogeneous of degree  $n$  in  $x$  and  $y$  then

$$\therefore x \frac{\partial z}{\partial x} + y \frac{\partial z}{\partial y} = \frac{n \phi(z)}{\phi'(z)}$$

Shortcut

So, given  $z = \log(x^2 + y^2)$

$\Rightarrow e^z = x^2 + y^2 \rightarrow$  homogeneous of degree 2

$\therefore x \frac{\partial z}{\partial x} + y \frac{\partial z}{\partial y}$

$= 2 \cdot \frac{e^z}{e^z} = 2$  Ans

Q. If  $z = \sin^{-1}\left(\frac{x^2 + y^2}{x - y}\right)$

$\Rightarrow x z_x + y z_y = 1 \cdot \frac{\sin z}{\cos z} = \tan z$  Ans

Q. If  $z = \tan^{-1}\left(\frac{x^3 + y^3}{x - y}\right)$  then  $x \frac{\partial z}{\partial x} + y \frac{\partial z}{\partial y} = ?$

$\Rightarrow \tan z = \frac{x^3 + y^3}{x - y}$  homogeneous of order 2.

$\therefore x \frac{\partial z}{\partial x} + y \frac{\partial z}{\partial y} = 2 \cdot \frac{\tan z}{\sec^2 z} = \sin 2z$  Ans

Q. If  $z = x^2 \sin^{-1}\left(\frac{y}{x}\right) - y^2 \tan^{-1}\left(\frac{x}{y}\right)$

then  $x \frac{\partial z}{\partial x} + y \frac{\partial z}{\partial y} = 2z$

$\because$  at  $\frac{y}{x} \neq 0$   $y=0$  &  $x=0$   
inverse fn not exists  
so it becomes  $z = x^2 - y^2$

Q. If  $z = (x^3 + y^3) e^{-x/y}$   $\Rightarrow x \frac{\partial z}{\partial x} + y \frac{\partial z}{\partial y} = 3z$

$\because$  when  $x=y=0$   
the  $e^{-x/y}$  does not  
exist.  
 $\therefore z = x^3 + y^3$   
homogeneous of  
order 3

## # Application of Euler's Theorem:-

If  $z$  is an homogeneous fn. of degree  $n$  in  $x$  and  $y$  then —

$$x^2 \frac{\partial^2 z}{\partial x^2} + 2xy \frac{\partial^2 z}{\partial x \partial y} + y^2 \frac{\partial^2 z}{\partial y^2} = n(n-1)z$$

Note:-

This formula is applicable directly if  $z$  is an algebraic fn.

Ex:

Q. If  $z = (x^2 + y^2)^{1/3}$  then  $x^2 \frac{\partial^2 z}{\partial x^2} + 2xy \frac{\partial^2 z}{\partial x \partial y} + y^2 \frac{\partial^2 z}{\partial y^2} = \frac{2}{3}(\frac{2}{3}-1)z$   
 $= -\frac{2}{9}z$

Q. If  $z = \frac{x^3 y^3}{x^2 + y^2}$  then  $x^2 \frac{\partial^2 z}{\partial x^2} + 2xy \frac{\partial^2 z}{\partial x \partial y} + y^2 \frac{\partial^2 z}{\partial y^2} = 4(4-1)z$   
 $= 12z$   
 $\hookrightarrow 6-2=4$

Q. If  $z = \tan^{-1} \left( \frac{x^3 + y^3}{x - y} \right)$  then  $x^2 \frac{\partial^2 z}{\partial x^2} + 2xy \frac{\partial^2 z}{\partial x \partial y} + y^2 \frac{\partial^2 z}{\partial y^2} =$

Shortcut

Note:-

If  $z$  is not algebraic then

$$x \frac{\partial z}{\partial x} + y \frac{\partial z}{\partial y} = \frac{n \phi(z)}{\phi'(z)} = F(z)$$

$$\text{and } x^2 \frac{\partial^2 z}{\partial x^2} + 2xy \frac{\partial^2 z}{\partial x \partial y} + y^2 \frac{\partial^2 z}{\partial y^2} = F(z) [F'(z) - 1]$$

$$\therefore F(z) = z \frac{\tan z}{\sec^2 z} = \sin 2z$$

$$\therefore F(z) [F'(z) - 1] = \sin 2z [2 \cos 2z - 1]$$

$$= \sin 4z - \sin 2z \quad \underline{\text{Ans}}$$

Q. If  $\log\left(\frac{x^4+y^4}{x+y}\right)$  then  $x^2 \frac{\partial^2 z}{\partial x^2} + 2xy \frac{\partial^2 z}{\partial x \partial y} + y^2 \frac{\partial^2 z}{\partial y^2} = ?$

$\rightarrow F(z) = 3 \cdot \frac{e^z}{e^z} = 3$

$F(z)[F'(z)-1] = 3[0-1] = \boxed{-3}$  Ans

# Explicit Function:-

Any function which we can express in the form of either  $y = f(x)$  or  $x = g(y)$  etc is called an explicit fn.

means  $\left. \begin{matrix} y = \text{purely fn of } x \\ x = \text{purely fn of } y \end{matrix} \right\} \Rightarrow \text{Explicit fn.}$

ex:-  $y = ax^2 + bx + c$

Ordinary Derivatives upto  $n^{\text{th}}$  order can be easily evaluated.

# Implicit Function:-

Any function which is not explicit is said to be an implicit function.

ex:-  $y = ax^2 + bxy + c$

Ordinary Derivatives upto  $n^{\text{th}}$  order evaluation becomes complex.

$\therefore$  Partial Derivative is used.

Shortcut

$$\frac{dy}{dx} = - \frac{\partial f / \partial x}{\partial f / \partial y}$$

$$\frac{d^2y}{dx^2} = - \frac{[q^2r - 2pqr + p^2t]}{q^3}$$

$p = \frac{\partial z}{\partial x}$

$q = \frac{\partial z}{\partial y}$

$r = \frac{\partial^2 z}{\partial x^2}$

$s = \frac{\partial^2 z}{\partial x \partial y}$  &  $t = \frac{\partial^2 z}{\partial y^2}$

\* If  $ax^2 + 2hxy + by^2 = 1$ .

Shortcut

$$\frac{d^2y}{dx^2} = \frac{h^2 - ab}{(hx + by)^3}$$

Ex:-

Q.  $2x^2 + 4xy + 3y^2 = 1$

$a = 2$

$b = 3$

$h = 2$

$$\Rightarrow \frac{d^2y}{dx^2} = \frac{2^2 - 2 \times 3}{(2x + 3y)^3} = \frac{-2}{(2x + 3y)^3} \quad \underline{\underline{Ans}}$$

Q.  $x^2 + xy + 2y^2 = 1$

$$\frac{d^2y}{dx^2} = \frac{\left(\frac{1}{2}\right)^2 - 2}{\left(\frac{1}{2}x + 2y\right)^3} = \frac{-14}{(x + 4y)^3} \quad \underline{\underline{Ans}}$$

\* If  $ax^2 + 2hxy + by^2 + 2gx + 2fy + c = 0$

Shortcut

$$\frac{d^2y}{dx^2} = \frac{abc + 2fgh - af^2 - bg^2 - ch^2}{(hx + by + f)^3}$$

ex:-

Q.  $2x^2 + 4xy + 3y^2 + 2x + 4y + 1 = 0$

$a = 2$

$b = 3$

$c = 1$

$h = 2$

$g = 1$

$f = 2$

$$\therefore \frac{d^2y}{dx^2} = \frac{6 + \cancel{2} - \cancel{2} - 3 - 4}{(2x + 3y + 2)^3} = \frac{-1}{(2x + 3y + 2)^3}$$

$$0. \quad y^2 - 5x + 4x^2 = 8$$

$$a = 1$$

$$b = 1$$

$$c = -8$$

$$h = 0$$

$$g = -\frac{5}{2}$$

$$f = 0.$$

$$\frac{d^2y}{dx^2} = \frac{-32 + \frac{25}{4}}{(1y)^3} = -\frac{153}{4y^3}$$

Alternate method (By Explicit)

$$y^2 = 8 + 5x - 4x^2$$

$$y = \sqrt{8 + 5x - 4x^2}$$

$\frac{d^2y}{dx^2}$  will become complex.

$\therefore$  Implicit shortcut is very Easy to use.

## # COMPOSITE FUNCTION

$$\rightarrow (g \circ f)(x) = g[f(x)].$$

$\rightarrow$  Any function in function is said to be a composite fn.  
or

$\rightarrow$  If  $z$  is a fn in  $x$  and  $y$  and  $x, y$  are functions in  $t$  then  $z$  is called a composite fn in  $t$ .

$$z = f(x, y)$$

\* Total Derivative of a composite fn :-

If  $z = f(x, y)$  and  $x = g(t)$ ,  $y = h(t)$  then the total derivative of  $z$  w.r. to  $t$  is denoted by -

$$\boxed{\frac{dz}{dt} = \frac{\partial z}{\partial x} \cdot \frac{dx}{dt} + \frac{\partial z}{\partial y} \cdot \frac{dy}{dt}}$$

$\rightarrow$  Total Derivative means combination of ordinary + partial derivative.

$$\frac{du}{dt} = \frac{\partial u}{\partial x} \frac{dx}{dt} + \frac{\partial u}{\partial y} \frac{dy}{dt} + \frac{\partial u}{\partial z} \frac{dz}{dt}$$

Q.  $z = e^x \sin y$

$x = \log t$

$y = t^2$


$\frac{dz}{dt} = ?$

$$\begin{aligned} \rightarrow \left. \begin{aligned} \frac{\partial z}{\partial x} &= e^x \sin y \\ \frac{\partial z}{\partial y} &= e^x \cos y \\ \frac{\partial x}{\partial t} &= \frac{1}{t} \\ \frac{dy}{dt} &= 2t \end{aligned} \right\} \therefore \frac{dz}{dt} &= e^x \sin y \left( \frac{1}{t} \right) + e^x \cos y (2t) \\ &= \frac{e^x}{t} [ \sin y + 2t^2 \cos y ] \end{aligned}$$

Q.  $u = x^2 + y^2 + z^2$

$x = e^{2t}, y = e^{2t} \cos 3t, z = e^{2t} \sin 3t$

then  $\frac{du}{dt} = ?$



will take time as in ur form.

Alternate Method.

$$u = (e^{2t})^2 + (e^{2t} \cos 3t)^2 + (e^{2t} \sin 3t)^2$$

$$= e^{4t} [ 1 + \cos^2 3t + \sin^2 3t ]$$

$$u = 2e^{4t}$$

$$\frac{du}{dt} = 8e^{4t}$$

0.  $u = x^2 - y^2$   
 $x = e^t \cos t$ ,  $y = e^t \sin t$

$$\left. \frac{du}{dt} \right|_{t=0} = ?$$

$$\rightarrow u = e^{2t} \cos^2 t - e^{2t} \sin^2 t$$

$$= e^{2t} (\cos^2 t - \sin^2 t)$$

$$u = e^{2t} \cos 2t$$

$$\Rightarrow \left. \frac{du}{dt} \right|_{t=0} = 2e^{2t} (-\sin 2t) + 2\cos 2t \cdot e^{2t}$$

$$= 2 \cdot 1 =$$

$$= 2 \underline{\underline{Ans}}$$

Q.  $u = x^3 y e^z$   
 $x = t$ ,  $y = t^2$ ,  $z = \log t$

$$\left. \frac{du}{dt} \right|_{t=2} = ?$$

$$u = t^3 t^2 e^{\log t}$$

$$u = t^5$$

$$\left. \frac{du}{dt} \right|_{t=2} = 5t^4 = 5 \times 2^4 = 80 \underline{\underline{Ans}}$$

Q. If  $x = r \cos \theta$ ,  $y = r \sin \theta$ .

then ①  $\frac{\partial^2 \theta}{\partial x^2} + \frac{\partial^2 \theta}{\partial y^2} = \underline{\hspace{2cm}}$

②  $\frac{\partial^2 \theta}{\partial x \partial y} = \underline{\hspace{2cm}}$

$$\frac{y \sin \theta}{y \cos \theta} = \frac{y}{x}$$

$$\Rightarrow \tan \theta = \frac{y}{x}$$

$$\theta = \tan^{-1} \frac{y}{x}$$

$$\frac{\partial \theta}{\partial x} = \frac{1}{1 + \frac{y^2}{x^2}} \times \frac{-y}{x^2}$$

$$\frac{\partial \theta}{\partial x} = \frac{-y}{x^2 + y^2}$$

$$\frac{\partial^2 \theta}{\partial x^2} = \frac{-y \times -1}{(x^2 + y^2)^2} \times 2x$$

$$= \frac{2xy}{(x^2 + y^2)^2}$$

Similarly  $\frac{\partial^2 \theta}{\partial y^2} = \frac{-2xy}{(x^2 + y^2)^2}$

$$\therefore \boxed{\frac{\partial^2 \theta}{\partial x^2} + \frac{\partial^2 \theta}{\partial y^2} = 0}$$


$$\therefore \frac{\partial \theta}{\partial x} = \frac{-y}{x^2 + y^2}$$

$$\frac{\partial \theta}{\partial x \partial y}$$

## Maxima and Minima.

### Maximum Value:-

If a continuous fn  $f(x)$  increases to a certain value and then decreases that value is called maximum value of the function.



### Minimum Value:-

If a continuous fn  $f(x)$  decreases to a certain value and then increases, that value is called minimum value of the fn.

### Note:-

- i) The maxima and minima occurs alternatively.
- ii) A fn can have several maximum and several minimum values.
- iii) The min<sup>m</sup> value may be greater than the max<sup>m</sup> value.
- iv) The least min<sup>m</sup> value is called the global minimum or universal minimum and the highest max<sup>m</sup> value is called the global max<sup>m</sup> or universal max<sup>m</sup>.

→ For a maxima or minima —

$$f'(x) = 0$$

$$x = \alpha, \beta, \lambda, \dots$$

$x = \alpha$ ,  $f''(\alpha) = 0$ ,  $f$  is minimum at  $x = \alpha$ . Min. value =  $f(\alpha)$

$x = \beta$ ,  $f''(\beta) < 0$ ,  $f$  is maximum at  $x = \beta$ . Max<sup>m</sup> value =  $f(\beta)$

$x = \lambda$ ,  $f''(\lambda) = 0$ ,  $f$  is stationary.

if  $f'''(\lambda) \neq 0$  then ' $\lambda$ ' is said to be inflection point.

Q.

$$f(x) = x^x$$

$$f'(x) = x^x(1 + \log x)$$

$$f''(x) = x^x\left(\frac{1}{x}\right) + (1 + \log x)x^x(1 + \log x)$$

$$f''(x) = x^{x-1} + x^x(1 + \log x)^2$$

$$f'(x) = 0$$

$$\Rightarrow x^x(1 + \log x) = 0$$

↳ can't be never 0

$$\therefore 1 + \log x = 0$$

$$\Rightarrow \log_e x = -1$$

$$\Rightarrow x = e^{-1} = \frac{1}{e}$$

$$f''(x) = f''\left(\frac{1}{e}\right) = \left(\frac{1}{e}\right)^{\frac{1}{e}-1} > 0$$

for  $f(x) = x^x$ ,  $f$  is min<sup>m</sup> at  $x = \frac{1}{e}$

$$\left. \begin{array}{l} 2^2 = 4 > 0 \\ 2^{-2} = \frac{1}{4} > 0 \end{array} \right\}$$

$$\therefore \text{Min}^m \text{ value} = f\left(\frac{1}{e}\right) = \left(\frac{1}{e}\right)^{\frac{1}{e}} = e^{-\frac{1}{e}} \text{ Ans.}$$

Imp Result

Q.

$f(x) = x^{\frac{1}{x}}$  has its max<sup>m</sup> at  $x = e$ .

$$\text{Max}^m \text{ value} = f(e) = e^{\frac{1}{e}}$$

Imp Result

\*\*\*\*

fn	Min value	Max value	At
$x^x$	$e^{-\frac{1}{e}}$	—	$x = \frac{1}{e}$
$x^{\frac{1}{x}}$	—	$e^{\frac{1}{e}}$	$x = e$

f

$$0. \quad f(x) = \frac{\log x}{x} = \frac{1}{x} \log x$$

$$f'(x) = \frac{1}{x} \left( \frac{1}{x} \right) + \log x \left( -\frac{1}{x^2} \right)$$

$$f'(x) = \frac{1}{x^2} (1 - \log x)$$

$$f''(x) = -\frac{2}{x^3} (1 - \log x) + \frac{1}{x^2} \left( -\frac{1}{x} \right)$$

$$f'(x) = 0.$$

$$\Rightarrow \frac{1}{x^2} (1 - \log x) = 0.$$

$$\Rightarrow \log x = 1$$

$$\therefore x = e.$$

$$f''(x) = f''(e) = 0 - \frac{1}{e^3} < 0$$

$\therefore f$  is max<sup>m</sup> at  $x = e$ .

$$\text{Max}^m \text{ value} = f(e) = \frac{\log e}{e} = \frac{1}{e}.$$

$$1. \quad f(x) = x^3 + \frac{3}{x}$$

$$f'(x) = 3x^2 - \frac{3}{x^2}$$

$$f''(x) = 6x + \frac{3 \times 2 x^{-3}}{1} = 6x + \frac{6}{x^3}$$

$$3x^2 - \frac{3}{x^2} = 0$$

$$\Rightarrow x^2 = \frac{1}{x^2}$$

$$\Rightarrow x^4 - 1 = 0$$

$$\Rightarrow (x^2 - 1)(x^2 + 1) = 0$$

$$\Rightarrow x^2 - 1 = 0$$

$$\Rightarrow x = \pm 1$$

$$f''(1) = 6 + 6 = 12 > 0 \rightarrow \text{min}^m \text{ at } x = 1$$

$$f''(-1) = -6 + 6 = 0 \rightarrow \text{max}^m \text{ at } x = -1$$

$$\rightarrow \text{Min}^m \text{ value} = 4$$

$$\rightarrow \text{Max}^m \text{ value} = -4$$

Q  $f(x) = x^2 e^{-x}$

$$f'(x) = e^{-x} \times (2x) + x^2 (-e^{-x}) = 0$$

$$= 2xe^{-x} - x^2 e^{-x} = 0$$

$$\Rightarrow e^{-x} (2x - x^2) = 0$$

$$\Rightarrow \cancel{e^{-x}} \times x(2-x) = 0$$

$$x = 0, 2.$$

$$f''(x) = -2xe^{-x} + e^{-x} \times 2 - [e^{-x}(2x) + x^2(-e^{-x})]$$

$$= e^{-x}(2 - 4x + x^2)$$

at  $x=0$   $f''(x) = 2 > 0 \rightarrow$  min value = 0  $\rightarrow$  min<sup>m</sup> at  $x=0$

at  $x=2$   $f''(x) = \frac{-2}{e^2} < 0 \rightarrow$  max<sup>m</sup> at  $x=2$

Max value =  $4e^{-2}$

Q  $f(x) = x^1 e^{-x}$  has its max at  $x = \underline{1}$ .

Trick \*\*\*

Remember in this type max<sup>m</sup> occurs at degree.

Q

$$f(x) = a \cos x + b \sin x + c$$

$$\text{Min value} = c - \sqrt{a^2 + b^2}$$

$$\text{Max value} = c + \sqrt{a^2 + b^2}$$

Shortcut

Q

$$f(x) = 3 \cos x + 4 \sin x + 2.$$

$$\text{Min value} = 2 - \sqrt{3^2 + 4^2} = 2 - 5 = -3$$

$$\text{Max value} = 2 + \sqrt{3^2 + 4^2} = 2 + 5 = 7$$

$$\begin{aligned}
 \text{Q. } f(x) &= 5 \cos x + 3 \cos\left(x + \frac{\pi}{3}\right) + 3 \\
 &= 5 \cos x + 3 \cos x \cdot \cos \frac{\pi}{3} - 3 \sin x \cdot \sin \frac{\pi}{3} + 3 \\
 &= \cos x \left(5 + \frac{3}{2}\right) - \frac{3\sqrt{3}}{2} \sin x + 3
 \end{aligned}$$

$$\begin{aligned}
 \therefore \text{Min value } &= c - \sqrt{a^2 + b^2} \\
 &= 3 - \sqrt{\left(\frac{13}{2}\right)^2 + \left(-\frac{3\sqrt{3}}{2}\right)^2} \\
 &= 3 - \sqrt{\frac{169}{4} + \frac{27}{4}} \\
 &= 3 - \sqrt{\frac{196}{4}} \\
 &= 3 - \frac{14}{2} \\
 &= -4
 \end{aligned}$$

$$\text{Max value} = 3 + 7 = 10$$

Q. Find the height of the cone of max<sup>m</sup> volume that can be inscribed in a sphere of radius 1 unit.

$$r^2 = 1 - h^2$$

$$\therefore \text{volume of cone} = \frac{1}{3} \pi r^2 h$$

$$= \frac{1}{3} \pi (1 - h^2) (1 + h)$$

$$V = \frac{\pi}{3} (1 + h - h^2 - h^3)$$


$$\frac{dV}{dh} = \frac{\pi}{3} (1 - 2h - 3h^2) = 0$$

$$\Rightarrow 1 - 2h - 3h^2 = 0$$

$$\Rightarrow 3h^2 + 2h - 1 = 0$$

$$h = \frac{-2 \pm \sqrt{4 + 12}}{6} = \frac{-2 \pm 4}{6} = \frac{-8}{6}, \frac{2}{6} \Rightarrow \frac{1}{3}$$

$$\therefore h = \frac{1}{3}$$



$$\therefore \text{height} = 1+h = 1 + \frac{1}{3} = \frac{4}{3} \checkmark$$

Q. find the height of the cylinder of maximum volume that can be inscribed in a sphere of radii  $a$  units.

$$\rightarrow r^2 = a^2 - \frac{h^2}{4}$$

$$\text{volume of cylinder} = \pi r^2 h$$

$$v = \pi r^2 h \\ = \pi \left( a^2 - \frac{h^2}{4} \right) h$$

$$v = \pi \left( a^2 h - \frac{h^3}{4} \right)$$

$$\frac{dv}{dh} = \pi \left( a^2 - \frac{3h^2}{4} \right) = 0$$

$$\Rightarrow a^2 - \frac{3h^2}{4} = 0$$


$$\Rightarrow 3h^2 = 4a^2$$

$$\Rightarrow h^2 = \frac{4a^2}{3}$$

$$h = \frac{2a}{\sqrt{3}}$$

Imp Result.

height for which max<sup>m</sup> volume of cylinder occurs in sphere.



Qf.  $z = f(x, y)$

$$P = \frac{\partial z}{\partial x}$$

$$Q = \frac{\partial z}{\partial y}$$

$$r = \frac{\partial^2 z}{\partial x^2}$$

$$s = \frac{\partial^2 z}{\partial x \partial y}$$

$$t = \frac{\partial^2 z}{\partial y^2}$$

$\rightarrow$  for a maxima and minima, Let  $P$  and  $Q = 0$   
(since 1<sup>st</sup> order derivative are only  $P$  &  $Q$ )

$\rightarrow$  since,  $P$  and  $Q$  are fn. in  $x$  and  $y$

So, we have two eq<sup>n</sup> in  $x$  &  $y$ .

$\rightarrow$  Solve these eq<sup>n</sup>s and get a relation bet<sup>n</sup>  $x$  and  $y$ .

Ex:-  $x = y, x = -y, y = 2x, x = 2y$  etc type

$\rightarrow$  Put this relation either in  $P = 0$  or in  $Q = 0$ ; then the eq<sup>n</sup> transformed to either purely in  $x$  or in  $y$

$\rightarrow$  Solve the Eq<sup>n</sup> and get the roots

Ex -  $x = x_0, x_1, x_2, \dots$  and  $y = y_0, y_1, y_2, \dots$  etc

$(x_0, y_0), (x_1, y_1), \dots$  are critical points.

→ Take a point  $x_0, y_0$ , find the values of  $r, s$  and  $t$

\* **Case-1** → If  $rt - s^2 > 0$  and  $r > 0$ , then the fn attains its minimum at that point.  
→ Min<sup>m</sup> value =  $f(x_0, y_0)$ .

\* **Case-2** → If  $rt - s^2 > 0$  and  $r < 0$ , then the fn attains its maximum at that point.  
→ Max<sup>m</sup> value =  $f(x_1, y_1)$

\* **Case-3** → If  $rt - s^2 < 0$  then the fn has no extremum.

**Case-4** → If  $rt - s^2 = 0$  then the case is doubtful and needs further investigation.

Q.  $f(x, y) = x^2 + y^2 + 6x + 12$  has

a) min<sup>m</sup> at  $(-3, 0)$

b) max<sup>m</sup> at  $(-3, 0)$

c) no extremum

d) none.

$$\begin{aligned} \rightarrow \left. \begin{aligned} p &= 2x + 6 \\ q &= \frac{\partial f}{\partial y} = 2y \\ r &= \frac{\partial^2 f}{\partial x^2} = 2 \\ s &= \frac{\partial^2 f}{\partial x \partial y} = 0 \\ t &= \frac{\partial^2 f}{\partial y^2} = 2 \end{aligned} \right\} \begin{aligned} rt - s^2 &= 2 \cdot 2 - 0^2 > 0 \text{ and } r > 0. \\ &\quad \underbrace{\hspace{10em}}_{\text{minimum}} \end{aligned} \end{aligned}$$

Q.  $f = 1 - x^2 - y^2$  has.

- (a) min at (0,0)
- (b) max at (0,0)
- (c) no extremum
- (d) none.

→ 
$$\left. \begin{aligned} P &= -2x \\ Q &= -2y \\ r &= -2 \\ S &= 0 \\ t &= -2. \end{aligned} \right\} r^2 - s^2 = 4 > 0 \quad \text{and} \quad -2 < 0.$$
  
maximum.

Q.  $f(x,y) = x^3 + y^3 - 3xy$ .

- (a) max at (1,-1)
- (b) min at (1,1)
- (c) max at (1,-1)
- (d) min at (1,1)

→ 
$$\left. \begin{aligned} P &= 3x^2 - 3y \\ Q &= 3y^2 - 3x \\ r &= 6x \\ S &= -3 \\ t &= 6y \end{aligned} \right\} \text{at } 1, -1$$
  
$$6 \times (-6) - 9 < 0 \quad \text{and} \quad r = 6 > 0.$$

~~maxm.~~  
no extremum  
or for multiple points check  
P and Q must be 0 at that point  
then consider max & min  
at 1,1

$$6 \times 6 - 9 > 0 \quad \text{and} \quad r = 6 > 0.$$

minm

Q.  $f(x, y) = x^2 + y^2 + xy + x - 4y + 5$  has \_\_\_\_\_

- (A) min at  $(2, -3)$
- (B) min at  $(-2, 3)$
- (C) max at  $(2, -3)$
- (D) max at  $(-2, 3)$

$$\begin{aligned} \rightarrow \quad & P = 2x + y + 1 \\ & Q = 2y + x - 4 \\ & r = 2 \\ & s = 1 \\ & t = 2 \end{aligned}$$

$$rt - s^2 = 4 - 1 > 0 \quad \& \quad r = 2 > 0.$$

minimum

Q.  $f(x, y) = 4x^2 + 6y^2 - 8x - 4y + 8$  has \_\_\_\_\_

- (A) min  $\frac{8}{3}$
- (B) max  $\frac{8}{3}$
- (C) min  $\frac{10}{3}$
- (D) max  $\frac{10}{3}$

$$\begin{aligned} \rightarrow \quad & P = 8x - 8 = 0 \Rightarrow x = 1 \\ & Q = 12y - 4 = 0 \Rightarrow y = \frac{1}{3} \\ & r = 8 \\ & s = 0 \\ & t = 12 \end{aligned}$$

$$rt - s^2 = 96 > 0 > 0 \quad \& \quad r = 8 > 0$$

minimum

$$\begin{aligned} \therefore f\left(1, \frac{1}{3}\right) &= 4 + 6\left(\frac{1}{3}\right)^2 - 8 - \frac{4}{3} + 8 \\ &= 4 + \frac{2}{3} - 8 - \frac{4}{3} + 8 \\ &= 4 - \frac{2}{3} \\ &= \frac{10}{3} \end{aligned}$$

Q.  $f(x, y) = \sin x + \sin y + \sin(x+y)$  has \_\_\_\_\_

- (a)  $\min \frac{\sqrt{3}}{2}$     (b)  $\max \frac{\sqrt{3}}{2}$     (c)  $\min \frac{3\sqrt{3}}{2}$     (d)  $\max \frac{3\sqrt{3}}{2}$

$$\rightarrow P = \cos x + \cos(x+y)$$

$$Q = \cos y + \cos(x+y)$$

$$r = \frac{\partial^2 f}{\partial x^2} = -\sin x - \sin(x+y) \Big|_{\frac{\pi}{3}, \frac{\pi}{3}} = -\sqrt{3}$$

$$s = \frac{\partial^2 f}{\partial x \partial y} = -\sin(x+y) \Big|_{\frac{\pi}{3}, \frac{\pi}{3}} = -\frac{\sqrt{3}}{2}$$

$$t = \frac{\partial^2 f}{\partial y^2} = -\sin y - \sin(x+y) \Big|_{\frac{\pi}{3}, \frac{\pi}{3}} = -\sqrt{3}$$

$\rightarrow$  Make  $P=0$  or  $Q=0$ .

$$\therefore \cos x + \cos(x+y) = 0$$

$$\cos y + \cos(x+y) = 0$$

---

$$\cos x - \cos y = 0.$$

$$\Rightarrow x = y$$

$$\therefore \cos x + \cos 2x = 0$$

$$\Rightarrow \cos 2x = -\cos x$$

$$\Rightarrow \cos 2x = \cos(\pi - x)$$

$$2x = \pi - x$$

$$\Rightarrow x = \frac{\pi}{3} = y$$

$$\therefore \Delta = r^2 - s^2 = 3 - \frac{3}{4} > 0 \quad \text{and } r = -\sqrt{3} < 0$$

$$\text{max}^m \cdot \frac{3\sqrt{3}}{2}$$

## # Definite and Indefinite Integrals:-

$$\int_a^b f(x) dx = \int_a^c f(x) dx + \int_c^b f(x) dx$$

$$\int_a^b f(x) dx = \int_a^b f(a+b-x) dx$$

$$\int_a^b \frac{f(x)}{f(x) + f(a+b-x)} dx = \frac{b-a}{2}$$

$$\int_0^a f(x) dx = \int_0^a f(a-x) dx$$

$$\int_{-a}^a f(x) dx = 2 \int_0^a f(x) dx, \text{ if } f(x) \text{ is even.}$$

$$= 0, \text{ if } f(x) \text{ is odd.}$$

$$\int_0^{2a} f(x) dx = 2 \int_0^a f(x) dx, \text{ if } f(2a-x) = f(x)$$

$$= 0, \text{ if } f(2a-x) = -f(x)$$

$$\int_0^{\pi/2} \frac{f(\sin x)}{f(\sin x) + f(\cos x)} dx = \int_0^{\pi/2} \frac{f(\tan x)}{f(\tan x) + f(\cot x)} dx = \frac{\pi}{4}$$

$$\int_0^{\pi/2} \sin^n x dx = \int_0^{\pi/2} \cos^n x dx = \frac{(n-1)(n-3)(n-5) \dots}{n(n-2)(n-4) \dots} \times k$$

$$\left\{ \begin{array}{l} k = 1, \text{ if } n \text{ is odd} \\ k = \frac{\pi}{2}, \text{ if } n \text{ is even} \end{array} \right\}$$

$$\int_0^{\pi/2} \sin^m x \cdot \cos^n x \, dx = \frac{(m-1)(m-3)\dots(n-1)(n-3)\dots}{(m+n)(m+n-2)(m+n-4)\dots} \times k$$

$$\left. \begin{array}{l} k=1, \text{ if either } m \text{ or } n \text{ or} \\ \text{both are odd} \\ k=\frac{\pi}{2}, \text{ if both } m \text{ and } n \text{ are} \\ \text{even} \end{array} \right\}$$

$$\int \cos x \, dx = \sin x$$

$$\int \sec^2 x \, dx = \tan x$$

$$\int \sec x \tan x \, dx = \sec x$$

$$\int \tan x \, dx = \log \sec x$$

$$\int \cot x \, dx = \log \sin x$$

$$\int \sinh x \, dx = \cosh x$$

$$\int \cosh x \, dx = \sinh x$$

$$\int \frac{1}{\sqrt{a^2-x^2}} \, dx = \sin^{-1}\left(\frac{x}{a}\right)$$

$$\int \frac{1}{\sqrt{a^2+x^2}} \, dx = \sinh^{-1}\left(\frac{x}{a}\right)$$

$$\int \frac{1}{\sqrt{x^2-a^2}} \, dx = \cosh^{-1}\left(\frac{x}{a}\right)$$

$$\int \frac{1}{a^2+x^2} \, dx = \frac{1}{a} \tan^{-1}\left(\frac{x}{a}\right)$$

$$\int \frac{1}{a^2-x^2} \, dx = \frac{1}{2a} \log \left( \frac{a+x}{a-x} \right)$$

$$\int \frac{1}{x^2-a^2} \, dx = \frac{1}{2a} \log \left( \frac{x-a}{x+a} \right)$$

$$\int \frac{f'(x)}{f(x)} dx = \log f(x)$$

$$\int \frac{f'(x)}{\sqrt{f(x)}} dx = 2\sqrt{f(x)}$$

$$\int \frac{f'(x)}{1+(f(x))^2} dx = \tan^{-1}[f(x)]$$

$$\int [f(x)]^n f'(x) dx = \frac{[f(x)]^{n+1}}{n+1}$$

$$\int e^{f(x)} \cdot f'(x) dx = e^{f(x)}$$

$$\int e^x [f(x) + f'(x)] dx = e^x \cdot f(x)$$

$$\int f(x) g(x) dx = f(x) \int g(x) dx - \int f'(x) \int g(x) dx dx$$

To choose  $f(x)$ , remember I L A T E

Shortcut  
\*\*\*  $\rightarrow \int \text{algebraic} \times e^x dx = e^x \left( \begin{array}{l} \text{algebraic with derivatives of} \\ \text{fn} \end{array} \right) \text{ alternate sign}$

$$\int x e^x dx = e^x (x-1)$$

$$\int x^2 e^x dx = e^x (x^2 - 2x + 2)$$

$$\int x^3 e^x dx = e^x (x^3 - 3x^2 + 6 - 6)$$

$$\int (x^2 + x) e^x dx = e^x [(x^2 + x) - (2x + 1) + 2]$$

Shortcut when  $e^{ax}$  is present

$$\int x e^{ax} dx = e^{ax} \left( \frac{x}{a} - \frac{1}{a^2} \right)$$

$$\int x^2 e^{ax} dx = e^{ax} \left[ \frac{x^2}{a} - \frac{2x}{a^2} + \frac{2}{a^3} \right]$$



Q.  $\int_0^2 |1-x| dx$  <sup>\*\*\*</sup> observe fn within limits  
 When modulus present.

$$= \int_0^2 |1-x| dx = \cancel{x^2/2 - x^2/2}$$

$$= \int_0^1 |1-x| dx + \int_1^2 |x-1| dx$$

$$= \left. x - \frac{x^2}{2} \right|_0^1 + \left. \frac{x^2}{2} - x \right|_1^2$$

$$\Rightarrow 1 - \frac{1}{2} + (2 - 2) - (\frac{1}{2} - 1)$$

$$= \frac{1}{2} + \frac{1}{2}$$

$$= 1. \underline{\text{Ans}}$$

<sup>\*\*\*</sup> When Greatest Integer fn present

Q.  $\int_{-1.5}^1 [x+1] dx$  observe limit bet<sup>n</sup> limits if integer lies  $\rightarrow$  split  
 if no integer lie bet<sup>n</sup> limits  $\rightarrow$  no need to split

~~1~~

$$\int_{-1.5}^{-1} [x+1] dx + \int_{-1}^0 [x+1] dx + \int_0^1 [x+1] dx$$

$$[x+1] = \begin{cases} -1, & -1.5 < x < -1 \\ 0, & -1 < x < 0 \\ 1, & 0 < x < 1 \end{cases}$$

$$\int_{-1.5}^{-1} -1 dx + \int_{-1}^0 0 dx + \int_0^1 1 dx$$

$$= -x \Big|_{-1.5}^{-1} + x \Big|_0^1$$

$$= -(-1 + 1.5) + (1 - 0) = -0.5 + 1 = 0.5 \underline{\text{Ans}}$$

Q.  $\int_1^2 x[x] dx$

$\int_1^2$

$x[x] = 2$

$$\int_1^2 x dx = \frac{x^2}{2} \Big|_1^2 = \left(\frac{2^2-1^2}{2}\right) = \frac{3}{2} \text{ Ans}$$

Q.  $\int_{-1}^1 \frac{1}{1+x^2} dx$  \*\*\*  
When limits  $-a$  to  $a$   
check even or odd.

$$= 2 \int_0^1 \frac{1}{1+x^2} dx$$

$$= 2 \tan^{-1} x \Big|_0^1$$

$$= 2 (\tan^{-1} 1 - \tan^{-1} 0)$$

$$= 2 \left(\frac{\pi}{4} - 0\right)$$

$$= \frac{\pi}{2} \text{ Ans}$$

Q.  $\int_{-a}^a \frac{\sqrt{a+x}}{\sqrt{a-x}}$  no even no odd.  
Always try to eliminate sq. root from  
numerator not from denominator  
WHEN  $\sqrt{f(x)}$  IS PRESENT.

$$= \int_{-a}^a \frac{a+x}{\sqrt{a^2-x^2}}$$

$$= \int_{-a}^a \frac{a}{\sqrt{a^2-x^2}} + \frac{x}{\sqrt{a^2-x^2}} dx$$

even odd.

$$= 2 \int_0^a \frac{a}{\sqrt{a^2-x^2}} dx = 2a \sin^{-1} \left(\frac{x}{a}\right) \Big|_0^a = 2a (\sin^{-1} 1 - \sin^{-1} 0)$$

$$= 2a \left(\frac{\pi}{2} - 0\right)$$

$$= a\pi \text{ Ans}$$

Q.  ~~$\int_0^{\pi} \frac{x \sin x}{1 + \cos^2 x} dx$~~

Q.  $\int_{-a}^a \sqrt{\frac{a-x}{a+x}} = a\pi$  (also)

~~Q. aπ~~ (B)  $\frac{a}{\pi}$  (C)  $\frac{\pi}{a}$  (D)  $\frac{1}{\pi a}$

When sign changes of  $x$  even though answer is same so forget the sign but remember the answer always.

\*\*\*

Q.  $I = \int_0^{\pi} \frac{x \sin x}{1 + \cos^2 x} dx = \frac{\pi^2}{4}$

Imp Result.

For Removal of <sup>algebraic</sup> a function always use —

$$\int_0^a f(x) dx = \int_0^a f(a-x) dx$$

$$I = \int_0^{\pi} \frac{(\pi-x) \sin(\pi-x)}{1 + \cos^2(\pi-x)} dx$$

$$I = \int_0^{\pi} \frac{(\pi-x) \sin x}{1 + \cos^2 x} dx$$

$$2I = \int_0^{\pi} \frac{(\cancel{\pi} + \pi - \cancel{x}) \sin x}{1 + \cos^2 x} dx$$

$$2I = \pi \int_0^{\pi} \frac{\sin x}{1 + (\cos x)^2} dx$$

$$\int \frac{f'(x)}{1 + [f(x)]^2} dx = \tan^{-1}[f(x)]$$

$$I = -\frac{\pi}{2} \left[ \tan^{-1}(\cos x) \right]_0^{\pi} = -\frac{\pi}{2} [\tan^{-1}(-1) - \tan^{-1}(1)]$$

$$= -\frac{\pi}{2} \left( -\frac{\pi}{4} - \frac{\pi}{4} \right) = -\frac{\pi}{2} \cdot -\frac{\pi}{2} = \frac{\pi^2}{4} \text{ Ans}$$

\*\*\*  

$$I = \int_0^{\pi/2} \log(\tan x) dx = 0$$

Imp Results

$$I = \int_0^{\pi/2} \log(\cot x) dx = 0$$

Q. 
$$I = \int_0^{\pi/2} \log(\tan x) dx$$

$$= \int_0^{\pi/2} \log\left(\tan\left(\frac{\pi}{2} - x\right)\right) dx = \int_0^{\pi/2} \log \cot x dx$$

Q. 
$$I = \int_0^{\pi/2} [\log(\tan x) + \log(\cot x)] dx$$

$$= \int_0^{\pi/2} \log\left(\frac{\tan x}{\cot x}\right) dx$$

$$= 0 \text{ Ans}$$

Whenever limit  $-a$  to  $a$  given.  
Check even or odd

Q. 
$$\int_{-\pi}^{\pi} x^4 \sin^5 x dx$$

$\downarrow$  even  $\times$  odd  
 $\downarrow$  odd  
 $\downarrow$  odd

$$= 0$$

odd

Q. 
$$\int_0^{\pi/2} \sin^5 x dx$$

If formula forget  
for Numerator  $\left\{ \begin{array}{l} \text{if power even} \rightarrow \text{write odd no} \\ \text{if power odd} \rightarrow \text{write even no} \end{array} \right.$

~~50~~  

$$= \frac{4 \times 2}{5 \times 3 \times 1} = \frac{8}{15}$$

for Denominator  $\left\{ \begin{array}{l} \text{if power even} \rightarrow \text{write even} \\ \text{if power odd} \rightarrow \text{write odd} \end{array} \right.$

$\left\{ \begin{array}{l} \text{If power = odd} \rightarrow \text{keep result as it is} \\ \text{If power = even} \rightarrow \text{multiply result by } \frac{\pi}{2} \end{array} \right.$

$$Q. \int_0^{\pi/2} \cos^7 x dx = \frac{6 \times 4 \times 2}{7 \times 5 \times 3} = \frac{16}{35}$$

$$Q. \int_0^{\pi/2} \sin^8 x dx = \frac{7 \times 5 \times 3 \times 1}{8 \times 6 \times 4 \times 2} \times \frac{\pi}{2} = \frac{35}{256} \pi$$

$$Q. \int_0^1 \frac{x^6 dx}{\sqrt{1-x^2}}$$

Let  $x = \sin \theta$   
 $dx = \cos \theta d\theta$

if  $x \rightarrow 0$   $\theta \rightarrow 0$

if  $x \rightarrow 1$   $\theta \rightarrow \frac{\pi}{2}$

$$1. \int_0^{\pi/2} \frac{\sin^6 \theta}{\cos^6 \theta} \cos \theta d\theta$$

$$= \frac{5 \times 3 \times 1}{6 \times 4 \times 2} \times \frac{\pi}{2} = \frac{5\pi}{32}$$

$$Q. \int_0^{\pi/2} \sin^4 x \cos^5 x dx$$

both are even then only multiply by  $\frac{\pi}{2}$ .

$$= \frac{3 \times 1 \times 4 \times 2}{9 \times 7 \times 5 \times 3 \times 1}$$

do individually as before in numerator

$$= \frac{8}{315}$$

In Denominator — Start from sum of powers and continue.

$$\begin{aligned}
 Q. \quad & \int_0^{\pi/2} \sin^6 x \cos^4 x \, dx \\
 & = \frac{5 \times 3 \times 1 \times 3 \times 1}{10 \times 8 \times 6 \times 4 \times 2} \times \frac{\pi}{2} = \frac{3\pi}{512}
 \end{aligned}$$

$$\begin{aligned}
 Q. \quad & \int_0^{\pi/2} \sin^7 x \cos^5 x \, dx \\
 & = \frac{2 \times 4 \times 2}{8 \times 6 \times 4 \times 2} = \frac{1}{24}
 \end{aligned}$$

$$Q. \quad \int_0^{\pi} \sin^3 x \, dx$$

$$\because \int_0^{2a} f(x) \, dx = 2 \int_0^a f(x) \, dx \quad \text{if } f(2a-x) = f(x)$$

$$\therefore f(\pi-x) = \sin^3(\pi-x) = \sin^3 \pi = f(x)$$

$$\begin{aligned}
 & \text{Converted in } \frac{\pi}{2} \\
 \therefore & 2 \int_0^{\pi/2} \sin^3 x \, dx
 \end{aligned}$$

$$= 2 \times \frac{2}{3 \times 1} = \frac{4}{3}$$

To use formula

## # Definite and Improper Integrals:-

An integral  $\int_a^b f(x) dx$  is said to be an improper integral if ①  $f$  becomes infinite in the interval of integration,

② one or both of the limits are infinite.

Ex:-  $\int_0^1 \frac{1}{x} dx$ ,  $\int_0^1 \sqrt{\frac{1+x}{1-x}} dx$ ,  $\int_0^3 \frac{1}{(x-1)^{2/3}} dx$ ,  
 $\int_1^{\infty} \frac{1}{x^3} dx$ ,  $\int_{-\infty}^0 \sin bx dx$ ,  $\int_{-\infty}^{\infty} \frac{1}{1+x^2} dx$ .

0.  $\int_0^1 \frac{1}{x} dx = \log x \Big|_0^1 = \log 1 - \log 0 = \log \frac{1}{0} = \log \infty = \infty$   
 Divergent

a.  $\int_0^1 \sqrt{\frac{1+x}{1-x}} dx$ . A/c to prev. discussion  
when limit was -1 to 1 ans was  $\pi$   
Here limit is 0 to 1 ans will be  $\frac{\pi}{2}$

But some extra term would be present also

$$= \int_0^1 \frac{1+x}{\sqrt{1-x^2}} + \frac{x}{\sqrt{1-x^2}} dx$$

$$= \int_0^1 \frac{1}{\sqrt{1-x^2}} dx + \int_0^1 \frac{-1-x^2-2x}{2\sqrt{1-x^2}} dx$$

$$= \left( \sin^{-1} x - \frac{1}{2} \times 2\sqrt{1-x^2} \right) \Big|_0^1$$

$$= \left( \sin^{-1} 1 - \sqrt{1-1} \right) - \left( \sin^{-1} 0 - \sqrt{1-0} \right)$$

$$= \frac{\pi}{2} + 1$$

$$\begin{aligned}
 \text{Q. } & \int_0^3 \frac{1}{(x-1)^{2/3}} dx \\
 &= \int_0^3 (x-1)^{-2/3} dx \\
 &= \left[ \frac{(x-1)^{-2/3+1}}{-\frac{2}{3}+1} \right]_0^3 \\
 &= \frac{(x-1)^{1/3}}{\frac{1}{3}} \Big|_0^3 \\
 &= 3 \left[ 2^{1/3} - \cancel{(0)} \right] \\
 &= 3 \left[ \sqrt[3]{2} + 1 \right] \text{ Ans }
 \end{aligned}$$

$$\begin{aligned}
 \text{Q. } & \int_1^{\infty} \frac{1}{x^3} dx \\
 &= \left( \frac{-1}{2x^2} \right)_1^{\infty} \\
 &= -\frac{1}{2} \left[ \frac{1}{\infty} - \frac{1}{1} \right] \\
 &= -\frac{1}{2} [0 - 1] \\
 &= \frac{1}{2} \text{ Ans }
 \end{aligned}$$

$$\begin{aligned}
 \text{Q. } & \int_{-\infty}^0 \sinh x dx \\
 &= \cosh x dx \Big|_{-\infty}^0 \\
 &= \frac{e^x + e^{-x}}{2} \Big|_{-\infty}^0 \\
 &= \frac{e^0 + e^{-0}}{2} - \frac{e^{-\infty} + e^{\infty}}{2} = -\infty \text{ Ans }
 \end{aligned}$$

$$Q. \int_{-\infty}^{\infty} \frac{1}{1+x^2} dx = 2 \int_0^{\infty} \frac{1}{1+x^2} dx$$

$$= 2 \tan^{-1} x \Big|_0^{\infty}$$

$$= 2 (\cancel{\tan^{-1}(\infty)} - \cancel{\tan^{-1}(0)})$$

$$= 2 \times \frac{\pi}{2}$$

$$= \pi \quad \underline{\text{Ans}}$$

$$Q. \int_0^{\infty} x e^{-x^2} dx$$

$$= \frac{-1}{2} \int_0^{\infty} e^{-x^2} (-2x) dx \quad \left\{ \int e^{f(x)} f'(x) = e^{f(x)} \right\}$$

$$= \frac{-1}{2} (e^{-x^2})_0^{\infty}$$

$$= \frac{-1}{2} [e^{-\infty} - e^{-0}]$$

$$= \frac{-1}{2} [0 - 1] = \frac{1}{2}$$

\*\*\*

Q.

$$\int_0^{\infty} e^{-x^2} dx$$

$$\left. \begin{aligned} \text{Let } t &= x^2 \\ dt &= 2x dx \\ dx &= \frac{dt}{2\sqrt{t}} \end{aligned} \right\}$$

do not take  $t = -x^2$

$$\Rightarrow x^2 = -t$$

sq never becomes

-ve

Wrong Idea

✗

$$x \rightarrow 0 \rightarrow t \rightarrow 0$$

$$x \rightarrow \infty \rightarrow t \rightarrow \infty$$

Imp Result.

\*\*\*  
 $\int \rightarrow$  means  $\Gamma$  fn

$$= \frac{1}{2} \int_0^{\infty} \frac{e^t}{\sqrt{t}} dt$$

$$= \frac{1}{2} \int_0^{\infty} e^t \cdot t^{-\frac{1}{2}} dt$$

$$= \frac{1}{2} \sqrt{\frac{1}{2}} = \frac{\sqrt{\pi}}{2}$$

$$\left\{ \begin{aligned} \Gamma(1) &= 1 \\ \Gamma\left(\frac{1}{2}\right) &= \sqrt{\pi} \end{aligned} \right.$$

Definite Improper Integral  
 When  $n$  becomes complex to evaluate  
 Take Help of  $\Gamma$  function.

$$\Gamma(n) = \int_0^{\infty} e^{-x} x^{n-1} dx$$

$$\Gamma(n+1) = n\Gamma(n), \text{ if } n > 0$$

$$\Gamma(n+1) = n!, \text{ if } n \in \mathbb{N}$$

Do clearly Prev. problem here.

$$Q. \int_0^{\infty} \int_0^{\infty} e^{-(x^2+y^2)} dx dy.$$

$$= \int_0^{\infty} \int_0^{\infty} e^{-x^2} \cdot e^{-y^2} dx dy.$$

$$= \int_0^{\infty} e^{-x^2} dx \int_0^{\infty} e^{-y^2} dy$$

$$= \frac{\sqrt{\pi}}{2} \times \frac{\sqrt{\pi}}{2} = \frac{\pi}{4}.$$

$$Q. \int_0^{\infty} \int_0^{\infty} \int_0^{\infty} e^{-(x^2+y^2+z^2)} dx dy dz$$

$$= \frac{\sqrt{\pi}}{2} \times \frac{\sqrt{\pi}}{2} \times \frac{\sqrt{\pi}}{2} = \frac{\sqrt{\pi}}{8} \text{ Ans}$$

Q.  $\int_0^{\infty} x^3 e^{-x} dx = 3!$  Power will become factorial

Shortcut

~~$= x^3 e^{-x} \left[ \frac{x^3}{-1} - \frac{3x^2}{-2} + \frac{6x}{-3} - \frac{6}{-4} \right]_0^{\infty} = 0 \times \infty$  (indeterminate form)~~

Q.  $\int_0^{\infty} \frac{x}{(x^2+9)^2} dx$

Let  $t = x^2 + 9$

$\Rightarrow dt = 2x dx \Rightarrow dx = \frac{dt}{2x}$

$x \rightarrow 0, t \rightarrow 9$

$x \rightarrow \infty, t \rightarrow \infty$

$= \int_9^{\infty} \frac{x}{t^2} \times \frac{dt}{2x}$

$= \frac{1}{2} \left[ -\frac{1}{t} \right]_9^{\infty}$

$= -\frac{1}{2} \left[ \frac{1}{\infty} - \frac{1}{9} \right]$

$= -\frac{1}{2} \left[ 0 - \frac{1}{9} \right]$

$= \frac{1}{18}$  Ans

Q.  $\int_{-\infty}^{\infty} \frac{x}{(x^2+9)^2} dx$   $\Rightarrow$  fn is odd  
 $\Rightarrow$  result = 0.

## Vector Calculus

### # Gradient of a scalar fn:-

Let  $\phi(x, y, z) = c$  be any scalar fn, then the grad of  $\phi$  is denoted by "grad  $\phi$ " or " $\Delta\phi$ " and is defined as -

$$\Delta\phi = i \frac{\partial\phi}{\partial x} + j \frac{\partial\phi}{\partial y} + k \frac{\partial\phi}{\partial z}$$

we know that -

$$\vec{r} = x\mathbf{i} + y\mathbf{j} + z\mathbf{k}$$

$$|\vec{r}| = \sqrt{x^2 + y^2 + z^2}$$

$$r^2 = x^2 + y^2 + z^2$$

$$\Rightarrow \frac{\partial r}{\partial x} = \frac{x}{r}, \quad \frac{\partial r}{\partial y} = \frac{y}{r}, \quad \frac{\partial r}{\partial z} = \frac{z}{r}$$

$$\Rightarrow \frac{\partial r}{\partial x} = \frac{x}{r}$$

$$\frac{\partial r}{\partial y} = \frac{y}{r}$$

$$\frac{\partial r}{\partial z} = \frac{z}{r}$$

$$\nabla r = i \frac{\partial r}{\partial x} + j \frac{\partial r}{\partial y} + k \frac{\partial r}{\partial z}$$

$$= i \left(\frac{x}{r}\right) + j \left(\frac{y}{r}\right) + k \left(\frac{z}{r}\right)$$

$$\nabla r = \frac{x\mathbf{i} + y\mathbf{j} + z\mathbf{k}}{r}$$

$$\therefore \boxed{\text{grad } r = \nabla r = \frac{\vec{r}}{r}}$$

$$\nabla(\log r) = \frac{1}{r} \nabla r = \frac{1}{r} \cdot \frac{\bar{r}}{r} = \frac{\bar{r}}{r^2}$$

$$\nabla\left(\frac{1}{r}\right) = -\frac{1}{r^2} \nabla r = -\frac{1}{r^2} \cdot \frac{\bar{r}}{r} = -\frac{\bar{r}}{r^3}$$

$$\nabla(r^n) = n r^{n-1} \nabla r = n r^{n-1} \frac{\bar{r}}{r} = n r^{n-2} \bar{r}$$

### # Tangent vector to a curve:-

Let  $\bar{r}(t)$  be the given vector curve, then  $\frac{d\bar{r}}{dt}$  is called the tangent vector to the given curve

$$\bar{r} = x\mathbf{i} + y\mathbf{j} + z\mathbf{k}$$

(fn of  $x$ )

$$\frac{d\bar{r}}{dt} = \frac{dx}{dt} \mathbf{i} + \frac{dy}{dt} \mathbf{j} + \frac{dz}{dt} \mathbf{k}$$

### # Normal to a surface:-

Let  $\phi(x, y, z) = c$  be any surface then

$\Delta\phi$  is called ~~is~~ the normal to the surface  $\phi$ .

and  $\frac{\Delta\phi}{|\Delta\phi|}$  is called the unit normal vector to the

Surface ' $\phi$ ' and is denoted by ' $N$ '

$$\text{i.e., } \phi(x, y, z) = c$$

$$\text{then } N = \frac{\nabla\phi}{|\nabla\phi|}$$

## # Directional Derivative:-

Let  $\phi(x, y, z) = c$  be any surface, then

$\nabla\phi \cdot e$  is called the directional derivative to the surface  $\phi$ . Where  $e$  is the unit ~~normal~~ vector in the dir<sup>n</sup> of given vector.


$$\text{i.e. } e = \frac{\vec{a}}{|\vec{a}|} \quad \checkmark$$

The max<sup>m</sup> value of the directional derivative or the greatest value of directional derivative or the magnitude of a gradient of a scalar fn  $\phi$  is

$$\checkmark \quad \text{defined by } |\Delta\phi|$$

## # The angle between the curves:-

The angle bet<sup>n</sup> the two curves is the angle bet<sup>n</sup> tangents at their point of intersection:



Note:-

If the two curves cut each other orthogonally then

$$a \cdot b = 0 \quad \left\{ \begin{array}{l} \because \theta \text{ will become } 90^\circ \\ \& \cos\theta = \cos 90^\circ = 0 \end{array} \right\}$$

## # The angle bet<sup>n</sup> two surfaces:-

The angle bet<sup>n</sup> two surfaces is the angle bet<sup>n</sup> normals drawn at their surfaces.

If  $f$  and  $g$  are any two surfaces and  $\theta$  is the angle bet<sup>n</sup> them then

$$\cos \theta = \frac{\nabla f \cdot \nabla g}{|\nabla f| |\nabla g|}$$

Note:-

If the two surfaces cuts each orthogonally then  $\nabla f \cdot \nabla g = 0$ .

Q. Find the unit normal vector to surface at the point  $(1, 2, -1)$  where  $\phi = x^3 + y^3 + 3xyz = 3$   
(surface)

$$\rightarrow \phi = x^3 + y^3 + 3xyz = 3.$$

$$\nabla \phi = (3x^2 + 3yz) \hat{i} + (3y^2 + 3xz) \hat{j} + (3xy) \hat{k}$$

$$\nabla \phi \Big|_{1,2,-1} = -3\hat{i} + 9\hat{j} + 6\hat{k}.$$

$\therefore N \hat{=} \frac{\nabla \phi}{|\nabla \phi|}$  is unit normal to surface

$$\therefore N = \frac{3(-\hat{i} + 3\hat{j} + 2\hat{k})}{3\sqrt{1+9+4}} = \frac{-\hat{i} + 3\hat{j} + 2\hat{k}}{\sqrt{14}}$$

Q. Unit normal to  $x^2y + 2xz = 4$  at  $(2, -2, 3)$

(A)  $\frac{i + 2j - 2k}{3}$

(B)  $\frac{i - 2j + 2k}{3}$

(C)  $\frac{-i + 2j + 2k}{3}$

(D)  $\frac{i + j + 2k}{3}$

$$\begin{aligned} \rightarrow \nabla\phi|_{2, -2, 3} &= (2xy + 2z)i + (x^2)j + (2x)k \\ &= -2i + 4j + 4k \end{aligned}$$

$$\therefore N = \frac{\nabla\phi}{|\nabla\phi|} = \frac{-i + j + 2k}{\sqrt{1+4+4}} = \frac{-i + j + 2k}{3}$$

2011

Q. U.N to  $x^2 + y^2 + z^2 = 1$  at  $(\frac{1}{\sqrt{2}}, 0, \frac{1}{\sqrt{2}})$

(A)  $\frac{i+j}{\sqrt{2}}$

(B)  $\frac{j+k}{\sqrt{2}}$

(C)  $\frac{i+k}{\sqrt{2}}$

(D)  $\frac{i+j+k}{\sqrt{3}}$

Type-1  
Q.

### DIRECTIONAL DERIVATIVE (4 types total)

find the directional derivative of  $f = xy + yz + zx$  at  $(1, 2, 0)$  in the dir<sup>n</sup> of  $i + 2j + 2k$ .

$$\begin{aligned} \rightarrow \nabla \phi &= i \frac{\partial \phi}{\partial x} + j \frac{\partial \phi}{\partial y} + k \frac{\partial \phi}{\partial z} & a &= i + 2j + 2k \\ \nabla f &= i(y+z) + j(x+z) + k(y+x) & e &= \frac{a}{|a|} = \frac{i+2j+2k}{\sqrt{1+4+4}} \\ \nabla f \Big|_{(1,2,0)} &= 2i + j + 3k & e &= \frac{i+2j+2k}{3} \end{aligned}$$

$$\begin{aligned} D \cdot D &= \nabla f \cdot e \\ &= 2\left(\frac{1}{3}\right) + 1\left(\frac{2}{3}\right) + 3\left(\frac{2}{3}\right) \\ &= \frac{10}{3} \quad \checkmark \end{aligned}$$

Q. Find D.D of  $f = 2xy + z^2$  at  $(1, -1, 3)$  in the dir<sup>n</sup> of  $i + 2j + 3k$ .

$$\begin{aligned} \rightarrow \nabla f \Big|_{1,-1,3} &= (2y)i + (2x)j + (2z)k & a &= i + 2j + 3k \\ &= -2i + 2j + 6k & e &= \frac{i+2j+3k}{\sqrt{1+4+9}} \\ & & e &= \frac{i+2j+3k}{\sqrt{14}} \end{aligned}$$

$$\begin{aligned} D \cdot D &= \nabla f \cdot e \\ &= -2\left(\frac{1}{\sqrt{14}}\right) + 2\left(\frac{2}{\sqrt{14}}\right) + 6\left(\frac{3}{\sqrt{14}}\right) \\ &= \frac{20}{\sqrt{14}} \quad \checkmark \end{aligned}$$

Type-II.

Q. find the directional derivative of  $f = x^2 - y^2 + 2z^2$  at  $P(1, 2, 3)$  in the dir<sup>n</sup> of  $\overline{PQ}$  where  $Q(5, 0, 4)$

$$\rightarrow a = \overline{PQ} = Q - P = 4i - 2j + k$$

$$e = \frac{a}{|a|} = \frac{4i - 2j + k}{\sqrt{16 + 4 + 1}} = \frac{4i - 2j + k}{\sqrt{21}}$$

dot product

$$\nabla f|_{1,2,3} = (2x)i - (2y)j + (4z)k = 2i - 4j + 12k$$

$$\therefore D \cdot D = \frac{8 + 8 + 12}{\sqrt{21}} = \frac{28}{\sqrt{21}} \quad \underline{\underline{Ans}}$$

Q. find the D.D of  $f = xy + yz + zx$  at  $P(1, 2, -1)$  in the dir<sup>n</sup> of  $\overline{PQ}$  where  $Q = 1, 2, 3$ .

$$\rightarrow \nabla f|_{1,2,-1} = (y+z)i + (x+z)j + (x+y)k = i + 3k$$

$$a = \overline{PQ} = Q - P = 4k$$

$$\therefore e = \frac{a}{|a|} = \frac{4k}{4}$$

$$e = k$$

$$\therefore \boxed{D \cdot D = 3}$$

### Type-III

Q. find the directional derivative of  $f = xy^2 + yz^2 + zx^2$  at  $(1,1,1)$  along the tangent to the curve  $x=t, y=t^2, z=t^3$

$$\nabla f \Big|_{(1,1,1)} = i(y^2 + 2xz) + j(2xy + z^2) + k(2yz + x^2)$$

$$\nabla f \Big|_{(1,1,1)} = (3i + 2j + 3k)$$

$$\therefore D \cdot D = \nabla f \cdot e$$

$$= \frac{3+6+9}{\sqrt{14}}$$

$$= \frac{18}{\sqrt{14}} \quad \checkmark$$

$$a = \frac{dr}{dt} = \frac{dx}{dt} i + \frac{dy}{dt} j + \frac{dz}{dt} k$$

$$x=t \quad \Big| \quad y=t^2 \quad \Big| \quad z=t^3$$

$$\frac{dx}{dt} = 1 \quad \Big| \quad \frac{dy}{dt} = 2t \quad \Big| \quad \frac{dz}{dt} = 3t^2$$

→ To take  $t$  value take easy relation.

→ If you choose complicated relation to find ' $t$ ' then after verifying we know that for other ~~points~~  <sup>$t$</sup>  our point varies (Ultimately time loss)

$$\therefore a = i + 2tj + 3t^2k$$

$$= i + 2j + 3k \quad \left\{ \because t=1 \right\}$$

$$\therefore e = \frac{a}{|a|} = \frac{i + 2j + 3k}{\sqrt{14}}$$

### Type-IV

Q. find the directional derivative of surface  $f = xy^2 + z^2$  at  $(1, 1, 1)$  in the dir<sup>n</sup> of normal to surface  $3xy^2 + y = z$  at  $(0, 1, 1)$

$$\nabla f = i(yz^2 + z) + j(xz^2) + k(2xy + 1)$$

$$\left. \nabla f \right|_{(1,1,1)} = 2i + j + 3k$$

$$\therefore D \circ D = \nabla f \cdot e$$

$$= \frac{2+1+3}{\sqrt{11}}$$

$$= \frac{4}{\sqrt{11}}$$

$$a = \nabla g|_{(0,1,1)} = i(3y^2) + j(6xy+1) + k(-1)$$

$$= 3i + j - k$$

$$e = \frac{a}{|a|} = \frac{3i + j - k}{\sqrt{11}}$$

Q. find the maximum value of the directional derivative to the surface  $\phi = x^2yz^3$  at  $(2, 1, -1)$

$$\rightarrow \nabla \phi = i(2xyz^3) + j(x^2z^3) + k(3x^2yz^2)$$

$$\left. \nabla \phi \right|_{(2,1,-1)} = -4i - 4j + 12k$$

$$|\nabla \phi| = \sqrt{16+16+144}$$

$$= \sqrt{176}$$

$$= \sqrt{16 \times 11}$$

$$= 4\sqrt{11}$$

Q. find the greatest value of dir<sup>n</sup> derivative of  $\phi = x^2yz$  at  $(1, 4, 1)$

$$\begin{aligned} \rightarrow \nabla \phi \Big|_{1,4,1} &= (2xyz) \mathbf{i} + (x^2z) \mathbf{j} + (x^2y) \mathbf{k} \\ &= 8 \mathbf{i} + \mathbf{j} + 4 \mathbf{k} \end{aligned}$$

$$\begin{aligned} |\nabla \phi| &= \sqrt{64 + 16 + 1} \\ &= \sqrt{81} \\ &= 9 \quad \checkmark \end{aligned}$$

Q. find the magnitude of gradient  $u = \frac{x^2}{2} + \frac{y^2}{3}$  at  $(1, 3)$

Ⓐ  $\frac{1}{\sqrt{10}}$    Ⓑ  $\frac{1}{\sqrt{5}}$    Ⓒ  $\sqrt{5}$    Ⓓ  $\sqrt{10}$

$$\begin{aligned} \rightarrow \nabla u \Big|_{1,3} &= \frac{2x}{2} \mathbf{i} + \frac{2y}{3} \mathbf{j} \\ &= \mathbf{i} + 2 \mathbf{j} \end{aligned}$$

$$|\nabla u| = \sqrt{1+4} = \sqrt{5}$$

Q. find the angle bet<sup>n</sup> surfaces  $x^2+y^2+z^2=9$  and  $x^2+y^2-z=3$  at the point  $(2, -1, 2)$

$$\begin{aligned} \rightarrow \nabla f \Big|_{2,-1,2} &= 2x \mathbf{i} + 2y \mathbf{j} + 2z \mathbf{k} = 4 \mathbf{i} - 2 \mathbf{j} + 4 \mathbf{k} \\ \nabla g &= 2x \mathbf{i} + 2y \mathbf{j} - \mathbf{k} = 4 \mathbf{i} - 2 \mathbf{j} - \mathbf{k} \end{aligned}$$

$$\cos \theta = \frac{\nabla f \cdot \nabla g}{|\nabla f| |\nabla g|} = \frac{16 + 4 - 4}{\sqrt{36} \sqrt{21}} = \frac{16}{36\sqrt{21}}$$

$$\therefore \theta = \cos^{-1} \left( \frac{8}{3\sqrt{21}} \right) \quad \checkmark$$

Q. find acute angle bet<sup>n</sup> the surfaces  $xy^2z = 3x + z^2$  and  $3x^2 - y^2 + 2z = 1$  at the point  $(1, -2, 1)$ .

$$\rightarrow \nabla f \Big|_{1, -2, 1} = i(y^2z - 3) + j(2xyz) + k(xy^2 - 2z)$$

$$= i(+7) + j(-4) + k(2)$$

$$\nabla g = 6xi - 2yj + 2k$$

$$(\nabla g)_{1, -2, 1} = 6i + 4j + 2k$$

$$\therefore \cos \theta = \frac{|6 - 16 + 4|}{\sqrt{1+16+4} \sqrt{36+16+4}}$$

Due to acute angle we will take only the  $\theta$ .

$$= \frac{6}{\sqrt{7 \times 3} \sqrt{7 \times 4 \times 2}}$$

$$= \frac{6^3}{2 \times 7 \times \sqrt{6}}$$

$$\theta = \cos^{-1} \left( \frac{3}{7\sqrt{6}} \right)$$

Q. find the const.  $a, b$  so that the surfaces  $ax^2 - byz = (a+2)x$  and  $4x^2y + z^3 = 4$  may intersect orthogonally at the point  $(1, 1, 2)$ .

$$\rightarrow \nabla f = (2ax - a - 2)i - bzj - byk$$

$$\Big|_{1, 1, 2} = (a-2)i - 2bj + bk$$

$$\nabla g = (8xy)i + (4x^2)j + (3z^2)k$$

$$\Big|_{1, 1, 2} = 8i + 4j + 12k$$

N/c to a-

$$\nabla f \cdot \nabla g = 0 \Rightarrow -8(a-2) - 8b + 12b = 0$$

$$\Rightarrow \boxed{b = 2a - 4} \quad \text{--- (1)}$$

A/c to Q- 1, -1, 2 lies also on surface ① -

$$\therefore a(1)^2 - b(-1)(2) = (a+2) \quad \text{--- ①}$$

$$\Rightarrow a + 2b = a + 2$$

$$\Rightarrow 2b = 2$$

$$\Rightarrow b = 1 \quad \underline{\underline{\text{Ans}}}$$

$\therefore$  from ① -

$$a = \frac{5}{2} \quad \underline{\underline{\text{Ans}}}$$

# Divergence of a vector function :-

Let  $\mathbf{F} = F_1 \mathbf{i} + F_2 \mathbf{j} + F_3 \mathbf{k}$  be any differentiable vector fn then the divergence of  $f$  is denoted by  $\text{div } \mathbf{F}$  or  $\nabla \cdot \mathbf{F}$  and is defined as -

$$\text{div } \mathbf{F} = \nabla \cdot \mathbf{F} = \frac{\partial F_1}{\partial x} + \frac{\partial F_2}{\partial y} + \frac{\partial F_3}{\partial z}$$

Ex:-  $\bar{r} = x\mathbf{i} + y\mathbf{j} + z\mathbf{k}$   
 $\text{div } \bar{r} = 1 + 1 + 1$   
 $\therefore \text{div } \bar{r} = \nabla \cdot \bar{r} = 3$

# Solenoidal Vector :-

The necessary and sufficient cond<sup>n</sup> for a vector  $f$  to be solenoidal is -

$$\nabla \cdot \mathbf{F} = 0$$

$\nabla \cdot \mathbf{F} = 0$

Note:-

Let  $\phi$  is a scalar and  $a$  is vector then—

$$\text{div}(\phi A) = \nabla \cdot (\phi A) = (\text{grad } \phi) \cdot A + \phi (\text{div } A)$$

$$\nabla \cdot (\phi A) = (\nabla \phi) \cdot A + \phi (\nabla \cdot A)$$

Q.  $\nabla^2 \left( \frac{1}{r} \right) = 0$  where  $r = |\vec{r}|$  and  $\vec{r} = xi + yj + zk$

Imp Result.

$$\rightarrow \nabla \cdot \nabla \left( \frac{1}{r} \right) = \nabla \cdot \left( -\frac{1}{r^2} \nabla r \right) = \nabla \cdot \left( -\frac{1}{r^2} \frac{\vec{r}}{r} \right) = \nabla \cdot \left( \frac{-1}{r^3} \cdot \vec{r} \right)$$

$$= \nabla \left( -\frac{1}{r^3} \right) \cdot \vec{r} + \left( -\frac{1}{r^3} \right) (\nabla \cdot \vec{r})$$

$$= \frac{3}{r^4} \nabla r \cdot \vec{r} - \frac{3}{r^3}$$

$$= \frac{3}{r^4} \frac{\vec{r}}{r} \cdot \vec{r} - \frac{3}{r^3}$$

$$= \frac{3}{r^4} \times r^2 - \frac{3}{r^3}$$

$$= \frac{3}{r^3} - \frac{3}{r^3}$$

$$= 0$$

\*\*\*

$$\text{SHORTCUT — } \nabla^2 [f(r)] = f''(r) + \frac{2}{r} f'(r)$$

$$\textcircled{1} \nabla^2 (\log r) = -\frac{1}{r^2} + \frac{2}{r} \cdot \frac{1}{r} = \frac{1}{r^2}$$

$$\textcircled{2} \nabla^2 (r^n) = n(n-1)r^{n-2} + \frac{2}{r} n r^{n-1}$$

$$= r^{n-2} [n(n-1) + 2n]$$

$$= r^{n-2} [n(n+1)]$$

## # Curl of a vector function :-

Let  $F = F_1 i + F_2 j + F_3 k$  be any differentiable vector function then the curl of  $F$  is denoted by  $\text{Curl } F$  or  $\nabla \times F$  and is defined as -

$$\nabla \times F = \begin{vmatrix} i & j & k \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ F_1 & F_2 & F_3 \end{vmatrix}$$

Ex:-  $\vec{r} = xi + yj + zk$

$$\begin{aligned} \text{Curl } \vec{r} = \nabla \times \vec{r} &= \begin{vmatrix} i & j & k \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ x & y & z \end{vmatrix} \\ &= i(0-0) - j(0-0) + k(0-0) \\ &= 0 \end{aligned}$$

$$\therefore \boxed{\text{Curl } \vec{r} = 0}$$

## # Irrotational Vectors :-

The necessary and sufficient cond<sup>n</sup> for a vector fn  $F$  to be irrotational is -

$$\boxed{\nabla \times F = 0}$$

Note :-

Let  $F$  be any non zero vector function then - ~~div~~

$$\text{div}(\text{Curl } F) = 0$$

$$\nabla \cdot (\nabla \times F) = 0.$$

Q. If  $F = (x+2y)i + (y+2z)j + (x+\lambda z)k$  is solenoidal then value of  $\lambda = ?$

$$\rightarrow \nabla \cdot F = 0 \quad (\text{For Solenoidal})$$

$$\Rightarrow \nabla \cdot [(x+2y)i + (y+2z)j + (x+\lambda z)k] = 0$$

$$\Rightarrow 1 + 1 + \lambda = 0$$

$$\Rightarrow \lambda = -2$$

Q. Find the value of  $\lambda$  so that —

$F = (\lambda x^2 y + yz)i + (4xy^2 + xz)j + 2xyzk$  is zero divergence.

$$\rightarrow \nabla \cdot F = 0$$

$$\Rightarrow 2\lambda xy + 8xy + 2xy = 0$$

$$\Rightarrow xy(2\lambda + 8 + 2) = 0$$

$$\Rightarrow \lambda + 4 + 1 = 0$$

$$\Rightarrow \lambda = -5 \quad \underline{\text{Ans}}$$

Q. If  $f = x^3 + y^3 + z^3 - 3xyz$  then  $\text{div}(\text{grad } f) = \underline{\hspace{2cm}}$

$$\rightarrow \nabla \cdot (\nabla f) = \nabla^2 f$$

$$= 6x + 6y + 6z$$

$$= 6(x + y + z)$$

Q. If  $F = 2xyi - x^2zj$  then  $(\text{curl } F)_{(1,1,1)} = \underline{\hspace{2cm}}$

$$\begin{aligned} \nabla \times F &= \begin{vmatrix} i & j & k \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ 2xy & -x^2z & 0 \end{vmatrix} = i(0+x^2) - j(0-0) + k(-2xz-2x^2) \\ &= i - 4k \quad \text{@ point } (1,1,1) \end{aligned}$$

Q. If  $F = 3x^2i + 5xy^2j + xyz^3k$   
 then  $(\text{div } F) = \underline{\hspace{2cm}}$   
1,2,3

→  $\nabla \cdot F = 6x + 10xy + 3xyz^2$   
 $= 6 + 20 + 54$   
 $= 80$  Ans

Q. If  $\nabla \cdot (r^n \bar{r}) = 0$  then  $n = \underline{\hspace{2cm}}$

→ ∵  $\nabla \cdot (\phi A) = (\nabla \phi) \cdot A + \phi (\nabla \cdot A)$   
 $\Rightarrow (\nabla r^n) \cdot \bar{r} + r^n (\nabla \cdot \bar{r}) = 0$   
 $\Rightarrow n r^{n-1} \nabla r \cdot \bar{r} + 3 r^n = 0$   
 $\Rightarrow n r^{n-2} \frac{\bar{r} \cdot \bar{r}}{r} + 3 r^n = 0$   
 $\Rightarrow n r^{n-2} r^2 + 3 r^n = 0$   
 $\Rightarrow n r^n + 3 r^n = 0$   
 $\Rightarrow n + 3 = 0$   
 $\Rightarrow n = -3$  ✓

Q. Find the constants  $a, b, c$  so that the vector for  
 $F = (x + 2y + az)i + (bx - 3y - z)j + (4x + cy + 2z)k$   
 is irrotational.

→  $\nabla \times F = 0 \Rightarrow$

	$i$	$j$	$k$
	$\frac{\partial}{\partial x}$	$\frac{\partial}{\partial y}$	$\frac{\partial}{\partial z}$
	$x + 2y + az$	$bx - 3y - z$	$4x + cy + 2z$

$$\Rightarrow i(c+1) - j(4-a) + k(b-2) = 0$$

$$\Rightarrow c+1 = 0 \Rightarrow c = -1$$

$$\text{and } 4-a = 0 \Rightarrow a = 4$$

$$\text{and } b-2 = 0 \Rightarrow b = 2$$

Ans

## # Vector Integration:-

① line integral  $\int$

② surface integral  $\iint$

③ volume integral  $\iiint$

PHOTOCOPY & BOOK CENTRE  
NEAR MADE EASY LANE NO. 4  
MAYAPURAM, SAIDULAJAB  
NEW DELHI-110038  
MOBILE: 980120202

### (i). Line Integral:-

Any integral which is evaluated along the curve is called the line integral

Let  $F$  be any differentiable vector function defined along 'curve  $c$ ', then the line integral of  $F$  is defined by —  $\int_c F \cdot d\vec{r}$

If  $c$  is a closed curve, it is denoted by —  $\oint_c F \cdot d\vec{r}$

### Cartesian form:-

$$\text{Let } F = F_1 \hat{i} + F_2 \hat{j} + F_3 \hat{k}$$

$$\vec{r} = x\hat{i} + y\hat{j} + z\hat{k}$$

$$d\vec{r} = dx\hat{i} + dy\hat{j} + dz\hat{k}$$

$$\int_c F \cdot d\vec{r} = \int_c (F_1 dx + F_2 dy + F_3 dz)$$

is called  
Cartesian form  
of line integral

Note:- If  $F(x, y, z)$  are fn. of  $t$ , then

$$\int_c F \cdot d\vec{r} = \int \left[ F_1 \frac{dx}{dt} + F_2 \frac{dy}{dt} + F_3 \frac{dz}{dt} \right] dt$$

Q. Evaluate  $\int_C F \cdot d\vec{r}$  where  $F = (2y+3)\vec{i} + xz\vec{j} + (yz-x)\vec{k}$

① Where  $C$  is the curve  $x = 2t^2$   $y = t$ ,  $z = t^3$  joining  $(0,0,0)$  to  $(2,1,1)$

② Where  $C$  is the curve of line ~~joining~~ <sup>joining</sup>  $(0,0,0) \rightarrow (0,0,1) \rightarrow (0,1,1) \rightarrow (2,1,1)$

$$\rightarrow (1). \int_C F \cdot d\vec{r} = \int_C \left[ F_1 \frac{dx}{dt} + F_2 \frac{dy}{dt} + F_3 \frac{dz}{dt} \right] dt$$

$$= \begin{array}{l} x = 2t^2 \quad | \quad y = t \quad | \quad z = t^3 \\ \frac{dx}{dt} = 4t \quad | \quad \frac{dy}{dt} = 1 \quad | \quad \frac{dz}{dt} = 3t^2 \end{array}$$

$$\begin{array}{l} \because y = t \\ \therefore y \rightarrow 0, t \rightarrow 0 \\ \quad y \rightarrow 1, t \rightarrow 1 \end{array}$$

$$= \int_{t=0}^1 \left[ \frac{(2y+3) \cdot 4t}{t} + \frac{xz \cdot 1}{2t^2} + \frac{(yz-x) \cdot 3t^2}{t \cdot 2t^2} \right] dt$$

$$= \int_{t=0}^1 [8t^2 + 12t + 2t^5 + 3t^6 - 6t^4] dt$$

$$= \left[ \frac{8t^3}{3} + \frac{12t^2}{2} + \frac{2t^6}{6} + \frac{3t^7}{7} - \frac{6t^5}{5} \right]_0^1$$

$$= \frac{8}{3} + 6 + \frac{1}{3} + \frac{3}{7} - \frac{6}{5}$$

$$= 9 + \frac{3}{7} - \frac{6}{5}$$

$$= \frac{288}{35}$$

$$\textcircled{2} \int_C \mathbf{F} \cdot d\mathbf{r} = \int_C F_1 dx + F_2 dy + F_3 dz.$$

Choose  $dx, dy, dz$   
 such that derivative  
 only that varies.  
 Others remain const. or  
 remains const. & other  
 two varies


① along  $\overline{OA}$

$$x=0, y=0$$

$$dx=0, dy=0.$$

$$\text{as } z=0 \text{ to } z=1$$

$$= \int_{OA} (yz-x) dz = 0 \quad \left\{ \begin{array}{l} \because x=0 \\ y=0 \end{array} \right\}$$



② along  $\overline{AB}$

$$x=0, z=1$$

$$dx=0, dz=0.$$

$$\text{as } y=0 \text{ to } 1.$$

$$= \int_{AB} xz dy = 0 \quad \left\{ \begin{array}{l} \because x=0 \end{array} \right\}$$

③ along  $\overline{BC}$

$$y=1, z=1$$

$$dy=0, dz=0.$$

$$\text{as } x=0 \text{ to } 2$$

$$= \int_{x=0}^2 (2y+3) dx = \int_{x=0}^2 [2(1)+3] dx = 5[x]_0^2 = 10$$

$$\therefore \int_C \mathbf{F} \cdot d\mathbf{r} = 0 + 0 + 10 = \underline{\underline{10}}$$

Q.  $\int F \cdot dr$  where  $F = 3x^2i + (2xz - y)j + zk$  where  $c$  is the st. line joining  $(0,0,0)$  to  $(2,1,3)$

$$\rightarrow \text{eqn of OA} = \frac{x-0}{2-0} = \frac{y-0}{1-0} = \frac{z-0}{3-0} = t$$

$$\Rightarrow \frac{x}{2} = \frac{y}{1} = \frac{z}{3} = t$$

$$\Rightarrow x = 2t, y = t, z = 3t$$

The eqn of a st. line joining  $(x_1, y_1, z_1)$  and  $(x_2, y_2, z_2)$  in the symmetric form is  $\frac{x-x_1}{x_2-x_1} = \frac{y-y_1}{y_2-y_1} = \frac{z-z_1}{z_2-z_1} = t$

Where  $t$  is some scalar.

$$\therefore \int_C F \cdot dr = \int_C \left[ F_1 \frac{dx}{dt} + F_2 \frac{dy}{dt} + F_3 \frac{dz}{dt} \right] dt$$

$$= \int_{t=0}^1 [3(4t^2)(2) + (2(2t)(3t) - t)(1) + 3t(3)] dt$$

$$= \int_{t=0}^1 (24t^2 + 12t^2 - t + 9t) dt$$

$$= \left[ 36 \frac{t^3}{3} + \frac{8t^2}{2} \right]_0^1 = 12 + 4 = 16$$

Q.  $\int_C F \cdot dr = ?$  where  $F = 2xyz i + x^2y j + x^2z k$  where  $c$  is the st. line joining  $(0,0,0)$  to  $(1,1,1)$

$\rightarrow$  eqn of st line —

$$\frac{x}{1} = \frac{y}{1} = \frac{z}{1} = t$$

$$x = t, y = t, z = t$$

$$\frac{dx}{dt} = \frac{dy}{dt} = \frac{dz}{dt} = 1$$

$$\begin{aligned} \therefore \int_C F \cdot dr &= \int_C F_1 \frac{dx}{dt} + F_2 \frac{dy}{dt} + F_3 \frac{dz}{dt} \\ &= \int_{t=0}^1 [2t^3(1) + t^3(1) + t^3(1)] dt \\ &= \int_{t=0}^1 4t^3 dt = 4 \frac{t^4}{4} \Big|_0^1 = 1 - 0 = 1 \end{aligned}$$

Q.  $\int_C F \cdot dr$ , where  $F = 3xy \mathbf{i} - y^2 \mathbf{j}$  and  $C$  is the curve  $y = 2x^2$  in the  $xy$  plane joining  $(0, 0)$  to  $(1, 2)$

$$\begin{aligned} \rightarrow \int_C F \cdot dr &= \int_C [3xy dx - y^2 dy] \\ &\quad \begin{array}{l} y = 2x^2 \\ dy = 4x dx \\ x \rightarrow 0 \text{ to } 1 \end{array} \\ &= \int_{x=0}^1 [3x(2x^2) dx - (2x^2)^2 4x dx] \\ &= \left| \frac{6x^4}{4} - \frac{16x^6}{6} \right|_0^1 = \frac{3}{2} - \frac{8}{3} = -\frac{7}{6} \end{aligned}$$

*variable term ki change krni limit krni*

Q.  $\int_C F \cdot dr = ?$  where  $F = (5xy - 6x^2) \mathbf{i} + (2y - 4x) \mathbf{j}$  and  $C$  is the curve  $y = x^3$  in the  $xy$  plane joining  $(1, 1)$  to  $(2, 8)$

$$\begin{aligned} \rightarrow \int_C F \cdot dr &= \int_C (5xy - 6x^2) dx + (2y - 4x) dy \\ &\quad \begin{array}{l} y = x^3 \\ dy = 3x^2 dx \\ x \rightarrow 1 \text{ to } 2 \end{array} \\ &= \int_{x=1}^2 (5x^4 - 6x^2) dx + (2x^3 - 4x) \cdot 3x^2 dx \end{aligned}$$

$$= \int_{x=1}^2 [5x^4 - 6x^2 + 6x^5 - 12x^3] dx$$

$$= \left[ \frac{5x^5}{5} - \frac{6x^3}{3} + \frac{6x^6}{6} - \frac{12x^4}{4} \right]_{1}^2$$

~~$$= \left[ \frac{5x^5}{5} - \frac{6x^3}{3} + \frac{6x^6}{6} - \frac{12x^4}{4} \right]_{1}^2$$~~

~~$$= 48$$~~

$$= (32 - 16 + 64 - 48) - (1 - 2 + 1 - 3)$$

$$= 32 + 3 = 35 \quad \underline{\underline{Ans}}$$

Q.  $\oint_C F \cdot d\vec{r} = ?$  where  $F = (2x - y + z)\mathbf{i} + (x + y - z^2)\mathbf{j} + (3x + 2y - 4z)\mathbf{k}$

where  $C$  is the circle in  $xy$  plane having the centre at origin and radius 3 units.

$$\rightarrow \int F \cdot d\vec{r} = \int (2x - y) dx + (x + y) dy$$

Circle is in  $xy$  plane  
 $\therefore$  do not consider  $dz$

Our Relation is —

$$x^2 + y^2 = 9$$

$$x = r \cos \theta = 3 \cos \theta$$

$$y = r \sin \theta = 3 \sin \theta$$

$$\Rightarrow dx = -3 \sin \theta d\theta$$

$$dy = 3 \cos \theta d\theta$$

as  $\theta \rightarrow 0$  to  $2\pi$  to complete a circle


$$= \int_{\theta=0}^{2\pi} 2(3 \cos \theta - 3 \sin \theta)(-3 \sin \theta d\theta) + (3 \cos \theta + 3 \sin \theta) \times (3 \cos \theta) d\theta$$

$$= \int_{\theta=0}^{2\pi} (-9(\sin \theta) \cos \theta + 9) d\theta$$

$$= \left[ -9 \frac{\sin^2 \theta}{2} + 9\theta \right]_0^{2\pi}$$

$$= 18\pi$$

Q.  $\oint_C F \cdot d\vec{r}$  where  $F = (x^2 + y^2)\vec{i} - 2xy\vec{j}$  where  $C$  is the rectangle bounded by  $x=0, x=a$   
 $y=0, y=b$ .



$$\int F \cdot d\vec{r} = \oint (x^2 + y^2) dx - 2xy dy$$

① along OA —  $y=0 \rightarrow dy=0$  (not changing)  
 $x \rightarrow 0$  to  $a$ .

$$\int F \cdot d\vec{r} = \int_{x=0}^a (x^2 + 0) dx = \left( \frac{x^3}{3} \right)_0^a = \frac{a^3}{3}$$

② along AB —  $x=a \rightarrow dx=0$  (not changing)  
 $y \rightarrow 0$  to  $b$

$$\int F \cdot d\vec{r} = \int_{y=0}^b -2ay dy = -2a \left( \frac{y^2}{2} \right)_0^b = -ab^2$$

③ along BC —  $y=b \rightarrow dy=0$  (not changing)  
 $x \rightarrow a$  to  $0$

$$\int F \cdot d\vec{r} = \int_{x=a}^0 (x^2 + b^2) dx = 0 - \frac{a^3}{3} - ab^2$$

④ along CO —  $x=0 \rightarrow dx=0$

$$\int F \cdot d\vec{r} = \int 0 - 0 = 0$$

$$\therefore \int F \cdot d\vec{r} = \frac{a^3}{3} - ab^2 - \frac{a^3}{3} - ab^2 - 0 = -2ab^2$$

## # Surface Integral :-

Any integral which is evaluated over surface is called a surface integral.

Let  $F$  be any differentiable vector field defined over a surface  $S$  then the surface integral is defined by  $\iint_S F \cdot N \, ds$  where  $N$  is the outer unit normal vector to given surface.

$$\iint_S F \cdot N \, ds = \iint_{R_1} F \cdot N \frac{dx \, dy}{|N \cdot k|} \quad \text{where } R_1 \text{ is projection in } xy \text{ plane.}$$

$$= \iint_{R_2} F \cdot N \frac{dy \, dz}{|N \cdot j|} \quad R_2 \text{ --- } yz \text{ plane}$$

$$= \iint_{R_3} F \cdot N \frac{dz \, dx}{|N \cdot i|} \quad R_3 \text{ --- } zx \text{ plane}$$

Q. Evaluate  $\iint_S F \cdot N \, ds$  where  $F = z\mathbf{i} + xy\mathbf{j} - 3y^2z\mathbf{k}$  and  $S$  is

the surface of cylinder  $x^2 + y^2 = 16$  included in the I<sup>st</sup>

octant bounded by  $z = 0$  &  $z = 5$ .

$\therefore z$  is given  $\therefore$  either  $xz$  plane or  $yz$  plane will be chosen for projection

Q. Evaluate  $\iint_S F \cdot N \, ds$  where  $F = 4xz\mathbf{i} - y^2\mathbf{j} + yz\mathbf{k}$  and  $S$  is the cube bounding  $0 \leq x, y, z \leq 1$ .

→ for POAR —

$$N = \mathbf{i} \quad \& \quad x = 1 \text{ (const)}$$

$$F \cdot N = 4xz = 4z$$

$$\int_S F \cdot N \, ds = \int_{y=0}^1 \int_{z=0}^1 4z \, dy \, dz$$

$$= 4 \left. \frac{z^2}{2} \right|_0^1 \left. y \right|_0^1$$

$$= 4 \times \frac{1}{2} \times 1$$


$$= 2$$

For SBOC —

$$N = -\mathbf{i}, \quad x = 0$$

$$F \cdot N = -4xz = 0$$

$$\therefore \iint F \cdot N \, ds = 0$$



For PQBS —

$$N = j, \quad y = 1.$$
$$\int_{\partial S} F \cdot N = \int_{x=0}^1 \int_{z=0}^1 -y^2 dx dz = -x \Big|_0^1 z \Big|_0^1 = -1$$

For OCRA —

$$N = j, \quad y = 0.$$

$$\int_S F \cdot N = 0.$$

For SPRC —

$$N = k, \quad z = 1.$$

$$\int_S F \cdot N ds = \int_{x=0}^1 \int_{y=0}^1 yz^{\frac{1}{2}} dx dy$$

$$= \frac{y^2}{2} \Big|_0^1 x \Big|_0^1$$

$$= \frac{1}{2}$$

For BQOA —  $z = 0, N = -k$

$$\int_S F \cdot N ds = \cancel{0} = 0$$

$$\therefore \iint_S F \cdot N ds = 2 + 0 + (-1) + 0 + \frac{1}{2}$$
$$= 1 + \frac{1}{2} = \frac{3}{2} \underline{\underline{\text{Ans}}}$$

## # Volume Integral :-

Any integral which is evaluated over a volume is called the volume integral.

Q.  $\iiint_V \nabla \cdot F \, dv$  where  $F = (2x^2 - 3z)\mathbf{i} - 2xy\mathbf{j} + xz\mathbf{k}$  where  $V$  is the region bounded by  $x=0, y=0, z=0$  and  $2x+2y+z=4$ .

$$\nabla \cdot F = \frac{\partial F_1}{\partial x} + \frac{\partial F_2}{\partial y} + \frac{\partial F_3}{\partial z}$$

$$\rightarrow \nabla \cdot F = 4x - 2x$$

~~(Previous)~~  
 $z \rightarrow$  in terms of  $x$  &  $y \rightarrow 0$  to  $4 - 2x - 2y$   
 $y \rightarrow$  ~~(Previous)~~  $\rightarrow 0$  to  $2 - x$   
 $x \rightarrow$  ~~(Previous)~~  $\rightarrow 0$  to  $2$ .

$$\begin{aligned} \iiint_V \nabla \cdot F \, dv &= \int_0^2 \int_0^{2-x} \int_0^{4-2x-2y} 2x \, dz \, dy \, dx \\ &= 2 \int_0^2 \int_0^{2-x} x(4-2x-2y) \, dy \, dx \\ &= 2 \int_0^2 \left[ 4x - \frac{2x^2}{2} - 2yx \right]_0^{2-x} dx \\ &= 2 \int_0^2 \left( 8 - 4x - 4 + x^2 - 4x + 2yx \right) dx \\ &= 2 \int_0^2 \left( x^2 - 8x + 4 + 2yx \right) dx \\ &= 2 \int_0^2 [4x - 2x^2 - 2xy] \, dx \, dy \\ &= 2 \int_0^2 [4xy - 2x^2y - xy^2]_0^{2-x} \, dx \\ &= 2 \int_0^2 x(2-x)^2 \, dx \\ &= 2 \left[ 4 \frac{x^2}{2} + \frac{x^4}{4} - \frac{4x^3}{3} \right]_0^2 \\ &= 2 \left[ 4 \times \frac{4}{2} + \frac{16^2}{4} - \frac{32}{3} \right] = \frac{8}{3} \text{ Ans } \end{aligned}$$

## Vector Transformation

### # Gauss Divergence Theorem :-

→ The Gauss divergence theorem uses the connection between surface to volume integrals.

→ Let  $S$  be the closed surface enclosed by a volume ' $V$ '. and let  $F$  be any differentiable vector fn then

$$\iint_S F \cdot N \, ds = \iiint_V (\text{div } F) \, dv = \iiint_V (\nabla \cdot F) \, dv.$$

### \* Cartesian form of Divergence Theorem :-

$$\iint_S (F_1 \, dy \, dz + F_2 \, dz \, dx + F_3 \, dx \, dy) = \iiint_V \left( \frac{\partial F_1}{\partial x} + \frac{\partial F_2}{\partial y} + \frac{\partial F_3}{\partial z} \right) dx \, dy \, dz.$$

Q. The value of  $\int_S (axi + byj + czk) \cdot N \, ds$ , where  $S$  is the surface of sphere  $x^2 + y^2 + z^2 = 1$ .

$$\rightarrow \nabla \cdot F = a + b + c$$

$$\therefore \int_S F \cdot N \, ds = \int_V (\text{div } F) \, dv$$

$$= (a+b+c) \frac{4\pi}{3} \times 1^3$$

$$= \frac{4\pi}{3} (a+b+c) \quad \underline{\underline{\text{Ans}}}$$

Q. Find  $\int_S (axi + byj + czk) \cdot N ds$  where  $S$  is sphere

$$x^2 + y^2 + z^2 = 1$$

or  $\int_S (ax^2 + by^2 + cz^2) ds$  where  $S$  is sphere  $x^2 + y^2 + z^2 = 1$

Same Problem

Imp Result  $= (a+b+c) \frac{4\pi}{3}$

Q.  $\int_S \vec{r} \cdot N ds$ , where 's' is the closed region enclosed by a volume 'v'.

$$\rightarrow \nabla \cdot \vec{r} = 3$$

$$\therefore \int_S \vec{r} \cdot N ds = \int_V (\nabla \cdot \vec{r}) dv = 3V \quad \underline{\underline{Ans}}$$

Q.  $\int_S \vec{r} \cdot N ds$  where  $S$  is sphere  $x^2 + y^2 + z^2 = a^2$

$$\rightarrow \nabla \cdot \vec{r} = 3$$

$$\therefore \int_S \vec{r} \cdot N ds = \int_V (\nabla \cdot \vec{r}) dv = \frac{4}{3}\pi a^3 \times 3 = 4\pi a^3 \quad \underline{\underline{Ans}}$$

Q.  $\int_S (x dy dz + y dz dx + z dx dy)$  where  $S$  is the cube of unit length.

$$\rightarrow \nabla \cdot F = 1 + 1 + 1 = 3$$

$$\int_S F \cdot N ds = \int_V (\text{div } F) dv = 3V = 3 \times 1^3 = 3 \quad \underline{\underline{Ans}}$$

Q.  $\int_{\mathbb{B}} (x+z) dy dz + (y+z) dz dx + (x+y) dx dy$  where  
 $\mathbb{B}$  is sphere  $x^2 + y^2 + z^2 = 4$ .

$\Rightarrow \nabla \cdot F = 1 + 1 + 0 = 2$

$\therefore \int_S F \cdot N ds = \int_V (\nabla \cdot F) dV = 2V = 2 \times \frac{4}{3} \pi \times 2^3$   
 $= 2 \cdot \frac{64\pi}{3}$  Ans

Q.  $\int_S (4xz \mathbf{i} - y^2 \mathbf{j} + yz \mathbf{k}) \cdot N ds$  where  $S$  is cube bounded  
 by  $0 \leq x, y, z \leq 1$ .

$\rightarrow \nabla \cdot F = 4z - 2y + y = 4z - y$

$\int_S F \cdot N ds = \int_V (\text{div } F) dV = \int_{x=0}^1 \int_{y=0}^1 \int_{z=0}^1 [4z - y] dx dy dz$

$= (x)_0^1 \left[ 4 \left( \frac{z^2}{2} \right)_0^1 (y)_0^1 - \left( \frac{y^2}{2} \right)_0^1 (z)_0^1 \right]$

$= (1-0) \left[ 2 - \frac{1}{2} \right]$

$= \frac{3}{2}$  Ans

Q.  $\int_S [(x^2 - yz) \mathbf{i} + (y^2 - zx) \mathbf{j} + (z^2 - xy) \mathbf{k}] \cdot N ds$  where

$S$  in the cuboid bounded by  $0 \leq x \leq a$   
 $0 \leq y \leq b$   
 $0 \leq z \leq c$

$\rightarrow \nabla \cdot F = 2x - 2y + 2z$

$\therefore \int_S F \cdot N ds = \int_V (\nabla \cdot F) dV = \int_{x=0}^a \int_{y=0}^b \int_{z=0}^c (2x - 2y + 2z) dx dy dz$

$$= [z^2yz + y^2zx + z^2xy]_{\substack{y=b \\ z=a}}$$

$$\substack{x=0 \\ y=0 \\ z=0}$$

$$= a^2bc + ab^2c + abc^2$$

$$= abc(a+b+c). \underline{\text{Ans}}$$

Q. A region of volume 10 cubic units is bounded by a closed surface 'S'. and  $N$  is the outer unit normal vector to S, then -

$$\int_S (4xi + 2yj - zk) \cdot N ds = \underline{\quad}$$

$$\rightarrow \nabla \cdot F = 4 + 2 - 1 = 5$$

$$\int_V 5 dv = 5V = 5 \times 10 = 50.$$

\*\*\*  
✓✓

(ONLY APPLICABLE WHEN R is also in only x-y plane, no z plane)

# GREEN'S THEOREM in a plane :-  $(\int \rightarrow \iint)$

Let 'R' be the closed region in the xy plane bounded by a closed and non intersecting curve 'c'

and Let P and Q are continuous fn of x and y possessing the first order partial derivatives, then

$$\oint_C P dx + Q dy = \iint_R \left( \frac{\partial Q}{\partial x} - \frac{\partial P}{\partial y} \right) dx dy.$$

Q. The value of  $\oint_C (2x-y)dx + (x+y)dy$  where  $C$  is the circle  $x^2+y^2=9$

$$\rightarrow \frac{\partial P}{\partial x} = -1, \quad \frac{\partial Q}{\partial x} = 1.$$

$\rightarrow$  Consto, so  
 no need to consider  
 limit.

Only for  $\square$  and  $\square$   
 we will take const. limit  
 of  $x$  and  $y$ .

for other  
 curves  
 except  
 rectangle  
 & square  
 $\rightarrow$  else where, we will take  
 one limit const and  
 other as a fn of former  
 one.

$$\therefore \iint_R (1 - (-1)) dx dy$$

$$= \int_R 2dR = 2R \cdot \frac{\pi r^2}{3} = 18\pi \text{ Ans}$$

Q.  $\oint_C \overbrace{(x^2+y^2)}^P dx - \overbrace{2xy}^Q dy$ , where  $C$  is the rectangle bounded  
 by  $x=0, x=a$   
 $y=0, y=b$ .

$$\rightarrow \frac{\partial P}{\partial y} = 2y, \quad \frac{\partial Q}{\partial x} = -2y$$

$$\iint_R (-2y - 2y) dx dy$$

$$= \int_{x=0}^a \int_{y=0}^b -4y dx dy$$

$$= (x)_0^a \left(-4 \frac{y^2}{2}\right)_0^b = a(-2b^2) = -2ab^2$$

a.  $\oint_C (x^2 - xy^3) dx + (y - 2xy) dy$  where  $C$  is the square bounded by the points  $(0,0)$   $(2,0)$   $(2,2)$   $(0,2)$

$$\rightarrow \frac{\partial P}{\partial y} = -3xy^2, \quad \frac{\partial Q}{\partial x} = -2y$$

$$\therefore \iint_R -2y - (-3xy^2) dx dy$$

$$= \int_{x=0}^2 \int_{y=0}^2 -2y + 3xy^2 dx dy$$

$$= -2 \left( \frac{y^2}{2} \right)_0^2 (x)_0^2 + 3 \left( \frac{x^2}{2} \right)_0^2 \left( \frac{y^3}{3} \right)_0^2$$

$$= -8 + 16 = 8 \quad \underline{\text{Ans}}$$

a.  $\oint_C (3x^2 - 8y^2) dx + (4y - 6xy) dy$  where  $C$  is the region bounded by  $y = \sqrt{x}$  and  $y = x^2$


$$\rightarrow \frac{\partial P}{\partial y} = -16y, \quad \frac{\partial Q}{\partial x} = -6y$$

$$\int_{x=0}^1 \int_{y=x^2}^{\sqrt{x}} (-6y + 16y) dx dy$$

$$= \int_0^1 10 \left( \frac{y^2}{2} \right)_{x^2}^{\sqrt{x}} dx$$

$$= 5 \int_0^1 (2 - x^4) dx$$

$$= 5 \left[ \frac{x^2}{2} - \frac{x^5}{5} \right]_0^1 = 5 \left( \frac{3}{10} \right) = \frac{3}{2} \quad \underline{\text{Ans}}$$



$$\begin{aligned} x^2 &= \sqrt{x} \\ x^4 &= x \\ x(x^3 - 1) &= 0 \\ x &= 0, 1 \\ y &= 0, 1 \end{aligned}$$

Q.  $\oint_C (3x^2 - 8y^2) dx + (4y - 6xy) dy$  where  $C$  is the region bounded by  $x=0, y=0, x+y=1$

$$\rightarrow \frac{\partial P}{\partial y} = -16y, \quad \frac{\partial Q}{\partial x} = -6y$$

$$\int_{x=0}^1 \int_{y=0}^{1-x} (-6y + 16y) dx dy$$

$$= \int_0^1 \int_{y=0}^{1-x} 10y dx dy$$


$$= \int_0^1 10 \left( \frac{y^2}{2} \right)_0^{1-x} dx$$

$$= 5 \int_0^1 (1-x)^2 dx$$

$$= 5 \left| \frac{(1-x)^3}{-3} \right|_0^1$$

$$= -\frac{5}{3}(0-1)$$

$$= \frac{5}{3} \quad \underline{\underline{\text{Ans}}}$$



# # STOKE'S THEOREM :- $\int \rightarrow \iint$

Let  $S$  be an open surface bounded by a closed and a non intersecting curve 'c' and  $F$  be any differentiable vector function then —

$$\oint_C F \cdot d\vec{r} = \iint_S (\text{curl } F) \cdot N \, ds = \iint_S (\nabla \times F) \cdot N \, ds$$

Q. Evaluate by the Stoke's theorem

$\oint_C F \cdot d\vec{r}$  where  $F = -y^3 i + x^3 j$  and  $S$  is the circular disc  $x^2 + y^2 \leq 1, z=0$ .

$$\rightarrow \nabla \times F = \begin{vmatrix} i & j & k \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ -y^3 & x^3 & 0 \end{vmatrix}$$

$$= i(0-0) - j(0-0) + k(3x^2+3y^2)$$

$$= 3(x^2+y^2)k$$

$$= 3k \quad \left\{ \because x^2+y^2 \leq 1 \right\}$$

Directly we can say  $N = \vec{k}$   $\left\{ \begin{array}{l} \because z=0 \text{ given} \\ \text{means } xy \text{ plane} \\ \text{and normal vector} \\ \text{will be only } k \end{array} \right\}$

$$\therefore \int \vec{k} \cdot \vec{k} \, ds$$


$$= 3 \int ds$$

$$= 3S$$

$$= 3\pi(1)^2$$

$$= 3\pi \underline{3\pi}$$

Q. The area bounded by  $y = x^2$  and  $y = x$  —



$$\begin{aligned} x^2 &= x \\ x(x-1) &= 0 \\ x &= 0, 1 \\ y &= 0, 1 \end{aligned}$$

Prop Trick

The area =  $\int_0^1 (x - x^2) dx$  from Graph

↳ Area will be always upper part - lower part.

$$= \left[ \frac{x^2}{2} - \frac{x^3}{3} \right]_0^1$$

$$= \frac{1}{2} - \frac{1}{3}$$

$$= \frac{1}{6} \text{ Ans}$$

Q. The area bounded by  $y^2 = 4ax$  &  $x^2 = 4ay$  is  $\frac{16a^3}{3}$

The area bounded by  $y^2 = 8x$  &  $x^2 = 8y$  is  $\frac{64}{3}$

\_\_\_\_\_  $y^2 = 4x$  &  $x^2 = 4y$  is  $\frac{16}{3}$

\_\_\_\_\_  $y^2 = x$  &  $x^2 = y$  is  $\frac{1}{3}$

\*\*\*  
Q.  
↑  
Prop Result

The arc length of the curve  $y = f(x)$  between

$$x = a \text{ \& \& } x = b \text{ is } \int_a^b \sqrt{1 + \left(\frac{dy}{dx}\right)^2} dx$$

Let Q.  $y = \frac{2}{3} x^{3/2}$  between  $x = 0$  and  $x = 1$

$$\frac{dy}{dx} = \frac{2}{3} \times \frac{3}{2} x^{3/2-1} = \sqrt{x}$$

$$\begin{aligned} \therefore \int_a^b \sqrt{1 + (\sqrt{x})^2} dx &= \int_a^b \sqrt{1+x} dx = \int_a^b (1+x)^{1/2} dx \\ &= \frac{(1+x)^{3/2}}{3/2} \Big|_0^1 = \frac{2}{3} [2\sqrt{2} - 1] \end{aligned}$$

Q. A parabolic arc  $y = \sqrt{x}$   $1 \leq x \leq 2$  is revolving around  $x$  axis then the volume of the solid revolution.

$$\rightarrow \boxed{\begin{aligned} \text{volume} &= \int_a^b \pi y^2 dx \quad (x\text{-axis}) \\ &= \int_a^b \pi x^2 dy \quad (y\text{-axis}) \end{aligned}}$$

$$\begin{aligned} \therefore \text{volume} &= \int_1^2 \pi (\sqrt{x})^2 dx \\ &= \pi \left( \frac{x^2}{2} \right)_1^2 = \frac{\pi}{2} (4-1) = \frac{3\pi}{2} \quad \underline{\text{Ans}} \end{aligned}$$

WB

(97)

Pg-19

$$\text{volume} = \int_a^b \pi x^2 dy$$

$$y^2 = 8x, \quad x=2$$

$$y^2 = 8(2)$$

$$y^2 = 16$$


$$\Rightarrow y = \pm 4$$

$$\begin{aligned} \therefore \text{volume} &= \int_{-4}^4 \pi x^2 dy \\ &= \frac{\pi}{8^2} \int_{-4}^4 (y^2)^2 dy \\ &= \frac{\pi}{64} \cdot 2 \int_0^4 y^4 dy \\ &= \frac{\pi}{32} \times \frac{y^5}{5} \Big|_0^4 \\ &= \frac{\pi}{32} \times \frac{4 \times 4 \times 4 \times 4 \times 4}{5} = \frac{32\pi}{5} \quad \underline{\text{Ans}} \end{aligned}$$

96

$$\int_0^2 \int_{y=x^2}^{\frac{y}{2}} f(x,y) dy dx$$


$\Rightarrow x = \frac{y}{2}$   
 $y = x^2$   
 $\Rightarrow x = \sqrt{y}$



$$\equiv \int_0^4 \int_{\frac{y}{2}}^{\sqrt{y}} f(x,y) dx dy \quad \underline{\text{Ans}}$$

87

$$\int_0^1 \int_{x=y}^{x=1} xy \sin(xy) dx dy = \int_0^1 \int_a^b xy \sin xy dy dx$$




By changing order of integration

$$= \int_0^1 \int_0^x xy \sin(xy) dy dx$$

88

$$\int_0^8 \int_{y=x/4}^2 f(x,y) dy dx \approx \int_0^8 \int_p^q f(x,y) dx dy$$




$$\approx \int_{y=0}^2 \int_{x=0}^{4y} f(x,y) dx dy$$

85

$$\int_0^1 \int_{x=2y}^{x=2} e^{x^2} dx dy$$

By changing order of integration—



$$\equiv \int_0^2 \int_0^{x/2} e^{x^2} dy dx$$

$$= \int_0^2 e^{x^2} (y)_0^{x/2} dx$$


$$= \int_0^2 e^{x^2} \frac{x}{2} dx$$

$$= \frac{1}{2 \times 2} \int_0^2 e^{x^2} 2x dx$$

$$= \frac{1}{4} (e^{x^2})_0^2 = \frac{e^4 - e^0}{4} = \frac{e^4 - 1}{4} \underline{\underline{Ans}}$$

81

$$\int_0^{\pi/2} \int_0^y \frac{\cos y}{y} dy dx$$



By changing order of integration —

$$\int_0^{\pi/2} \int_0^y \frac{\cos y}{y} dx dy$$

$$= \int_0^{\pi/2} \frac{\cos y}{y} (x)_0^y dy$$

$$= \int_0^{\pi/2} \cos y dy$$

$$= (\sin y)_{\pi/2}^0$$


$$= \sin \frac{\pi}{2} - \sin 0$$

$$= 1 - 0$$

$$= \underline{\underline{1}}$$

~~77~~  
78

$$\int_0^{\infty} \int_x^{\infty} \frac{1}{y} e^{-y/2} dy dx$$




$$\approx \int_0^{\infty} \int_0^y \frac{1}{y} e^{-y/2} dx dy$$

$$= \int_0^{\infty} \int_0^y \frac{1}{y} e^{-y/2} (x)_0^y dy$$

$$= -2 \left[ e^{-y/2} \right]_0^{\infty} = -2 [e^{-\infty} - e^{-0}] = -2(0-1) = 2$$

77

$$\int_0^1 \int_y^1 y \sqrt{1+x^3} dx dy$$



$$\approx \int_0^1 \int_0^x y \sqrt{1+x^3} dy dx$$

$$\approx \int_0^1 \sqrt{1+x^3} \left( \frac{y^2}{2} \right)_0^x dx$$

$$= \int_0^1 \sqrt{1+x^3} \frac{x^2}{2} dx$$

$$= \frac{1}{2} \times \frac{1}{3} \int_0^1 (1+x^3)^{1/2} \cdot 3x^2 dx$$

$$= \frac{1}{3} \left| \frac{(1+x^3)^{3/2}}{\frac{3}{2}} \right|_0^1$$

$$= \frac{1}{9} [2\sqrt{2}-1]$$

41  $f(n) = \int_0^{\pi/4} \tan^n x \, dx$  then  $f(3) + f(1) = ?$

$$\rightarrow f(3) + f(1) = \int_0^{\pi/4} (\tan^3 x + \tan x) \, dx.$$

$$= \int_0^{\pi/4} (\tan x) (\sec^2 x) \, dx$$

$$= \left( \frac{\tan^2 x}{2} \right)_0^{\pi/4} = \frac{1}{2} \underline{Ans}$$

60  $\int_0^{\pi/2} \frac{\cos x - \sin x}{1 + \sin x \cos x} \, dx = ?$

$$\therefore \int_0^{\pi/2} f(x) \, dx = \int_0^a f(a-x) \, dx$$

$$I = \int_0^{\pi/2} \frac{\sin x - \cos x}{1 + \cos x \sin x} \, dx$$

$$2I = \int_0^{\pi/2} \frac{\cos x - \sin x - \sin x - \cos x}{1 + \sin x \cos x}$$

$$= 0$$

$$\Rightarrow I = 0 \quad \underline{Ans}$$

56

$P(x, y, z)$   
 $z^2 = 1 + xy$   
 $O(0,0,0)$

$$OP = \sqrt{x^2 + y^2 + z^2}$$

$$= \sqrt{x^2 + y^2 + 1 + xy}$$

$$f = x^2 + y^2 + xy + 1$$

$$P = \frac{\partial f}{\partial x} = 2x + y = 0$$

$$Q = \frac{\partial f}{\partial y} = 2y + x = 0$$

$$x - y = 0$$

$$x = y$$

~~$$3x = 0$$~~  
~~$$x = 0$$~~

$$3x = 0$$

$$\Rightarrow x = 0$$

$$y = 0$$

Put  $x=y$ .

$$\therefore OP = \sqrt{0+0+0+1}$$

$$= 1$$

53

$x = 2$  after making  $f''(x) = 0$

Since maximum of slope needed not maximum of curve demanding

45

$$x = uv, \quad y = \frac{v}{u}$$

$$f(x, y) dx dy = \iint f(uv, \frac{v}{u}) \phi(u, v) du dv$$

$$\phi(u, v) = \begin{vmatrix} \frac{\partial x}{\partial u} & \frac{\partial x}{\partial v} \\ \frac{\partial y}{\partial u} & \frac{\partial y}{\partial v} \end{vmatrix} = \begin{vmatrix} v & u \\ -\frac{v}{u^2} & \frac{1}{u} \end{vmatrix}$$

$$= \frac{2v}{u}$$

Q. A curve  $c$  is defined as  $x = a \cos^3 \theta$   
 $y = a \sin^3 \theta$  in  $[0, \frac{\pi}{2}]$

What will be point  $P$  on curve  $c$  where tangent to curve is parallel to chord joining points  $(a, 0)$  &  $(0, a)$ .

$$\begin{aligned} \rightarrow & \quad x = a \cos^3 \theta, \quad y = a \sin^3 \theta. \quad \leftarrow \\ & \frac{a \times 3 \sin^2 \theta \times \cos^2 \theta}{a \times 3 \cos^2 \theta \times -\sin^2 \theta} = \frac{a-0}{0-a} \\ \Rightarrow & \quad \tan \theta = \frac{a}{a} \\ \Rightarrow & \quad \theta = \frac{\pi}{4}. \end{aligned}$$

$\therefore$  putting values —

we get —

$$\left( \frac{a}{2\sqrt{2}}, \frac{a}{2\sqrt{2}} \right)$$

Q. 19 — correct  $y = a(1 - \cos \theta)$ .

Q. ⑧.  $\lim_{n \rightarrow \infty} \left( \frac{n!}{n^n} \right)^{\frac{1}{n}}$

$$d = \left( \frac{n!}{n^n} \right)^{\frac{1}{n}}$$

$$\log d = \frac{1}{n} [\log n! - n \log n]$$

~~$$\log d = \frac{1}{n} [\log n! - n \log n]$$~~

$$= \frac{1}{n} \left[ \log \frac{1 \times 2 \times 3 \times \dots \times n}{n \times n \times n \dots \times n} \right]$$

$$= \frac{1}{n} \left[ \log\left(\frac{1}{n}\right) + \log\left(\frac{2}{n}\right) + \dots \right]$$

Imp formula

$$\lim_{n \rightarrow \infty} \left( \frac{1}{n} \sum_{r=1}^n \log\left(\frac{r}{n}\right) \right) = \int_0^1 \log x \, dx$$

$$= [x \log x - x]_0^1$$

$$\log 1 = (1 \log 1 - 1)$$

$$l = e^{-1} = \frac{1}{e} \text{ Ans}$$

$$\text{Q6} \quad \lim_{n \rightarrow \infty} \left[ \frac{1}{1+n} + \frac{1}{2+n} + \dots + \frac{1}{n+n} \right]$$

$$= \lim_{n \rightarrow \infty} \frac{1}{n} \left[ \frac{1}{1+\frac{1}{n}} + \frac{1}{1+\frac{2}{n}} + \dots \right]$$

$$= \lim_{n \rightarrow \infty} \left( \frac{1}{n} \sum_{r=1}^n \frac{1}{1+\left(\frac{r}{n}\right)} \right)$$

$$= \int_0^1 \log \frac{1}{1+x} \, dx$$

$$= \log(1+x) \Big|_0^1$$

$$= \log 2 - \log 1$$

$$= \log 2 \text{ Ans}$$

$$\textcircled{5} \quad \lim_{n \rightarrow \infty} \left[ \frac{n}{n^2} + \frac{n}{n^2+1^2} + \dots + \frac{n}{n^2+(n-1)^2} \right]$$

$$\lim_{n \rightarrow \infty} \frac{n}{n^2} \left[ \sum_{r=0}^{n-1} \frac{1}{1 + \left(\frac{r}{n}\right)^2} \right]$$

$$= \int_0^1 \frac{1}{1+x^2} dx$$

$$= \left( \tan^{-1} x \right)_0^1$$

$$= \tan^{-1} 1 - \tan^{-1} 0$$

$$= \frac{\pi}{4} - 0 = \frac{\pi}{4} \quad \text{Ans}$$

$$\textcircled{7} \quad \lim_{n \rightarrow \infty} \sum_{k=1}^n \frac{1}{3n+k}$$

$$= \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{k=1}^n \frac{1}{3 + \left(\frac{k}{n}\right)}$$

$$= \int_0^1 \frac{1}{3+x} dx = \left[ \log(3+x) \right]_0^1 = \log 4 - \log 3 = \log \frac{4}{3} \quad \text{Ans}$$

## DIFFERENTIAL EQ<sup>n</sup>

An eq<sup>n</sup> consisting of Differential coeff.  $\frac{dy}{dx}$ ,  $\frac{d^2y}{dx^2}$ ,  $\frac{d^3y}{dx^3}$ , ...  $\frac{d^ny}{dx^n}$

is called a differential eq<sup>n</sup>.

\* order -  $\frac{d^ny}{dx^n}$  is order

\* Degree - The degree of diff eq<sup>n</sup> is the degree of the highest order

$$\frac{d^2y}{dx^2} + 5\left(\frac{dy}{dx}\right)^3 + 6y = 0 \quad \begin{array}{l} 0 \rightarrow 2 \\ D \rightarrow 1 \end{array}$$

$$\left(\frac{d^2y}{dx^2}\right)^3 + 5\frac{dy}{dx} + 2y = k\left(\frac{d^3y}{dx^3}\right)^2 \quad \begin{array}{l} 0 \rightarrow 3 \\ D \rightarrow 2 \end{array}$$

$$\left(\frac{d^2y}{dx^2}\right)^2 + \sqrt{1 + \left(\frac{dy}{dx}\right)^2} = 5y \quad \begin{array}{l} D \rightarrow 4 \\ 0 \rightarrow 2 \end{array}$$

$$\left(\frac{d^2y}{dx^2}\right)^3 + 5 \cos\left(\frac{dy}{dx}\right) + 6y = 0 \quad \begin{array}{l} D \rightarrow \text{not defined determined} \\ 0 \rightarrow 2 \end{array}$$

### # Formation of a Differential Eq<sup>n</sup> :-

A diff. eq<sup>n</sup> can be formed by eliminating arbitrary constants in the solution.

Ex:-  $y^2 = 4ax$

$$2y \frac{dy}{dx} = 4a$$

$$y^2 = \frac{2y}{2} \cdot x \cdot \frac{2y}{dx}$$

$$2x \frac{dy}{dx} - y = 0$$

$$0 \quad x^2 + y^2 = a^2$$

$$2x + 2y \frac{dy}{dx} = 0$$

$$\boxed{x + y \frac{dy}{dx} = 0}$$

Note :-

Differentiate only  
that much time  
upto these <sup>no. of</sup> arbitrary  
constants. <sup>^</sup>

$$Q. \quad y = ae^x + be^{-x}$$

$$\frac{dy}{dx} = ae^x - be^{-x}$$

$$\frac{d^2y}{dx^2} = ae^x + be^{-x}$$

$$\boxed{\frac{d^2y}{dx^2} = y}$$

$$Q. \quad y = e^x (a \cos x + b \sin x)$$

$$\frac{dy}{dx} = y_1 = e^x (a \cos x + b \sin x) + e^x (-a \sin x + b \cos x)$$

$$y_1 - y = e^x (-a \sin x + b \cos x)$$

$$y_2 - y_1 = \underbrace{e^x [-a \sin x + b \cos x]}_{y_1 - y} + \underbrace{e^x [-a \cos x + b \sin x]}_{-y}$$

$$y_2 - y_1 = y_1 - y - y$$

$$\frac{d^2y}{dx^2} - 2\frac{dy}{dx} + 2y = 0$$

# Solution of a Differential Eq<sup>n</sup> :-

\* Variable Seperable Method :-

$$\textcircled{1} \quad \frac{dy}{dx} + \sqrt{\frac{1+y^2}{1+x^2}} = 0$$

$$\frac{dy}{dx} = -\frac{\sqrt{1+y^2}}{\sqrt{1+x^2}}$$

$$\int \frac{dy}{\sqrt{1+y^2}} = \int -\frac{dx}{\sqrt{1+x^2}}$$

$$\Rightarrow \boxed{\sinh^{-1} y = -\sinh^{-1} x + c}$$

$$\textcircled{2} \quad \frac{dy}{dx} = e^{x+y}$$

$$\frac{dy}{dx} = e^x \cdot e^y$$

$$\int dy e^{-y} = \int e^x dx$$

$$-e^{-y} = e^x + c$$

$$\Rightarrow e^x + e^{-y} + c = 0$$

$$\textcircled{3} \quad \frac{dy}{dx} = x^2 e^{x-y}$$

$$\int \frac{dy}{e^y} = \int x^2 e^x dx$$

$$e^y = e^x [x^2 - 2x + 2] + c$$

~~e<sup>y</sup>~~

$$\textcircled{4} \quad \frac{dy}{dx} + y^2 = 0$$

$$\int dy y^{-2} = -\int dx$$

$$\frac{y^{-2+1}}{-1} = -x + C$$

$$\Rightarrow -y^{-1} = -x + C$$

$$\Rightarrow y^{-1} + x + C = 0 \quad \text{or} \quad \boxed{y = \frac{1}{x+C}}$$

$$\textcircled{5} \quad e^x \tan y dx + (1+e^x) \sec^2 y dy = 0$$

$$e^x \tan y dx = -(1+e^x) \sec^2 y dy$$

$$\frac{e^x dx}{1+e^x} = -\frac{\sec^2 y dy}{\tan y}$$

$$\int \frac{e^x}{1+e^x} dx + \int \frac{\sec^2 y}{\tan y} dy = \int 0$$

$$\log(1+e^x) + \log \tan y = \log C$$

$$\Rightarrow \boxed{(1+e^x) \tan y = C}$$

⑥

## II<sup>nd</sup> Method

\* Eq<sup>n</sup> Reducible to variable-seperable form:-

If the given diff. eq<sup>n</sup> is not directly variable seperable form, by making substitution, again we can transform and under variable seperable form.

$$Q. \quad \frac{dy}{dx} = (4x+y+1)^2$$

$$\hookrightarrow \text{Let } z = 4x+y+1$$

$$\frac{dz}{dx} = 4 + \frac{dy}{dx}$$

$$4 + \frac{dy}{dx} = 4 + (4x+y+1)^2$$

$$\Rightarrow \frac{dz}{dx} = 4 + z^2$$

$$\Rightarrow \int \frac{dz}{z^2+4} = \int dx$$

$$\Rightarrow \frac{1}{2} \tan^{-1} \frac{z}{2} = x + c$$

$$\Rightarrow \boxed{\frac{1}{2} \tan^{-1} \left( \frac{4x+y+1}{2} \right) = x + c}$$

$$Q. \quad 1 + \frac{dy}{dx} = 1 + \sin(x+y) + \cos(x+y)$$

$$\frac{dz}{dx} = 1 + \sin z + \cos z$$

$$x+y = z$$

$$1 + \frac{dy}{dx} = \frac{dz}{dx}$$

$$\Rightarrow \int \frac{dz}{1 + \sin z + \cos z} = \int dx$$

Remember

WHENEVER INTEGRAL IS -

in form of  $\frac{1}{a+b \sin x}$

$\int \frac{1}{a+b \cos x}$

$\int \frac{1}{a \cos x + b \sin x} dx$

$$\text{Put } t = \tan \frac{x}{2} \Rightarrow dx = \frac{2dt}{1+t^2}$$

$$\sin x = \frac{2t}{1+t^2}$$

$$\cos x = \frac{1-t^2}{1+t^2}$$

$$\int \frac{1}{1 + \frac{2t}{1+t^2} + \frac{1-t^2}{1+t^2}} \times \frac{2dt}{1+t^2} = x + c$$

$$\int \frac{2dt}{(1+t^2+2t+1-t^2)} = x + c$$

$$\int \frac{2}{2(1+t)} dt = x + c$$

$$\log(1+t) = x + c$$

$$\log\left[1 + \tan \frac{x+y}{2}\right] = x + c$$

## # HOMOGENEOUS DIFFERENTIABLE EQUATION:-

A diff. eq<sup>n</sup> of the form -

$\frac{dy}{dx} = \frac{f(x,y)}{g(x,y)}$  is said to be homogeneous differential

eq<sup>n</sup>, if both  $f(x,y)$  and  $g(x,y)$  are homogeneous functions of the same degree.

$$y = vx \Rightarrow v = \frac{y}{x}$$

$$\boxed{\frac{dy}{dx} = v + x \frac{dv}{dx}}$$

When  $\frac{dx}{dy} = \frac{f(x,y)}{g(x,y)}$

$$x = vy \Rightarrow v = \frac{x}{y}$$

$$\boxed{\frac{dx}{dy} = v + y \frac{dv}{dy}}$$

### ILATE

Ex:-  $\frac{dy}{dx} = \frac{x^2+y^2}{2xy}$

$$\frac{dx}{dy} = \frac{2xy}{x^2+y}$$

$$\therefore \frac{dy}{dx} = \frac{x^2+y^2}{2xy}$$

$$v + x \frac{dv}{dx} = \frac{\cancel{x^2} + v^2 \cancel{x^2}}{2x \cdot v \cancel{x}}$$

$$x \frac{dv}{dx} = \frac{1+v^2}{2v} - v$$

$$x \frac{dv}{dx} = \frac{1-v^2}{2v}$$

→ which form should be selected for substitution?

↳ निम्ने numerator में प्सादा no. of terms रोगा

$$\int \frac{-2v}{1-v^2} dv = \int \frac{-dx}{x}$$

$$\Rightarrow \log(1-v^2) = \log \frac{C}{x}$$

$$\Rightarrow 1 - \frac{y^2}{x^2} = \frac{C}{x}$$

$$\Rightarrow \frac{x^2 - y^2}{x^2} = \frac{C}{x}$$

$$\Rightarrow \boxed{x^2 - y^2 = Cx}$$

$$\frac{1}{x} \rightarrow \log Cx$$

$$-\frac{1}{x} \rightarrow \log \frac{C}{x}$$

$$\frac{2}{x} \rightarrow \log Cx^2$$

$$-\frac{2}{x} \rightarrow \log \frac{C}{x^2}$$

$$\frac{3}{x} \rightarrow \log Cx^3$$

$$-\frac{3}{x} \rightarrow \log \frac{C}{x^3}$$

Q.  $x^2y dx - (x^3 + y^3) dy = 0$ .

$$\frac{dy}{dx} = \frac{x^2y}{x^3 + y^3}$$

$$\Rightarrow \frac{dx}{dy} = \frac{x^3 + y^3}{x^2y}$$

$$v + y \frac{dv}{dy} = \frac{v^3 \cancel{y^3} + y^3}{v^2 y^3}$$

$$v + y \frac{dv}{dy} = \frac{v^3 + 1}{v^2}$$

$$y \frac{dv}{dy} = \frac{1}{v^2}$$

$$\Rightarrow \int \frac{dy}{y} = \int v^2 dv$$

$$\Rightarrow \log cy = \frac{v^3}{3}$$

$$\Rightarrow \log cy = \frac{x^3}{3y^3} \quad \underline{\underline{Ans}}$$

$$Q. \quad x \frac{dy}{dx} = y + x \tan\left(\frac{y}{x}\right)$$

$$\frac{dy}{dx} = \frac{y}{x} + \tan \frac{y}{x}$$

$$y + x \frac{dv}{dx} = y + \tan v$$

$$\int \frac{dv}{\tan v} = \int \frac{dx}{x}$$

$$\log \sin v = \log Cx$$

$$\boxed{\sin\left(\frac{y}{x}\right) = Cx}$$

INDIAN PHOTOSTAT & BOOK CENTRE  
 40D WESTEND MARKET, SHRIJEEVARAB  
 NEW DELHI-110030  
 Mob: 9560127702

$$Q. \quad (1 + e^{xy}) dx + e^{xy} \left(1 - \frac{x}{y}\right) dy = 0$$

$$\frac{dx}{dy} = \frac{e^{xy} \left(\frac{x}{y} - 1\right)}{1 + e^{xy}}$$

$$v + y \frac{dv}{dy} = \frac{e^v (v-1)}{1 + e^v}$$

$$y \frac{dv}{dy} = \frac{e^v (v-1)}{1 + e^v} - v$$

$$y \frac{dv}{dy} = \frac{v/e^v - e^v - v - v/e^v}{1 + e^v}$$

$$y \frac{dv}{dy} = \frac{-(v + e^v)}{1 + e^v}$$

$$\int \frac{(1 + e^v)}{v + e^v} dv = \int -\frac{dy}{y}$$

$$\log(v + e^v) = \log \frac{C}{y}$$

$$\Rightarrow \frac{x}{y} + e^{xy} = \frac{C}{y} \Rightarrow \boxed{x + y \cdot e^{xy} = C}$$

Triq → Angle विलीन  
 Algebra → उचित power  
 numerator में

exponents  $\frac{x}{y} - \frac{dx}{dy}$   
 $\frac{y}{x} - \frac{dy}{y}$

# # NON HOMOGENEOUS DIFFERENTIAL FUNCTION:-

A diff. eq<sup>n</sup> of the form —

$$\frac{dy}{dx} = \frac{a_1x' + b_1y' + c_1}{a_2x' + b_2y' + c_2} \text{ is said to be a non}$$

homogeneous differential eq<sup>n</sup>

Case-1:- IF  $a_1b_2 - a_2b_1 \neq 0$ , then the substitution is —

$$x = X + h$$

$$y = Y + k$$

where  $h, k$  are constants to be determined.

Case-2:- IF  $a_1b_2 - a_2b_1 = 0$ , then ~~the~~ some part of numerator and denominator are equal and that part is considered as substitution.

Q. Find the substitution that transforms a non homogeneous diff eq<sup>n</sup>  $\frac{dy}{dx} = \frac{2x+2y-2}{3x+y-5}$  to homogeneous form.

$\hookrightarrow \neq 0 \rightarrow 1^{st} \text{ form}$

$\rightarrow$

$$\frac{dy}{dx} = \frac{2(x+h) + 2(Y+k) - 2}{3(x+h) + (Y+k) - 5}$$

$h$	$k$	$1$	$h$	$k$	
2	2	-2	2	2	$\rightarrow$ numerators
3	1	-5	3	1	$\rightarrow$ Denominators

$\rightarrow$  write like this only

$$\frac{h}{-10+2} = \frac{k}{-6+10} = \frac{1}{2-6}$$

$$\frac{h}{-8} = \frac{k}{4} = \frac{1}{-4} \quad \checkmark$$

$$h = 2, k = -1 \quad \checkmark$$

$$\Rightarrow \left. \begin{aligned} x &= X + 2 \\ y &= Y - 1 \end{aligned} \right\} \quad \checkmark$$

Q. Find Substitution —

$$\frac{dy}{dx} = \frac{x+2y-3}{2x+y-3}$$

$\hookrightarrow \neq 0 \rightarrow 1^{\text{st}} \text{ form}$

h	k	1	h	k
1	2	-3	1	2
2	1	-3	2	1

$$\frac{h}{-6+3} = \frac{k}{-6+3} = \frac{1}{1-4}$$

$$\frac{h}{-3} = \frac{k}{-3} = \frac{1}{-3}$$

$$h = 1, k = 1 \quad \leftarrow$$

$$\begin{aligned} x &= X+1 \\ y &= Y+1 \end{aligned}$$

Ans.

Q. Find the Substitution —

$$\frac{dy}{dx} = \frac{2x+3y+1}{4x+6y+1}$$

$\hookrightarrow = 0 \rightarrow 2^{\text{nd}} \text{ form}$

$$\frac{dy}{dx} = \frac{(2x+3y)+1}{2(2x+3y)+1}$$

$$z = 2x+3y$$

$$0. \quad \frac{dy}{dx} = \frac{x+2y+3}{y+x+5}$$

$$\frac{dy}{dx} = \frac{x+2y+3}{2x+y+5}$$

↳  $\neq 0 \rightarrow$  1<sup>st</sup> form.

h	k	l	h	k
1	2	3	1	2
2	1	5	2	1

$$\frac{h}{10-3} = \frac{k}{6-5} = \frac{l}{1-4}$$

$$\frac{h}{7} = \frac{k}{1} = \frac{l}{-3}$$

$$h = -\frac{7}{3}, \quad k = -\frac{1}{3}$$

$$\boxed{\begin{aligned} x &= X + \frac{7}{3} \\ y &= Y - \frac{1}{3} \end{aligned}}$$

Ans

## # EXACT DIFFERENTIAL EQN:-

A diff. eqn of the form  $Mdx + Ndy = 0$  is said to be exact if

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

$$\text{Gen. Sol}^n \rightarrow \int^x M dx + \int (\text{terms of } N \text{ not containing } x) dy = c$$

Ex:-  $(x^2 - ay)dx + (y^2 - ax)dy = 0$

$$\frac{\partial M}{\partial y} = -a \quad , \quad \frac{\partial N}{\partial x} = -a$$

exact

$$\int (x^2 - ay) dx + \int y^2 dy = c$$

$$\frac{x^3}{3} - axy + \frac{y^3}{3} = c$$

$$\Rightarrow x^3 + y^3 - 3axy = 3c \quad \text{Ans}$$

MORE  
Efficient  
Shortcut

### GENERALIZATION:-

Ex:-  $(ax + by + g)dx + (hx + ky + f)dy = 0$

$$\frac{\partial M}{\partial y} = h \quad , \quad \frac{\partial N}{\partial x} = h$$

$$\frac{ax^2}{2} + hxy + gx + \frac{by^2}{2} + fy = c$$

Ans

a.  $e^x + \tan y dx + (1 + e^x) \sec^2 y dy = 0$

$$\frac{\partial M}{\partial y} = e^x \sec^2 y \quad , \quad \frac{\partial N}{\partial x} = e^x \sec^2 y$$

exact

Sol<sup>n</sup>

$$\therefore e^x + \tan y + \tan y = c$$

$$\Rightarrow (e^x + 1) \tan y = c$$

$$Q. (x^2 + y^2) dx + 2xy dy = 0$$

$$\frac{\partial M}{\partial y} = 2y \quad , \quad \frac{\partial N}{\partial x} = 2y$$

Sol<sup>n</sup> ————— exact —————

$$\boxed{\frac{x^3}{3} + y^2 x = C}$$

$$Q. (1 + e^{\frac{x}{y}}) dx + e^{\frac{x}{y}} (1 - \frac{x}{y}) dy = 0$$

$$\text{exact} \left( \begin{array}{l} \frac{\partial M}{\partial y} = e^{\frac{x}{y}} \cdot x \cdot \left(-\frac{1}{y^2}\right) \\ \frac{\partial N}{\partial x} = e^{\frac{x}{y}} \left(\frac{1}{y}\right) + \left(1 - \frac{x}{y}\right) e^{\frac{x}{y}} \left(\frac{1}{y}\right) \\ = e^{\frac{x}{y}} \left(-\frac{1}{y} + \frac{1}{y} - \frac{x}{y^2}\right) \end{array} \right)$$

Sol<sup>n</sup>

$$\boxed{x + ye^{\frac{x}{y}} = C}$$

## # Non-Exact Differential Eq<sup>n</sup>:-

A diff eq<sup>n</sup>  $Mdx + Ndy = 0$  is said to be non exact,  
if  $\frac{\partial M}{\partial y} \neq \frac{\partial N}{\partial x}$

METHOD-1 (Method to form a suitable Integrating factor)

IMPORTANT INTEGRATING FACTORS:-

- \*\*\*
- ①  $x dy + y dx = d(xy)$
  - ②  $\frac{x dy - y dx}{x^2} = d\left(\frac{y}{x}\right)$
  - ③  $\frac{x dy - y dx}{xy} = d\left(\log \frac{y}{x}\right)$
  - ④  $\frac{x dy - y dx}{x^2 + y^2} = d\left(\tan^{-1} \frac{y}{x}\right)$
  - ⑤  $\frac{y dx - x dy}{y^2} = d\left(\frac{x}{y}\right)$
  - ⑥  $\frac{y dx - x dy}{xy} = d\left(\log \frac{x}{y}\right)$
  - ⑦  $\frac{y dx - x dy}{x^2 + y^2} = d\left(\tan^{-1} \frac{x}{y}\right)$
  - ⑧  $\frac{y e^x dx - e^x dy}{y^2} = d\left(\frac{e^x}{y}\right)$
  - ⑨  $\frac{2xy dx - x^2 dy}{y^2} = d\left(\frac{x^2}{y}\right)$
  - ⑩  $\frac{x dx + y dy}{x^2 + y^2} = d\left[\frac{1}{2} \log(x^2 + y^2)\right]$

Q.  $y(1+xy)dx + x(1-xy)dy = 0.$

$$(y+xy^2)dx + (x-x^2y)dy = 0.$$

$$\frac{\partial M}{\partial y} = 1+2xy, \quad \frac{\partial N}{\partial x} = 1-2xy$$

Non exact

→ we can make it perfect IF but considering 1st term carefully

$$y dx + xy^2 dx + xdy - x^2y dy = 0$$

Perfect IF

$$\frac{d(xy)}{(xy)^2} + \cancel{xy} \frac{[y dx - x dy]}{(xy)^2} = \frac{0}{(xy)^2}$$

$$\Rightarrow \int \frac{1}{(xy)^2} dx + \int d(\log \frac{x}{y}) = 0$$

$$\Rightarrow \boxed{-\frac{1}{xy} + \log \frac{x}{y} = C}$$

∴  $\frac{dx}{x^2} = -\frac{1}{x}$

Q ①  $y dx - x dy + (1+x^2)dx + x^2 \sin y dy = 0.$

by taking common it is also integrable

$$\frac{y dx - x dy}{x^2} + \frac{(1+x^2) dx}{x^2} + \frac{x^2 \sin y dy}{x^2}$$

$$\Rightarrow - \int \frac{(x dy - y dx)}{x^2} + \int \left(\frac{1}{x^2} + 1\right) dx + \int \sin y dy = 0$$

$$\Rightarrow - \int d\left(\frac{y}{x}\right) + \int \left(\frac{1}{x^2} + 1\right) dx + \int \sin y dy = 0$$

$$\Rightarrow \boxed{-\frac{y}{x} - \frac{1}{x} + x - \cos y = C}$$

$$a. \int \frac{axy^2 dx}{y^2} + \int \frac{ye^x dx - e^x dy}{y^2} = \int 0$$

$$\Rightarrow \int ax dx + \int d\left(\frac{e^x}{y}\right) = \int 0$$

$$\Rightarrow \boxed{\frac{ax^2}{2} + \frac{e^x}{y} = c} \quad \checkmark$$

$$a. (y^2 e^x + 2xy) dx - x^2 dy$$

$$\Rightarrow \frac{y^2 e^x dx + 2xy dx}{y^2} - x^2 dy = 0$$

$$\Rightarrow \int e^x dx + \int d\left(\frac{x^2}{y}\right) = \int 0$$

$$\Rightarrow \boxed{e^x + \frac{x^2}{y} = c} \quad \checkmark$$

$$a. \frac{2x^2 y^2 dx}{y^2} + \frac{ye^x dx - e^x dy}{y^2} - \frac{y^3 dy}{y^2} = 0$$

$$\Rightarrow \int 2x^2 dx + \int d\left(\frac{e^x}{y}\right) - \int y dy = \int 0$$

$$\Rightarrow \boxed{\frac{2x^3}{3} + \frac{e^x}{y} - \frac{y^2}{2} = c} \quad \checkmark$$

NON EXACT DIFFERENTIAL EQ<sup>n</sup> 1-

METHOD-2

If  $Mdx + Ndy = 0$  is non exact, but homogeneous and if

~~$Mx + Ny \neq 0$~~  then  $\frac{1}{Mx+Ny}$  is an I.F.

After multiplying in diff. eq<sup>n</sup>  $\rightarrow$  non exact becomes exact.

Then for sol<sup>n</sup> will follow procedure of Exact diff. eq<sup>n</sup>.

$d=2$  <sup>Homogeneous</sup>  $d=2$

$\rightarrow$  Q. ①  $(x^2+y^2)dx - 2xydy = 0$

$$\frac{\partial M}{\partial y} = +2y \quad , \quad \frac{\partial N}{\partial x} = -2y$$

$$\begin{aligned} \text{I.F.} &= \frac{1}{Mx+Ny} = \frac{1}{x(x^2+y^2) + y(-2xy)} = \frac{1}{x^3 + xy^2 - 2xy^2} \\ &= \frac{1}{x^3 - xy^2} \\ &= \frac{1}{x(x^2 - y^2)} \end{aligned}$$

Now multiply by I.F. —

$$\int \frac{x^2+y^2}{x(x^2-y^2)} dx - \frac{2xy}{x(x^2-y^2)} dy = 0.$$

$$\Rightarrow \int \left[ \frac{2x}{x^2-y^2} - \frac{1}{x} \right] dx = \log C$$

$$\Rightarrow \cancel{\log} \frac{x^2-y^2}{x} = \log C$$

$$\Rightarrow \boxed{x^2-y^2 = Cx} \quad \checkmark$$

Calculate I.F. —

⑥

$$\textcircled{1} \quad x^2 y dx - (x^3 + y^3) dy = 0.$$

$$\frac{\partial M}{\partial y} = 0 \quad Mx + Ny \neq 0 \quad \leftarrow$$

$$\therefore x^3 y + x^3 y + y^4 \neq 0$$

$$\therefore \frac{1}{Mx + Ny} = -\frac{1}{y^4}$$

Note:-

In Exam option will be provided as  $\frac{1}{y^4}$

coz,  $(-)$  and const will not effect integration and ~~by~~ in process of solving  $(-)$  or const can be taken out common and can be send to right where  $0$  is given.

$$\textcircled{2} \quad (x^2 y - xy^2) dx - (x^3 - 2x^2 y) dy$$

$$\begin{aligned} Mx + Ny &= x^3 y - x^2 y^2 - x^3 y + 2x^3 y^2 \\ &= x^2 y^2 \end{aligned}$$

$$\text{I.F.} = \frac{1}{Mx + Ny} = \frac{1}{x^2 y^2}$$

$$\textcircled{3} \quad xy dx - (x^2 + 2y^2) dy = 0.$$

$$Mx + Ny = x^3 y - x^3 y - 2y^3$$

$$\therefore \text{I.F.} = \frac{1}{Mx + Ny} = \frac{1}{-2y^3} \approx \frac{1}{y^3} \quad \underline{\underline{\text{Ans}}}$$

no importance

# NON EXACT DIFFERENTIAL EQ<sup>N</sup>:-

## METHOD-3:-

If  $Mdx + Ndy = 0$  is non exact, but is in the form of

$$y \frac{f(xy)}{x} dx + x \frac{g(xy)}{y} dy = 0 \text{ and if}$$

$$Mx - Ny \neq 0$$

then  $\frac{1}{Mx - Ny}$  is an I.F. =

$$\Rightarrow y(1+xy)dx + x(1-xy)dy = 0$$

$$Mx - Ny = xy + x^2y^2$$

$$\therefore \frac{1}{Mx - Ny} = \frac{1}{xy(1+xy)}$$

$$Q 1) y(x^2y^2 + xy + 1)dx + x(x^2y^2 - xy + 1)dy = 0$$

for fast calculation of  $\frac{1}{Mx - Ny}$  - cancel ~~at~~ terms having same ~~to~~ order and some const. and increase power or multiply by  $\frac{1}{x} \times \frac{1}{y}$

$$\therefore \frac{1}{Mx - Ny} = \frac{1}{x^2y^2}$$

$$Q 2) y(x^3y^3 + x^2y^2 + xy + 1)dx + x(x^3y^3 - x^2y^2 - xy + 1)dy = 0$$

$$\frac{1}{My - Nx} = \frac{1}{x^3y^3 + x^2y^2}$$

$$Q 3) y(x^2y^2 + 1)dx + x(1 - 2x^2y^2)dy = 0$$

$$\frac{1}{My - Nx} = \frac{1}{x^3y^3}$$

$$Q 4) y(x^2y^2 \cos xy + \cos xy)dx + x[x^2y^2 \sin xy - \cos xy]dy = 0$$

$$\frac{1}{xy \cos xy}$$



$$(2) \quad (y^4 + 2y) dx + (xy^3 + 2y^4 - 4x) dy = 0$$

$$\frac{\partial M}{\partial y} = 4y^3 + 2$$

$$\frac{\partial N}{\partial x} = y^3 - 4$$

2 terms  
 $\therefore$  divisible by  $M$   
 as it has also  
 2 terms

$$\frac{1}{M} \left( \frac{\partial N}{\partial x} - \frac{\partial M}{\partial y} \right)$$

$$= \frac{1}{y^4 + 2y} \left( \cancel{4y^3 + 2} y^3 - 4 - 4y^3 + 2 \right)$$

$$= \frac{1}{y^4 + 2y} \left( -3y^3 - 2 \right)$$

$$= \frac{-3}{y} \rightarrow I.F. = \frac{1}{y^3}$$

$$(3) \quad \left( y + \frac{y^3}{3} + \frac{x^2}{2} \right) dx + \left( \frac{x + xy^2}{4} \right) dy = 0$$

$$\frac{\partial M}{\partial y} = 1 + \frac{xy^2}{3}$$

$$\frac{\partial N}{\partial x} = \frac{1}{4} + \frac{y^2}{4}$$

$$\frac{1}{M} \left( \frac{\partial M}{\partial y} - \frac{\partial N}{\partial x} \right)$$

$$= \frac{4}{x + xy^2} \left( 1 + y^2 - \frac{1}{4} - \frac{y^2}{4} \right)$$

$$= \frac{4}{x(1+y^2)} \cdot 3 \left( \frac{3}{4} + \frac{1}{4} y^2 \right)$$

$$= \frac{3}{x} \rightarrow I.F. = \frac{x^3}{x}$$

$$a. (3x^2y^4 + 2xy) dx + (2x^3y^3 - x^2) dy = 0.$$

$$\frac{\partial M}{\partial y} = 12x^2y^3 + 2x$$

$$\frac{\partial N}{\partial x} = 6x^2y^3 - 2x$$

$\xrightarrow{\quad}$  बीच में + sign आर एट दट  
 divisible by ~~M~~ not  $M'$

$$\frac{1}{M} \left( \frac{\partial M}{\partial x} - \frac{\partial M}{\partial y} \right)$$

$$= \frac{1}{3x^2y^4 + 2xy} (-6x^2y^3 + 4x)$$

$$= -\frac{2}{y} \rightarrow I.F \rightarrow \frac{1}{y^2}$$

## # Linear Eq<sup>n</sup>:-

$\frac{dy}{dx} + Py = Q$ , Where P and Q are said to be a <sup>fn of x, called</sup>

Linear eq<sup>n</sup>.

$$I.F = e^{\int P dx}$$

General sol<sup>n</sup> -

$$y \underset{I.F}{e^{\int P dx}} = \int (Q \cdot \underset{I.F}{e^{\int P dx}}) dx + C$$

Ex:-  
→ ①

$\frac{dx}{dy} + Px = Q$ , Where P and Q are said to be fn. of y

$$I.F = e^{\int P dy}$$

General sol<sup>n</sup> -

$$x e^{\int P dy} = \int (Q e^{\int P dy}) dy + C$$

①  $\frac{dy}{dx} + \frac{y}{x} = \frac{\log x}{x}$

$$P = \frac{1}{x}$$

$$Q = \frac{\log x}{x}$$

$$I.F = e^{\int P dx} = e^{\int \frac{1}{x} dx} = x$$

Sol<sup>n</sup> -

$$y \cdot x = \int \log x dx + C$$

$$\Rightarrow yx = x \log x - x + C \quad \underline{\text{Ans}}$$

②

$$(2-x^2) \frac{dy}{dx} - 2xy = x^2$$

$$\frac{dy}{dx} = \frac{2xy}{2-x^2} = \frac{x^2}{2-x^2}$$

Integrating factor (I.F.) =  $e^{\int P dx} = e^{\int \frac{2x}{2-x^2} dx} = e^{\log(2-x^2)}$   
 $= 2-x^2$

G. Sol<sup>n</sup> -

$$y(2-x^2) = \int \frac{x^2}{2-x^2} (2-x^2) dx + C$$

$$= \frac{x^3}{3} + C$$

③

$$\frac{dy}{dx} + 2xy = e^{-x^2}, \quad y(0) = 1$$

I.F. =  $e^{\int P dx} = e^{\int 2x dx} = e^{x^2} = e^{x^2}$

G. Sol<sup>n</sup> -

$$y e^{x^2} = \int e^{-x^2} \cdot e^{x^2} dx + C$$

$$\boxed{y e^{x^2} = x + C}$$

When  $y(0) = 1$

$$1 \cdot e^0 = C \Rightarrow C = 1$$

$$\therefore \boxed{y e^{x^2} = x + 1}$$

$$\textcircled{4} \quad \frac{dy}{dx} + \frac{1}{x \log x} y = \frac{2 \log x}{x \log x} \quad \text{Q}$$

$$\text{I.F} = e^{\int \frac{1}{x \log x} dx} = e^{\int \frac{1}{\log x} \cdot \frac{1}{x} dx} = e^{\log(\log x)} = \log x$$

G. Sol<sup>n</sup>

$$y \log x = \int \frac{2 \log x}{x \log x} \times \cancel{\log x} dx + C$$

$$y \log x = \int \frac{2 (\log x)^2}{x} dx + C$$

$$y \log x = 2 \log x$$

$$\boxed{y = \log x + \frac{C}{\log x}}$$

$$\textcircled{5} \quad (x + 2y^3) \frac{dy}{dx} = y$$

$$\frac{dx}{dy} = \frac{x}{y} + \frac{2y^5 - 2}{y}$$

$$\frac{dx}{dy} - \frac{x}{y} = 2y^2 \quad \text{Q}$$

$$\text{I.F} = e^{\int -\frac{1}{y} dy} = \frac{1}{y}$$

G. Sol<sup>n</sup>

$$x \cdot \frac{1}{y} = \int 2y^2 \times \frac{1}{y} dy + C$$

$$\frac{x}{y} = y^2 + C$$

$$\Rightarrow \boxed{x = y^3 + Cy}$$

$$\text{Q 5 } (1+y^2)dx + (x - e^{\tan^{-1}y}) dy = 0$$

$$(1+y^2)dx = (e^{\tan^{-1}y} - x) dy$$

$$\Rightarrow \frac{dx}{dy} = \frac{e^{\tan^{-1}y}}{1+y^2} - \frac{x}{1+y^2}$$

$$\Rightarrow \frac{dx}{dy} + \left(\frac{x}{1+y^2}\right) = \frac{e^{\tan^{-1}y}}{1+y^2} \quad \text{Q}$$

$$\therefore \text{I.F} = e^{\int P dy} = e^{\int \frac{1}{1+y^2} dy} = e^{\tan^{-1}y}$$

General Sol<sup>n</sup>—

$$x \cdot e^{\tan^{-1}y} = \int \frac{e^{\tan^{-1}y}}{1+y^2} \cdot e^{\tan^{-1}y} dy + C.$$

$$= \frac{1}{2} \int \frac{e^{2 \tan^{-1}y}}{1+y^2} dy + C$$

$$\boxed{x e^{\tan^{-1}y} = \frac{1}{2} e^{2 \tan^{-1}y} + C.}$$

FOR PHOTOS AT A BOOK CENTER  
ADDITIONAL MADE EASY LANE NO. 4  
WESTEND WARD, SURULAIAB  
NEW DELHI-110038  
MOB: 9560127202

# # BERNOULLI'S EQ<sup>n</sup> :-

A diff. eq<sup>n</sup> of the form  $\frac{dy}{dx} + Py = Q \cdot y^n$ , where

P and Q are fn. of x if said to be a Bernoulli's eq<sup>n</sup>

$$\therefore y^{-n} \frac{dy}{dx} + P y^{1-n} = Q$$

$$\Rightarrow z = y^{1-n}$$

$$\frac{dz}{dx} = (1-n)y^{-n} \frac{dy}{dx}$$

$$\frac{dy}{dx} \rightarrow \frac{dz}{dx}$$

$$y^{-n} \frac{dy}{dx} = \frac{1}{(1-n)} \frac{dz}{dx}$$

$$\frac{1}{(1-n)} \frac{dz}{dx} + Pz = Q$$

$$\Rightarrow \frac{dz}{dx} + P(1-n)z = Q(1-n)$$

$$* \text{ I.F} = e^{\int P(1-n) dx}$$

Gen. sol<sup>n</sup>.

$$* y^{1-n} \cdot \underset{\text{I.F}}{e^{\int P(1-n) dx}} = \int (Q(1-n) \cdot \underset{\text{I.F}}{e^{\int P(1-n) dx}}) dx + C$$

$$\frac{dx}{dy} + Px = Q^n$$

$$\text{I.F} = e^{\int P(1-n) dy}$$

Gen sol<sup>n</sup>

$$x^{1-n} \times \text{I.F} = \int (Q(1-n) \times \text{I.F}) dy + C$$

Find I.F.

$$\textcircled{1} \frac{dy}{dx} + \frac{y}{x} = x^2 y^3$$

$$I.F = e^{\int \frac{1}{x} (1-n) dx} = e^{(1-n) \log x} = x^{1-n} = x^{-2} = \frac{1}{x^2}$$

$$\textcircled{2} \frac{dy}{dx} (xy + x^2 y^3) = 1$$

~~$$\Rightarrow \frac{dy}{dx} - xy = x^2 y^3$$~~

~~$$I.F = e^{\int \frac{1}{x} (1-n) dx} = e^{\frac{x^2}{2} - nx} = e^{-2x}$$~~

$$\Rightarrow \frac{dx}{dy} - xy = y^3 x^{2n}$$

$$I.F = e^{\int -y (1-n) dy} = e^{\int -y (-2) dy} = e^{\int +y dy} = e^{\frac{y^2}{2}}$$

# Differential Eq<sup>n</sup> with constant coefficient :-

A diff eq<sup>n</sup> of the form -

$$a_0 \frac{d^n y}{dx^n} + a_1 \frac{d^{n-1} y}{dx^{n-1}} + a_2 \frac{d^{n-2} y}{dx^{n-2}} + \dots + a_n y = Q.$$

Where  $a_0, a_1, a_2, \dots, a_n$  are constants and  $Q$  be the function of  $x$  is said to be differential equation with constant coefficient.

$$\text{Let } \frac{d}{dx} = D$$

$$(a_0 D^n + a_1 D^{n-1} + a_2 D^{n-2} + \dots + a_n) y = Q.$$

$$\boxed{f(D) y = Q}$$

This eq<sup>n</sup> may contain two types of sol<sup>n</sup>

- one is called complementary fn
- other is called particular integral.

Note:-

If  $Q=0$ , then complementary fn is called the general sol<sup>n</sup>  
if  $Q \neq 0$ , then only particular integral must be existed.

# To Find Complimentary fn (C.F) :-

Replace 'D' by m,  $f(m)=0$  is called auxiliary eq<sup>n</sup>

$$a_0 m^n + a_1 m^{n-1} + a_2 m^{n-2} + \dots + a_n = 0.$$

Case-1.

Roots are real and Unequal.

Let  $m = \alpha, \beta, \gamma, \dots$

$$\therefore y_c = C_1 e^{\alpha x} + C_2 e^{\beta x} + C_3 e^{\gamma x} + \dots$$

Case-2

Roots are real and equal.

Let  $m = \alpha, \alpha, \beta$

$$y_c = (C_1 + C_2 x) e^{\alpha x} + C_3 e^{\beta x}$$

Let  $m = \alpha, \alpha, \alpha, \beta$ .

$$y_c = (C_1 + C_2 x + C_3 x^2) e^{\alpha x} + C_4 e^{\beta x}$$

Let  $m = \alpha, \alpha, \beta, \beta$

$$y_c = (C_1 + C_2 x) e^{\alpha x} + (C_3 + C_4 x) e^{\beta x}$$

Ex:-

$$\frac{d^2 y}{dx^2} - 4 \frac{dy}{dx} + 4y = 0.$$

$$m^2 - 4m + 4 = 0.$$

$$(m-2)^2 = 0.$$

$$m = 2, 2$$

$$y = (C_1 + C_2 x) e^{2x}$$

### Case-3

Roots are complex —

$$\text{Let } m = \alpha \pm i\beta$$

$$y_c = [C_1 \cos \beta x + C_2 \sin \beta x] e^{\alpha x}$$

$$\text{Let } m = \alpha \pm i\beta, \alpha \pm i\beta.$$

$$y_c = [(C_1 + C_2 x) \cos \beta x + (C_3 + C_4 x) \sin \beta x] e^{\alpha x}$$

Ex:-

$$\frac{d^2 y}{dx^2} - 2 \frac{dy}{dx} + 4y = 0$$

$$m^2 - 2m + 4 = 0$$

$$m = \frac{2 \pm \sqrt{4 - 16}}{2}$$

$$= \frac{2 \pm 2\sqrt{-3}}{2}$$

$$= 1 \pm i\sqrt{3}$$

$$= 1 \pm \sqrt{3} i$$

$$\therefore y = [C_1 \cos \sqrt{3} x + C_2 \sin \sqrt{3} x] e^{x}$$

# To find the roots of Eq<sup>n</sup>:-

For 3<sup>rd</sup> degree

i) Every 3<sup>rd</sup> degree eq<sup>n</sup> must contain at least 1 real root.

ii) If ~~some~~ <sub>sum</sub> of all the coeff = 0, then  $m = 1$  satisfies eq<sup>n</sup>  
↳ one root  
or  $(m-1)$  is a factor.

iii) If sum of coeff. of even powers = sum of coeff. of odd powers  
then  $m = -1$  satisfies the eq<sup>n</sup>. or  $(m+1)$  will be factor.  
↳ one root

$$\textcircled{1} (D^3 - 3D + 2)y = 0$$

$$m^3 - 3m + 2 = 0$$

Sum = 0  $\rightarrow$   $m=1$  is root  $\checkmark$

root	$m=1$	1	0	-3	2	$\rightarrow m^3$
		0	1	1	-2	
o always	$m=1$	1	1	-2	0	$\rightarrow m^2$
		0	1	2		
		1	2	0		$\rightarrow m$

$$m+2=0$$

$$m=-2$$

$\therefore m=1, 1 \text{ \& } -2$  are roots.

$$\textcircled{2} [D^3 - 4D^2 + 5D - 2]y = 0.$$

$$m^3 - 4m^2 + 5m - 2 = 0$$

$$1 - 4 + 5 - 2 = 0.$$

$m=1$	1	-4	5	-2
	0	1	-3	2
$m=1$	1	-3	2	
	0	1	-2	
	1	-2	0	

$$m-2=0$$

$$m=2$$

$\therefore m=1, 1, 2$   $\triangle$



$$\textcircled{4} (D^4 - D^3 - 9D^2 - 11D - 4)y = 0$$

$$m^4 - m^3 - 9m^2 - 11m - 4 = 0$$

$$\begin{array}{cccccc} 1 & -1 & -9 & -11 & -4 & \\ & \swarrow & \swarrow & \swarrow & \swarrow & \\ & & -12 & & & \\ & & & & & \end{array} \Rightarrow m = -1$$

$$\begin{array}{l} m = -1 \\ \hline 1 \quad -1 \quad -9 \quad -11 \quad -4 \\ 0 \quad -1 \quad 2 \quad 7 \quad 4 \\ \hline 1 \quad -2 \quad -7 \quad -4 \\ \quad \swarrow \quad \swarrow \quad \swarrow \\ \quad \quad -6 \quad -6 \quad -6 \\ \hline 0 \quad -1 \quad 3 \quad 4 \\ \hline m = -1 \\ \hline 1 \quad -3 \quad -4 \\ \quad \swarrow \quad \swarrow \\ \quad \quad -3 \quad -3 \\ \hline 0 \quad -1 \quad 4 \\ \hline 1 \quad -4 \end{array}$$

$$\begin{aligned} m - 4 &= 0 \\ m &= 4 \end{aligned}$$

$$\therefore m = -1, -1, -1, 4$$

$$y = (C_1 + C_2 x + C_3 x^2) e^{-x} + C_4 e^{4x}$$

$$\textcircled{5} (D^2 - 3D + 4)y = 0$$

$$m^2 - 3m + 4 = 0$$

$$\begin{aligned} (m-4)(m-?) \quad m &= \frac{3 \pm \sqrt{9-16}}{2} \\ &= \frac{3 \pm \sqrt{7}i}{2} \end{aligned}$$

$$y = \left( C_1 \cos \frac{\sqrt{7}x}{2} + C_2 \sin \frac{\sqrt{7}x}{2} \right) e^{\frac{3x}{2}}$$

$$\textcircled{6} [D^2 - (a+b)D + ab]y$$

$$m^2 - (a+b)m + ab = 0$$

$$(m-a)(m-b) = 0$$


$$m = a, b$$


$$\therefore y = C_1 e^{ax} + C_2 e^{bx}$$

# To find the particular Integral:-

$$F(D)y = Q$$

$$\text{then } y_p = PI = \frac{1}{F(D)} Q$$





Case-III. To find P.I of the form  $\frac{1}{f(D)} \sin ax$ ,  $\frac{1}{f(D)} \cos ax$ , replace  $D^2$  by  $-a^2$  if  $\phi(-a^2) \neq 0$ .

$$\begin{aligned} \frac{1}{D^2-2D+5} \sin 2x &= \frac{1}{-4-2D+5} \sin 2x = \frac{1}{1-2D} \times \frac{1+2D}{1+2D} \sin 2x \\ &= \frac{1+2D}{1-4D^2} \sin 2x = \frac{(1+2D) \sin 2x}{1-4(-4)} \\ &= \frac{\sin 2x + 4 \cos 2x}{17} \end{aligned}$$

Case-IV. To find P.I of the form  $\frac{1}{D^2+a^2} \sin ax$ ,  $\frac{1}{D^2+a^2} \cos ax$ ,

$$\frac{1}{D^2+a^2} \sin ax = \frac{x}{2D} \sin ax = \frac{x}{2} \left( \frac{-\cos ax}{a} \right)$$

$\frac{1}{D^2+a^2} \sin ax = \frac{-x}{2a} \cos ax$ $\frac{1}{D^2+a^2} \cos ax = \frac{x}{2a} \sin ax$
--

$$\frac{1}{D^2+4} \sin 2x = \frac{-x}{4} \cos 2x$$

$$\frac{1}{D^2+9} \cos 3x = \frac{x}{6} \sin 3x$$

$$\frac{1}{D^2+16} \sin 4x = \frac{-x}{8} \cos 4x$$

$$** \frac{1}{D^2+1} \cos 3x = \frac{1}{-9+6} \cos 3x = \frac{1}{3} \cos 3x$$

$$\textcircled{1} (D^2 - D - 2)y = \cos 2x$$

$$\textcircled{2} (D^2 - 4)y = \sin^2 x$$

$$\textcircled{3} (D^2 + 1)y = \sin 2x \cdot \cos x$$

$$\rightarrow \textcircled{1} \frac{1}{D^2 - D - 2} \cos 2x = \frac{1}{-4 - D - 2} \cos 2x = \frac{1}{-6 + D} \times \frac{-6 + D}{-6 + D} \cos 2x$$

$$= \frac{-6 + D}{36 - D^2} \cos 2x$$

$$= \left[ \frac{(-6 + D) \cos 2x}{36 - (-4)} \right]$$

$$= \frac{-6 \cos 2x - 2 \sin 2x}{40}$$

$$= -\frac{2}{40} (3 \cos 2x + \sin 2x)$$

$$= -\frac{1}{20} (3 \cos 2x + \sin 2x)$$

$$\textcircled{2} \frac{1}{D^2 - 4} \sin^2 x = \frac{1}{D^2 - 4} \left[ \frac{1 - \cos 2x}{2} \right] = \frac{1}{2} \left[ \frac{1}{D^2 - 4} (1) - \frac{1}{D^2 - 4} (\cos 2x) \right]$$

$$\begin{aligned} &= \frac{1}{D^2 - 4} e^{0x} \\ &= \frac{1}{D^2 - D} (5) = \frac{x}{2D - 1} (5) \end{aligned}$$

$$= \frac{1}{2} \left[ \frac{1}{0 - 4} \cdot 1 - \frac{1}{-4 - 4} \cos 2x \right]$$

$$= -\frac{1}{8} + \frac{1}{16} \cos 2x$$

$$\textcircled{3} (D^2+1)y = \sin 2x \cdot \cos x$$

$$= \frac{1}{2(D^2+1)} 2 \sin 2x \cdot \cos x$$

$$= \frac{1}{2} \left[ \frac{1 \sin 3x}{(D^2+1)} + \frac{1 \sin x}{D^2+1} \right]$$

$$\approx \frac{1}{2} \left[ \frac{1}{-9+1} \sin 3x - \frac{x}{2(1)} \cos x \right]$$

$$\approx -\frac{1}{16} \sin 3x - \frac{x}{4} \cos x$$

SHORTCUT

$$\rightarrow \text{if } (D^2+a^2)y = \tan ax$$

$$y_p = \frac{1}{a^2} \cos ax \cdot \log \left( \tan \left( \frac{\pi}{4} + \frac{ax}{2} \right) \right)$$

$$\rightarrow \text{if } (D^2+a^2)y = \sec ax$$

$$y_p = \frac{x}{a} \sin ax + \frac{1}{a^2} \cos ax \log(\cos ax)$$

ex:  $(D^2+4)y = \tan 2x$

$$y_p = \frac{1}{4} \cos 2x \cdot \log \left( \tan \left( \frac{\pi}{4} + x \right) \right)$$

ex:  $(D^2+9)y = \sec 3x$

$$y_p = \frac{x}{3} \sin 3x + \frac{1}{9} \cos 3x \cdot \log(\cos 3x)$$

Case-V.

To find the P.I of the form  $\frac{1}{f(D)} (x^m)$ .

①  $(D^2-4)y = x^2$

$$\begin{aligned} & \frac{1}{D^2-4} x^2 \\ &= \frac{1}{-4 \left[ 1 - \frac{D^2}{4} \right]} x^2 \\ &= -\frac{1}{4} \left[ 1 - \frac{D^2}{4} \right]^{-1} x^2 \\ &= -\frac{1}{4} \left[ 1 + \frac{D^2}{4} + \left( \frac{D^2}{4} \right)^2 + \dots \right] x^2 \\ &= -\frac{1}{4} \left[ x^2 + \frac{x^0}{4 \cdot 2} \right] \\ &= -\frac{1}{4} \left[ x^2 + \frac{1}{2} \right] \end{aligned}$$

②  $[D^3+8]y = x^4+2x+1$

$$\begin{aligned} &= \frac{1}{D^3+8} (x^4+2x+1) \\ &= \frac{1}{8 \left[ 1 + \frac{D^3}{8} \right]} (x^4+2x+1) \\ &= \frac{1}{8} \left[ 1 + \frac{D^3}{8} \right]^{-1} (x^4+2x+1) \\ &= \frac{1}{8} \left[ 1 - \frac{D^3}{8} + \frac{D^6}{8} + \dots \right] (x^4+2x+1) \\ &= \frac{1}{8} \left[ (x^4+2x+1) - \frac{1}{8} (24x+0+0) \right] \\ &= \frac{1}{8} (x^4 - x + 1) \end{aligned}$$

$$(1+D)^{-1} = 1 - D + D^2 - D^3 + \dots$$

$$(1-D)^{-1} = 1 + D + D^2 + D^3 + \dots$$

$$(1+D)^{-2} = 1 - 2D + 3D^2 - 4D^3 + \dots$$

$$(1-D)^{-2} = 1 + 2D + 3D^2 + 4D^3 + \dots$$

Remember

$$D^3(x^2) = 0$$

$$D^n(x^n) = n!$$

$$D^{n+1}(x^n) = 0$$

$$\textcircled{3} \quad (D^2 + 2D + 1)y = x^2 + x$$

$$\Rightarrow \frac{1}{(D+1)^2} (x^2 + x)$$

$$= [1 + D]^{-2} (x^2 + x)$$

$$= [1 - 2D + 3D^2] (x^2 + x)$$

$$= (x^2 + x) - 2(2x + 1) + 3(2 + 0)$$

$$= x^2 + x - 4x - 2 + 6$$

$$= x^2 - 3x + 4$$

Case-VI To find PI of the form  $\frac{1}{f(D)} e^{ax} \cdot v$ , where  $v$  is a fn. of  $x$ .

$$\boxed{\frac{1}{f(D)} e^{ax} \cdot v = e^{ax} \frac{1}{f(D+a)} \cdot v}$$

$$\textcircled{1} \quad (D^2 - 2D + 1)y = x^2 e^x$$

$$\textcircled{2} \quad (D^2 - 2D + 5)y = e^x \sin x$$

$$\textcircled{3} \quad (D^2 - 4D + 4)y = e^{2x} \cos 3x$$

$$\textcircled{4} \quad (D^3 + 3D^2 + 3D + 1)y = x^2 e^{-x}$$

$$\rightarrow \textcircled{1} \quad \frac{1}{(D-1)^2} e^x x^2 = e^x \cdot \frac{1}{(D+1-1)^2} x^2 = e^x \left[ \frac{1}{D^2} x^2 \right]$$

$$= e^x \frac{x^2}{2}$$

represents integration

$$\textcircled{2} \quad \frac{1}{D^2 - 2D + 5} (e^x \sin x)$$

$$= e^x \left[ \frac{1}{(D+1)^2 - 2(D+1) + 5} \sin x \right]$$

$$= e^x \left[ \frac{1}{D^2 + 1 + \cancel{D} - 2D - 2 + 5} \sin x \right]$$

$$= e^x \left[ \frac{1}{\cancel{D} + 4} \sin x \right]$$

$$= e^x \left[ \frac{1}{-1+4} \sin x \right]$$

$$= \frac{e^x \sin x}{3}$$

$$\textcircled{3} \quad \frac{1}{(D^2 - 4D + 4)} e^{2x} \cos 3x$$

$$= e^{2x} \left[ \frac{1}{(\cancel{D})^2} \cos 3x \right]$$

$$= e^{2x} \left[ \frac{1}{D^2} \cos 3x \right]$$

$$= e^{2x} \frac{1}{-9} \cos 3x$$

$$= -\frac{e^{2x}}{9} \cos 3x$$

AND PHOTOSTY & BOOK CENTRE  
 ADD. NEAR MADE EASY LANE NO. 4  
 WESTEND MARKET, SAIDUL ALAM  
 NEW DELHI - 110030  
 MOB: 9809127702

$$\textcircled{4} \frac{1}{(D^3 + 3D^2 + 3D + 1)} e^{-x} x^2$$

$$= \frac{1}{(D+1)^3} e^{-x} x^2$$

$$= e^{-x} \frac{1}{(D+1)^3} x^2$$

$$= e^{-x} \left[ \frac{1}{D^3} x^2 \right]$$

$$\int \int \int x^2 = \frac{x^3}{3} = \frac{x^4}{12} = \frac{x^5}{60}$$

$$= e^{-x} \left( \frac{x^5}{60} \right) \underline{\underline{Ans}}$$

Case-VII. To find the P.I of the form  $\frac{1}{f(D)} x \cdot v$ , where  $v$  is a fn of  $x$

(Trigonometric fn)

$$\frac{1}{f(D)} x \cdot v = x \cdot \frac{1}{f(D)} \cdot v - \frac{f'(D)}{[f(D)]^2} \cdot v$$

$$\textcircled{1} (D^2 + 4)y = x \cos x$$

$$\textcircled{2} (D^2 - 2D + 1)y = x e^x \sin x$$

$$\textcircled{1} \frac{1}{D^2 + 4} (x \cos x)$$

$$= x \cdot \frac{1}{(D^2 + 4)} \cos x - \frac{2D}{(D^2 + 4)^2} \cdot \cos x$$

$$= x \cdot \frac{1}{-1 + 4} \cos x - \frac{2(-\sin x)}{(-1 + 4)^2}$$

$$= \frac{x}{3} \cos x + \frac{2}{9} \sin x$$

$$\textcircled{2} \quad \frac{1}{(D-1)^2} e^x \cdot x \cdot \sin x$$

$$= e^x \left[ \frac{1}{(D+1-1)^2} x \sin x \right]$$

$$= e^x \left[ \frac{1}{D^2} x \sin x \right]$$

$$= e^x \left[ x \frac{1}{-a^2} \sin x - \frac{2D}{(-a^2)^2} \sin x \right]$$

$$= e^x \left[ x \frac{1}{-1} \sin x - \frac{2 \cos x}{(-1)^2} \right]$$

$$= e^x (-x \sin x - 2 \cos x) \text{.}$$

# Differential Eq<sup>n</sup> with variable co-efficients :-

or

Cauchy Euler form :-

---

→ A diff eq<sup>n</sup> is of the form —

$$a_0 x^n \frac{d^2 y}{dx^2} + a_1 x^{n-1} \frac{dy}{dx} + a_2 x^{n-2} y + \dots + a_n y = Q$$

where  $a_0, a_1, a_2, \dots, a_n$  are constants

and  $Q$  be the function of  $x$ .

is said to be Diff<sup>n</sup> Eq<sup>n</sup> with Variable Coefficients.

$$x = e^z \Rightarrow z = \log x.$$

$$\frac{dz}{dx} = \frac{1}{x} \quad \& \quad \frac{dx}{dz} = x$$

$$\frac{d}{dz} = \frac{d}{dx} \cdot \frac{dx}{dz} \xrightarrow{x} = x \frac{d}{dx} = xD = \frac{d}{dz} = \theta$$

Now—

$$\frac{d}{dz} \left( x \frac{d}{dx} \right) = \frac{d^2}{dz^2}$$

$$\frac{d}{dx} \left( x \frac{d}{dx} \right) \cdot \frac{dx}{dz} = \theta^2$$

$$x \left( x \frac{d^2}{dx^2} + \frac{d}{dx} \cdot 1 \right) = \theta^2$$

$$\therefore x^2 \frac{d^2}{dx^2} + x \frac{dx}{dx} = \theta^2$$

$$x^2 \frac{d^2}{dx^2} + \theta = \theta^2$$

$$x^2 \frac{d^2}{dx^2} = \theta(\theta - 1)$$

$$\therefore x^2 \frac{d^2}{dx^2} = x^2 D^2 = \theta(\theta - 1)$$

\*\*\*  
 Imp  
 Substitution  
 to Remember

Let  $x \frac{d}{dx} = xD = \theta$ .

$$x^2 \frac{d^2}{dx^2} = x^2 D^2 = \theta(\theta-1) = \theta^2 - \theta$$

$$x^3 \frac{d^3}{dx^3} = x^3 D^3 = \theta(\theta-1)(\theta-2) = \theta^3 - 3\theta^2 + 2\theta$$

Variable Coeff.

①  $(x^2 D^2 - xD - 3)y = 0$

$$= [\theta^2 - \theta - \theta - 3]y = 0$$

$$= m^2 - 2m - 3 = 0$$

$$= (m-3)(m+1) = 0$$

$$m = 3, -1$$

$$y = C_1 e^{3x} + C_2 e^{-x}$$

$$= C_1 (e^x)^3 + C_2 (e^x)^{-1}$$

$$y = C_1 x^3 + C_2 x^{-1}$$

observe the diff<sup>n</sup> bet<sup>n</sup> const coeff  
 approach

$$D^2 - D + 2 = 0$$

$$m^2 - m + 2 = 0$$

②  $[x^3 D^3 + 3x^2 D^2 - 2xD + 2]y = 0$

$$[\theta^3 - 3\theta^2 + 2\theta + 3\theta^2 - 3\theta - 2\theta + 2]y = 0$$

$$[\theta^3 - 3\theta + 2]y = 0$$

$$m^3 - 3m + 2 = 0$$

~~$$m^3 - 3m + 2 = 0$$~~

$\therefore m = 1$  satisfies.

	1	0	-3	2
$m=1$	0	1	1	-2
$m=1$	1	1	-2	0
	0	1	2	0
	1	2	0	

$$m+2=0 \Rightarrow m = -2$$

$$\therefore y = (c_1 + c_2 z) e^z + c_3 e^{-2z}$$

$$\checkmark \boxed{y = (c_1 + c_2 \log x) x + c_3 x^{-2}}$$

$$\textcircled{3} \quad [x^2 D^2 + xD - 4] y = 0, \quad x(0) = 0 \\ y(1) = 1$$

$$[D^2 - 0 + 0 - 4] y = 0$$

$$m^2 - 4 = 0$$

$$m = \pm\sqrt{4} = \pm 2 = 2, -2$$

$$\therefore y = c_1 e^{2z} + c_2 e^{-2z}$$

$$\Rightarrow y = c_1 x^2 + c_2 x^{-2}$$

$$\Rightarrow y = c_1 x^2 + \frac{c_2}{x^2}$$

$$\Rightarrow x^2 y = c_1 x^4 + c_2$$

$$\therefore x(0) = 0$$

$$\Rightarrow \cancel{0 = c_1 \cdot 0^4 + c_2}$$

$$0 \cdot 0 = c_1(0) + c_2$$

$$\Rightarrow \boxed{c_2 = 0}$$

$$y(1) = 1$$

$$\Rightarrow 1^2 \cdot 1 = c_1(1)^4 + c_2$$

$$1 = c_1 + 0$$

$$\boxed{c_1 = 1}$$

$$\therefore y = 1(x^2) + \frac{0}{x^2}$$

$$\checkmark \boxed{y = x^2}$$

$$Q. [x^4 D^3 + 2x^3 D^2 - x^2 D + x] y = 1$$

$$\Rightarrow (x^3 D^3 + 2x^2 D^2 - x D + 1) y = \frac{1}{x}$$

$$\Rightarrow (\theta^3 - 3\theta^2 + 2\theta - \theta + 1) y = e^{-z}$$

$$\Rightarrow (\theta^3 - \theta^2 - \theta + 1) y = e^{-z}$$

$$m^3 - m^2 - m + 1 = 0$$

$$m^2(m-1) - 1(m-1) = 0$$

$$(m-1)(m^2-1) = 0$$

$$(m-1)(m-1)(m+1) = 0$$

$$m = 1, 1, -1$$

$$\therefore y_c = (c_1 + c_2 z) e^z + c_3 e^{-z}$$

$$\Rightarrow y_c = (c_1 + c_2 \log x) x + c_3 x^{-1}$$

$$\Rightarrow y_p = \frac{1}{\theta^3 - \theta^2 - \theta + 1} e^{-z}$$

$$= \frac{z}{3\theta^2 - 2\theta - 1} e^{-z}$$

$$= \frac{z}{3(-1)^2 - 2(-1) - 1} e^{-z} = \frac{z e^{-z}}{4}$$

$$= \frac{\log x}{4} \times \frac{1}{x} = \frac{1}{4x} \log x$$

$$\textcircled{5} \quad (x^2 D^2 - x D + 1)y = 0.$$

$$\theta^2 - \theta - \theta + 1 = 0$$

$$m^2 - 2m + 1 = 0$$

$$m = +1, +1$$

$$= (C_1 + C_2 \log x) e^{z \rightarrow x}$$

$$= (C_1 + C_2 \log x) x.$$

$$\textcircled{6} \quad x^2 \frac{d^3 y}{dx^3} + 2x \frac{d^2 y}{dx^2} - 2 \frac{dy}{dx} = 0$$

$$x^3 \frac{d^3 y}{dx^3} + 2x^2 \frac{d^2 y}{dx^2} - 2x \frac{dy}{dx} = 0.$$

$$\Rightarrow (x^3 D^3 + 2x^2 D^2 - 2x D)y = 0$$

$$\Rightarrow [\theta^3 - 3\theta^2 + 2\theta + 2\theta^2 - 2\theta - 2\theta] y = 0$$

$$\Rightarrow m^3 - 3m^2 - 2m = 0$$

$$\Rightarrow m(m^2 - m - 2) = 0$$

$$\Rightarrow m(m-2)(m+1) = 0$$

$$m = 0, 2, -1$$

$$\therefore y = C_1 e^{0z} + C_2 e^{2z} + C_3 e^{-z}$$

$$\checkmark \boxed{y = C_1 + C_2 x^2 + C_3 x^{-1}}$$

$$\textcircled{7} \quad 4x^2y'' + 12xy' + 3y = 0$$

$$(4x^2D^2 + 12xD + 3)y = 0$$

$$\Rightarrow (4\theta^2 - 4\theta + 12\theta + 3)y = 0$$

$$\Rightarrow 4m^2 + 8m + 3 = 0.$$

$$\cancel{m = \frac{-8 \pm \sqrt{64 - 48}}{2} = \frac{-8 \pm \sqrt{16}}{2} = \frac{-8 \pm 4}{2} = \frac{-8+4}{2} = -2, \frac{-8-4}{2} = -6}$$

$$m = -\frac{1}{2}, -\frac{3}{2}$$

$$\therefore y = c_1 x^{-\frac{1}{2}} + c_2 x^{-\frac{3}{2}}$$

$$\textcircled{8} \quad x^2y'' + xy' - y = 0$$

$$[\theta^2 - \theta - 1]y = 0$$

$$m^2 - 1 = 0$$

$$m = \pm 1$$

$$y = c_1 e^x + c_2 e^{-x}$$

$$\boxed{y = c_1 x + c_2 x^{-1}}$$

## # Orthogonal Trajectories:-

→ The family of curves which cuts every member of the given family orthogonally is called the orthogonal Trajectory of the given family.

$$\longrightarrow f(x, y, c) = 0 \quad \text{--- ①}$$

replace

$$\longrightarrow F\left(x, y, \frac{dy}{dx}\right) = 0 \quad \text{--- ②}$$

replace  $\left(\frac{dy}{dx}\right)$  by  $\left(\frac{-dx}{dy}\right)$

$$\longrightarrow g\left(x, y, -\frac{dx}{dy}\right) = 0 \quad \text{--- ③}$$

The sol<sup>n</sup> of eq<sup>n</sup> ③ is called the orthogonal trajectory of the given family.

Note:-

If eq<sup>n</sup> ② and eq<sup>n</sup> ③ are identical, then the curve is said to be self orthogonal.

Q. Find the Orthogonal Trajectory of family of lines passing through the origin—

i) family of lines

ii) family of Circles having the centre at the origin.

iii) family of Parabolas

iv) family of ellipses.

~~\*\*\*~~  
If fn given, then  
Steps —

find  $\frac{dy}{dx}$

replace  $\frac{dy}{dx}$  by  $\frac{-dx}{dy}$

and integrate & solve.

That will represent  
orthogonal of that fn

$$\textcircled{1} \rightarrow y = mx \quad \text{---} \textcircled{1}$$

$$\frac{dy}{dx} = m$$

$$y = x \frac{dy}{dx} \quad \text{---} \textcircled{2}$$


$$y = -x \frac{dx}{dy} \quad \text{---} \textcircled{3}$$

$$\int y dy + \int x dx = \int 0$$

$$\frac{y^2}{2} + \frac{x^2}{2} = c$$

$$x^2 + y^2 = 2c$$

$$\boxed{x^2 + y^2 = r^2} \rightarrow \text{Circle}$$



$$\textcircled{2} \quad y^2 = 4ax \quad \text{---} \textcircled{1}$$

$$2y \frac{dy}{dx} = 4a$$

$$y^2 = 2y \frac{dy}{dx} x$$

$$\cancel{2 \frac{dy}{y}} = \cancel{\int \frac{dx}{x}}$$

$$2x \frac{dy}{dx} = y \quad \text{---} \textcircled{2}$$

$$y = -2x \frac{dx}{dy} \quad \text{---} \textcircled{3}$$

$$\int y dy = \int -2x dx$$

$$\frac{y^2}{2} = -\frac{2x^2}{2} + c$$

$$\Rightarrow \boxed{\frac{x^2 + y^2}{2} = c} \rightarrow \text{Ellipse} \dots$$

$$\rightarrow x^2 - y^2 = a^2 \quad \text{--- ①}$$

~~000~~

$$x - y \frac{dy}{dx} = 0$$

$$x + y \frac{dx}{dy} = 0$$

$$x dy + y dx = 0$$

$$\int d(xy) = \int 0$$

$$\boxed{xy = c}$$

Conclusion - Orthogonal Trajectories of family of ---

circle  $\xrightarrow{is}$  st line.

st line  $\xrightarrow{is}$  circle

Parabola  $\xrightarrow{is}$  ellipse

Hyperbola  $\xrightarrow{is}$  Rectangular Hyperbola.

$$\textcircled{2} \quad y^2 = 4a(x+a) \quad \text{--- ①}$$

$$2yy_1 = 4a \cdot 1 \Rightarrow a = \frac{yy_1}{2}$$

$$y^2 = 2yy_1 \left( x + \frac{yy_1}{2} \right)$$

$$y = 2xy_1 + yy_1^2 \quad \text{--- ②}$$

replace  $y_1$  by  $-\frac{1}{y_1}$

$$y = -\frac{2x}{y_1} + \frac{y}{y_1^2}$$

$$yy_1^2 = -2xy_1 + y$$

$$yy_1^2 + 2xy_1 = y \quad \text{--- (3)}$$

$$\therefore \text{eq}^n \text{ (2)} = \text{eq}^n \text{ (3)}$$

$\therefore$  Curve is said to be self orthogonal. ✓

#

$$\frac{x^2}{a^2 + \lambda^2} + \frac{y^2}{b^2 + \lambda^2} = 1$$

Where  $\lambda$  is a parameter

Imp Result

is also self-Orthogonal. ✓

#

$$x^2 + y^2 + 2gx + c = 0$$

Imp Concept

$$x^2 + y^2 + 2fy + c = 0$$

$$\text{O.T of } x^2 + y^2 + 2gx - 5 = 0 \text{ is ---}$$

$$x^2 + y^2 - 2fy + 5 = 0$$

→ always take opposite coefficient sign.

#

$$f(r, \theta, c) = 0 \quad \text{--- (1)}$$

$$F(r, \theta, \frac{dr}{d\theta}) = 0 \quad \text{--- (2)}$$

replace  $\frac{dr}{d\theta}$  by  $-r^2 \frac{d\theta}{dr}$

$$g(r, \theta, -r^2 \frac{d\theta}{dr}) = 0 \quad \text{--- (3)}$$

The sol<sup>n</sup> of eq<sup>n</sup> (3) is called the Orthogonal of the given family.

$$\textcircled{1} \rightarrow r = a\theta$$

$$\log r = \log a + \log \theta$$

$$\frac{1}{r} \frac{dr}{d\theta} = 0 + \frac{1}{\theta}$$

$$\frac{1}{r} \times -r^2 \frac{d\theta}{dr} = \frac{1}{\theta}$$

$$\int \frac{dr}{r} = - \int \frac{d\theta}{\theta}$$

$$\log r = -\frac{\theta^2}{2} + C$$

$$r = e^{-\frac{\theta^2}{2} + C}$$

$$r = e^{-\frac{\theta^2}{2}} \cdot e^C$$

$$r = k e^{-\frac{\theta^2}{2}}$$

$\textcircled{2}$

$$r = \frac{2a}{1 - \cos \theta}$$

$$\log r = \log 2a - \log (1 - \cos \theta)$$

$$\frac{1}{r} \frac{dr}{d\theta} = 0 - \frac{1}{1 - \cos \theta} \cdot \sin \theta$$

$$\frac{1}{r} \times -r^2 \frac{d\theta}{dr} = -\frac{\sin \theta}{1 - \cos \theta}$$

$$r \frac{d\theta}{dr} = \frac{\sin \theta}{1 - \cos \theta}$$

$$\Rightarrow \frac{dr}{r} = \frac{1 - \cos \theta}{\sin \theta} d\theta$$

$\frac{1 - \cos \theta}{\sin \theta} = \frac{2 \sin^2 \frac{\theta}{2}}{2 \sin \frac{\theta}{2} \cos \frac{\theta}{2}} = \frac{\sin \frac{\theta}{2}}{\cos \frac{\theta}{2}}$

$$\Rightarrow \int \frac{dr}{r} = \int \frac{\sin \frac{\theta}{2}}{\cos \frac{\theta}{2}} d\theta$$

$$\Rightarrow \log r = -2 \log \cos \frac{\theta}{2} + \log C$$

$$\Rightarrow \log r = \log \frac{c}{\cos^2 \frac{\theta}{2}}$$

$$\Rightarrow r = \frac{c}{\cos^2 \frac{\theta}{2}}$$

$$\Rightarrow r = \frac{c}{\frac{1 + \cos \theta}{2}}$$

$$\Rightarrow \boxed{r = \frac{2c}{1 + \cos \theta}}$$

i.e.,  $\boxed{r = \frac{2a}{1 - \cos \theta}}$   $\xleftrightarrow{\text{Orthogonal}}$   $\boxed{r = \frac{2c}{1 + \cos \theta}}$

③  $r^n = a^n \sin^n \theta$

$$n \log r = n \log a + \log \sin^n \theta$$

$$\frac{n}{r} \frac{dr}{d\theta} = \frac{n \cos \theta}{\sin \theta}$$

$$\frac{n}{r} - r^2 \frac{d\theta}{dr} = \frac{n \cos \theta}{\sin^n \theta}$$

$$- n r \frac{d\theta}{dr} = \frac{n \cos \theta}{\sin^n \theta}$$

$$\theta \frac{dr}{r} = \frac{1}{n} \int -n \frac{\sin \theta}{\cos \theta} d\theta$$

$$\log r = \frac{1}{n} \log \cos \theta + \log c$$

$$n \log r = \log \cos \theta + n \log c$$

$$\log r^n = \log \cos \theta + \log c^n$$

$$\boxed{r^n = c^n \cos \theta}$$

i.e.,  $\boxed{r^n = a^n \sin^n \theta}$   $\xleftrightarrow{\text{Orthogonal}}$   $\boxed{r^n = c^n \cos \theta}$

Q. The diff. eq<sup>n</sup> of family of curves is given by  $\frac{dy}{dx} = \frac{x}{y}$   
 then the system of orthogonal Trajectories of the given family of curves —

$$\rightarrow \frac{dy}{dx} = \frac{x}{y}$$

$$-\frac{dx}{dy} = \frac{x}{y}$$

$$-\frac{dx}{x} = \frac{dy}{y}$$

$$-\log x = \log y + \log C$$

$$\Rightarrow \boxed{xy = C}$$

This is the orthogonal Trajectory of family  $\frac{dy}{dx} = \frac{x}{y}$

### # Differential Eq<sup>n</sup> of First Order but not first Degree :-

— A differential eq<sup>n</sup> of the form —

$$a_0 \left(\frac{dy}{dx}\right)^n + a_1 \left(\frac{dy}{dx}\right)^{n-1} + a_2 \left(\frac{dy}{dx}\right)^{n-2} + \dots + a_n = 0.$$

Where  $a_0, a_1, a_2, \dots, a_n$  are fn. of  $x$  and  $y$  or constants  
 is said to be differential eq<sup>n</sup> of 1<sup>st</sup> order but not first degree.

$$\text{Let } \frac{dy}{dx} = P$$

$$\boxed{a_0 P^n + a_1 P^{n-1} + a_2 P^{n-2} + \dots + a_n = 0}$$

$$f(x, y, P) = 0$$

— This eq<sup>n</sup> containing 3 variables  $x, y$  and  $P$ , therefore it may contain 3 types of solution:

→ It may be solvable for  $P$ , may be solvable for  $y$ , may be solvable for  $x$ .

\* The eq<sup>n</sup> solvable for P -

→ If it is solvable for P, then the eq<sup>n</sup> must be factorized.

$$\rightarrow [P - f_1(x, y)] [P - f_2(x, y)] [P - f_3(x, y)] \dots [P - f_n(x, y)] = 0$$

then  $P - f_1(x, y) = 0$

replace by  $\frac{dy}{dx}$

$$\boxed{\frac{dy}{dx} = f_1(x, y)}$$

ex:- ①  $P^2 - 5P + 6 = 0$  where  $P = \frac{dy}{dx}$ .

$$(P-3)(P-2) = 0.$$

$$P-2 = 0$$

$$\frac{dy}{dx} - 2 = 0.$$

$$\int dy = \int 2 dx = 0$$

$$y - 2x - C_1 = 0 \quad \text{--- ①}$$

$$y - 3x - C_2 = 0 \quad \text{--- ②}$$

$$\boxed{(y - 2x - C_1)(y - 3x - C_2) = 0}$$

②  $x^2 P^2 + xyP - 6y^2 = 0$

$$\Rightarrow x^2 P^2 + 3xyP - 2xyP - 6y^2 = 0$$

$$\Rightarrow xP(xP + 3y) - 2y(xP + 3y) = 0$$

$$\Rightarrow (xP + 3y)(xP - 2y) = 0$$

$$xP + 3y = 0$$

$$x \frac{dy}{dx} + 3y = 0$$

$$\frac{dy}{dx} = -\frac{3}{x} y = 0$$

$$y = e^{\int P dx} = \int \frac{0}{x} e^{\int P dx} + C$$

$$y(x^3) - C_1 = 0 \quad \text{--- ①}$$

$$x \frac{dy}{dx} - 2y = 0.$$

$$\frac{dy}{dx} - \frac{2}{x} y = 0.$$

$$y \left( \frac{1}{x^2} \right) - C_2 = 0 \quad \text{--- ②}$$

Soln  
obtained

$$\boxed{(x^3 y - C_1)(y - C_2 x^2) = 0}$$

$$\textcircled{2} \quad xyp^2 - (x^2 + y^2)p + xy = 0.$$

$$xy p^2 - x^2 p - y^2 p + xy = 0.$$

$$xp(y p - x) - y(y p - x) = 0$$

$$(xp - y)(yp - x) = 0.$$

$$xp - y = 0.$$

$$x \frac{dy}{dx} - y = 0$$

$$\frac{dy}{dx} - \frac{y}{x} = 0.$$

$$y \left( \frac{1}{x} \right) - C_2 = 0 \quad \text{--- ①}$$

$$yp - x = 0.$$

$$y \frac{dy}{dx} - x = 0$$

$$y dy - x dx = 0$$

$$\frac{y^2}{2} - \frac{x^2}{2} = C_2$$

$$\Rightarrow y^2 - x^2 - 2C_2 = 0 \quad \text{--- ②}$$

$$\boxed{(y - C_1 x)(y^2 - x^2 - 2C_2) = 0}$$

Trick  
When to apply linear approach  
When to apply variable separable?  
→  $x \frac{dy}{dx}$  or  $y \frac{dx}{dy}$  → linear  
→  $x \frac{dx}{dy}$  or  $y \frac{dy}{dx}$  → Variable separable

\* When factorization is not possible, then Solvable for P is not possible  
two case arises —

When Solvable for y ?	When Solvable for x ?
① The degree of y must be 1.	① The degree of x must be 1
② Separate y from x & P $y = f(x, P)$	② Separate x from y & P $x = f(y, P)$
③ diff. both side w.r.to 'x'	③ Diff. both side w.r.to 'y'.
④ Replace $\frac{dy}{dx}$ by P	④ Replace $\frac{dx}{dy}$ by $\frac{1}{P}$ .
⑤ neglect the terms or cancel the terms which does not contain $\frac{dP}{dx}$ on both sides.	⑥ Neglect the term or Cancel the term which does not contain $\frac{dP}{dy}$ on both sides.

ex:-  $y + px = P^2 x^4$  — ①

∴ degree of y = 1 → solvable for y ✓

∴ all coeff are 1, 1, 1 ⇒ means factorization not possible ⇒ not solvable for P ✗

$$y = P^2 x^4 - px$$

$$\frac{dy}{dx} = P^2(4x^3) + x^4 \left( 2P \frac{dP}{dx} \right) - P(1) - x \left( \frac{dP}{dx} \right)$$

$$P - 4P^2 x^3 + P = (2Px^4 - x) \frac{dP}{dx}$$

$$2P(1 - 2Px^3) = -x(1 - 2Px^3) \frac{dP}{dx}$$

$$2P = -x \frac{dP}{dx}$$

$$\Rightarrow \int \frac{dP}{P} = \int -2 \frac{dx}{x}$$

$$\Rightarrow \log P = \log \frac{C}{x^2}$$

$$\boxed{P = \frac{C}{x^2}}$$

2 methods now to get sol<sup>n</sup> of diff<sup>n</sup> eq<sup>n</sup> in Q.

either put  $P = \frac{dy}{dx}$

$$\frac{dy}{dx} = \frac{C}{x^2}$$

and integrate and solve.

$$dy - \frac{C dx}{x^2} = 0$$

$$\boxed{y + \frac{C}{x} = C}$$

sol<sup>n</sup>

Put in eq<sup>n</sup> of Question.

$$y + px = \frac{C}{x^2} \cdot x^2$$

$$\Rightarrow \boxed{y + \frac{C}{x} = C}$$

sol<sup>n</sup>

Ans

Q2

$$y = 2px + y^2 p^3$$

$$2px = y - y^2 p^3$$

$$x = \frac{y}{2p} - \frac{y^2 p^3}{2p}$$

Ans

$$2x = \frac{1}{p} \cdot y - y^2 p^2$$

$$2 \frac{dx}{dy} = \frac{1}{p} \cdot 1 + y \left( -\frac{1}{p^2} \frac{dp}{dy} \right) - 2y p^2 - y^2 \left( 2p \frac{dp}{dy} \right)$$

$$2 \cdot \frac{1}{p} = \frac{1}{p} - \frac{y}{p^2} \frac{dp}{dy} - 2y p^2 - 2p y^2 \frac{dp}{dy}$$

$$\frac{2}{p} - \frac{1}{p} + 2y p^2 = \left( -\frac{y}{p^2} - 2p y^2 \right) \frac{dp}{dy}$$

$$\frac{(1 + 2y p^3)}{p} = -y \left( \frac{1 + 2y p^3}{p^2} \right) \frac{dp}{dy}$$

~~y~~

$$1 = -\frac{y}{P} \frac{dP}{dy}$$

$$\int \frac{dP}{P} = \int -\frac{dy}{y}$$

$$\log P = \log \frac{C}{y}$$

$$\boxed{P = \frac{C}{y}} \quad \text{Put in (A)}$$

$$\therefore y = 2x \cdot \frac{C}{y} + y^2 \frac{C^3}{y^3}$$

$$\boxed{y^2 = 2Cx + C^3}$$

Shortcut

$$\boxed{y = Px + \phi(P)}$$

replace P by c

$$\boxed{y = cx + \phi(c)}$$

ex: - ①  $y = Px - P^2$

gen soln =  $\boxed{y = cx - c^2}$

②  $\log(Px - y) = P$

$\therefore Px - y = e^P$

$y = Px - e^P$

$\boxed{y = cx - e^c}$

## Numerical Methods

I: Sol<sup>n</sup> of non-linear equation:-

\*\*\*

non linear means ?

linear  $\rightarrow$  degree 1.

non-linear  $\rightarrow$  degree  $\neq 1$

Let  $f(x) = 0$  be any non linear equation to find the approximate root of  $f(x) = 0$

choose any two values of  $x = a, b$ , particularly adjacent values in such a way that their functional values must have opposite signs.

ex:-  $f(a) < 0, f(b) > 0$

Then, we can say that root lies bet<sup>n</sup> a and b.

This approximate root can be find by —

- ① method of bisection —
- ② Regular Falsi method
- ③ Newton - Raphson

\* Method of Bisection:-

$$x = a, b$$

$$f(a) < 0, f(b) > 0.$$

$$\frac{f(a) \quad 0 \quad f(b)}{\quad \quad \quad}$$

$$x_1 = \frac{a+b}{2}$$

$$f(x_1) < 0$$

$$f(a), f(x_1), f(b)$$

$< \quad < \quad >$

$$x_2 = \frac{x_1 + b}{2}$$

$$f(x_2) > 0$$

x.

$$x_3 = \frac{x_1 + x_2}{2}$$

upto 13<sup>th</sup> - 14<sup>th</sup> step

The rate of convergence in bisection method is very-very slow.

→ Comparatively, the Bisection method, Regula-falsi method has a little fast & rate of convergence.

\* Regula-falsi Method:-

$$x_1 = \frac{a f(b) - b f(a)}{f(b) - f(a)}$$

\* Newton-Raphson method:-

→ has a very fast rate of convergence.

→ and the convergence of Newton Raphson method is 2<sup>nd</sup> order convergence or Quadratic convergence.

initial guess (99% chance given for Q.)  
if chance not given then how to get  $x_0$

$$x_1 = x_0 - \frac{f(x_0)}{f'(x_0)}$$

$$x_2 = x_1 - \frac{f(x_1)}{f'(x_1)}$$

$$x_3 = x_2 - \frac{f(x_2)}{f'(x_2)}$$

$$x_{n+1} = x_n - \frac{f(x_n)}{f'(x_n)}$$

↳ take  $f(1) \rightarrow x_0$   
 $f(2) \rightarrow x_0$   
} whatever is nearer to origin that value will be taken.  
when  $f(1)$  and  $f(2)$  must have opposite signs.

$$\textcircled{1} \quad f(x) = x^3 + 2x - 5 = 0$$

$$x_0 = 1 \quad \text{find } x_1:$$

$$\rightarrow f'(x) = 3x^2 + 2$$

$$x_1 = 1 - \frac{x^3 + 2x - 5}{3x^2 + 2}$$

$$= 1 - \left[ \frac{1 + 2 - 5}{5} \right]$$

$$= 1 + \frac{2}{5}$$

$$x_1 = 1.4 \quad \underline{\underline{\text{Ans}}}$$

$$\textcircled{2} \quad f(x) = x^3 + 3x - 7 = 0$$

$$x_0 = 1.$$

$$x_1 = ?$$

$$\rightarrow x_1 = 1 - \frac{x^3 + 3x - 7}{3x^2 + 3}$$

$$= 1 - \frac{-8}{6}$$

$$= 1 + \frac{1}{2}$$

$$= 1.5 \quad \underline{\underline{\text{Ans}}}$$

$$\textcircled{3} \quad f(x) = x^3 - 2x - 5 = 0.$$

$$f(0) = -5 < 0$$

$$f(1) = 1 - 2 - 5 = -6 < 0$$

$$f(2) = 8 - 4 - 5 = -1 < 0 \quad \rightarrow x_0 = 2 \text{ is chosen}$$

$$f(3) = 27 - 6 - 5 = 16 > 0.$$

$$x_0 = 2$$

$$x_1 = 2 - \frac{(-1)}{3(2^2) - 2} = 2 + \frac{1}{10} = 2.1$$

$$\textcircled{4} \quad f(x) = e^x - 1 = 0 \quad @ 0.71828$$

$$x_0 = -1 \quad @ 0.34567$$

$$x_1 = ? \quad @ 0.12326$$

$$\quad \quad \quad @ 0.00000$$

$$\rightarrow f'(x) = e^x$$

$$x_1 = -1 - \frac{e^{-1} - 1}{e^{-1}}$$

$$= -1 - 1 + e$$

$$= e - 2$$

$$= 2.71 - 2$$

$$= \underline{\underline{0.71}}$$

$$\textcircled{5} \quad f(x) = x^2 - 2x - 1$$

$$x_0 = 2$$

$$x_2 = \underline{\quad}$$

$$\textcircled{a} 2.425 \quad \textcircled{b} 2.423 \quad \textcircled{c} 2.419 \quad \textcircled{d} 2.417$$

$$\rightarrow f'(x) = 2x - 2$$

$$x_1 = 2 - \frac{-1}{2} = 2 + 0.5 = 2.5$$

$$x_2 = 2.5 - \frac{6.25 - 5 - 1}{5 - 2}$$

$$= 2.5 - \frac{0.25}{3}$$

$$= 2.5 - 0.083$$

$$= \underline{\underline{2.417}}$$

$$= \underline{\underline{2.417}}$$

ADD NEAR MADE SASI LINE NO. 7  
WESTERD WAFER, SINDHUR  
NEW PATEL, 110000  
MUM. 400012/1001

Q.  $f(x) = x^4 - 3x + 1 = 0$

$x_0 = 0$

$x_1 = \underline{\hspace{2cm}}$

- (A)  $-\frac{1}{3}$  (B)  $\frac{1}{3}$  (C) 3 (D) -3

$\rightarrow f'(x) = 4x^3 - 3$

$x_1 = 0 - \frac{1}{-3}$

$= \frac{1}{3}$  Ans

Q: The approximate root of eq<sup>n</sup>  $f(x) = (x-1)^2 + x - 3 = 0$  is to be solved by the Newton Raphson method. The method will fail in the very first iteration if the initial guess  $x_0 = \underline{\hspace{2cm}}$

- (A) 0 (B) 0.5 (C) 1.0 (D) 1.5

$\rightarrow f(x) = x^2 - 2x + 1 + x - 3$   
 $= x^2 - x - 2$

$f'(x) = 2x - 1$

$x_1 = x_0 - \frac{f(x_0)}{f'(x_0)}$

$= x_0 - \frac{x_0^2 - x_0 - 2}{2x_0 - 1} \rightarrow 0$  method will fail.

i.e. method will fail if  $2x_0 - 1 = 0$

$\Rightarrow x_0 = \underline{\underline{0.5}}$

Q The Newton Raphson iteration  $x_{n+1} = \frac{1}{2} \left[ x_n + \frac{R}{x_n} \right]$  can be used to

Imp  
Concept

compute the

- a) sq. root of R
- b) sq. of R
- c) logarithmic of R
- d) Reciprocal of R.

$$\Rightarrow x_{n+1} = x_n = \alpha.$$

$$\therefore \alpha = \frac{1}{2} \left[ \alpha + \frac{R}{\alpha} \right].$$

$$\Rightarrow 2\alpha - \alpha = \frac{R}{\alpha}$$

$$\Rightarrow \alpha^2 = R$$

$$\Rightarrow \alpha = \sqrt{R}$$

→ used to compute sq. root of no.

ex:-  $x_{n+1} = \frac{1}{2} \left[ x_n + \frac{117}{x_n} \right]$  converges to  $\sqrt{117}$ .

Q  $f(x) = x^3 - x^2 + 4x - 4 = 0$

$$x_0 = 2$$

$$x_1 = \underline{\hspace{2cm}}$$

a)  $\frac{2}{3}$ , b)  $\frac{4}{3}$ , c) 1, d)  $\frac{3}{2}$ .

$$\rightarrow f'(x) = 3x^2 - 2x + 4$$

$$x_1 = 2 - \left[ \frac{8^2}{\frac{12}{3}} \right]$$

$$= \frac{4}{3}$$

$$=$$

Q. The recursion relation  $x = e^{-x}$  is to be solved by Newton Raphson method, then which of the following is true?

- a)  $x_{n+1} = e^{-x_n}$   
 b)  $x_{n+1} = x_n - e^{-x_n}$   
 c)  $x_{n+1} = \frac{(1+x_n)e^{-x_n}}{(1+e^{-x_n})}$   
 d)  $x_{n+1} = \frac{x_n^2 - e^{-x_n}(1+x_n) - 1}{x_n - e^{-x_n}}$

→  $f(x) = x - e^{-x}$   
 $f'(x) = 1 + e^{-x}$   
 $f'(x_n) = 1 + e^{-x_n}$  → This will come into denominator.  
 ∴ option c

Q. Consider the series —

$$x_{n+1} = \frac{x_n}{2} + \frac{9}{8x_n}$$

$x_0 = 0.5$  is obtained from the Newton Raphson method.

The series converges to —

→  $x_{n+1} = x_n = \alpha$  (Take it always when given Series Converges)

$$\Rightarrow \alpha = \frac{\alpha}{2} + \frac{9}{8\alpha}$$

$$\Rightarrow \alpha = \frac{4\alpha^2 + 9}{8\alpha}$$

$$\Rightarrow 4\alpha^2 = 9$$

$$\Rightarrow \alpha^2 = \frac{9}{4}$$

$$\Rightarrow \alpha = \sqrt{9/4} = \frac{3}{2} = 1.5 \text{ Ans}$$

Q. The sol<sup>n</sup> of the variables  $x_1$  and  $x_2$  for the equations -

$$U = 10x_2 \sin x_1 - 0.8 = 0 \quad \text{--- ①}$$

$$V = 10x_2^2 - 10x_2 \cos x_1 - 0.6 = 0 \quad \text{--- ②}$$

is to be solved by the Newton Raphson method.

Assuming the initial value,  $x_1 = 0.0$

Jacobian Matrix =  $\dots$   $x_2 = 1.0$ .

→ Jacobian Matrix - 
$$\begin{bmatrix} \frac{\partial U}{\partial x_1} & \frac{\partial U}{\partial x_2} \\ \frac{\partial V}{\partial x_1} & \frac{\partial V}{\partial x_2} \end{bmatrix}$$


$$\frac{\partial U}{\partial x_1} = 10x_2 \cos x_1 = 10(1)(1) = 10$$

$$\frac{\partial U}{\partial x_2} = 10 \sin x_1 = 10(0) = 0.$$

$$\frac{\partial V}{\partial x_1} = 10x_2 \sin x_1 = 10(1)(0) = 0.$$

$$\frac{\partial V}{\partial x_2} = 20x_2 - 10 \cos x_1 = 20(1) - 10(1) = 10.$$

## # Numerical Integration:-



$$h = \frac{b-a}{n}$$

### ① Trapezoidal Rule:-

$$\int_a^b f(x) dx = \frac{h}{2} [(y_0 + y_n) + 2(y_1 + y_2 + y_3 + \dots + y_{n-1})]$$

### ② Simpson's $\frac{1}{3}$ rd Rule:-

$$\int_a^b f(x) dx = \frac{h}{3} [(y_0 + y_n) + 4(y_1 + y_3 + y_5 + \dots) + 2(y_2 + y_4 + y_6 + \dots)]$$

### ③ Simpson's $\frac{3}{8}$ th Rule:-

$$\int_a^b f(x) dx = \frac{3h}{8} [(y_0 + y_n) + 3(y_1 + y_2 + y_4 + y_5 + y_7 + \dots) + 2(y_3 + y_6 + y_9 + \dots)]$$

### Note:-

The accuracy of Trapezoidal rule or order of integration of Trapezoidal rule is  $O(h^2)$  → accuracy up to 2<sup>nd</sup> place of decimal

The accuracy of Simpson's  $\frac{1}{3}$ rd rule =  $O(h^4)$  → accuracy up to 4<sup>th</sup> place of decimal

The accuracy of Simpson's  $\frac{3}{8}$ th rule is  $O(h^5)$  → accuracy up to 5<sup>th</sup> place of decimal

$$9. \int_0^1 \frac{1}{1+x^2} dx.$$

By Trapezoidal & Simpson's rule and dividing the interval into 4 equal parts. and Derive the value of  $\pi$ .

$$\rightarrow \int_0^1 \frac{1}{1+x^2} dx$$

$$h = \frac{b-a}{n} = \frac{1-0}{4} = \frac{1}{4} = 0.25$$

$$y = \frac{1}{1+x^2}$$

x	0	0.25	0.50	0.75	1.
y	1.0000	0.9412	0.8000	0.6400	0.5000

$$\frac{1}{1+\frac{1}{16}}$$

$$\frac{1}{1+\frac{1}{4}}$$

$$\frac{1}{1+\frac{9}{16}}$$

$$\therefore \int_0^1 \frac{1}{1+x^2} dx = \frac{0.25}{2} \left[ (1.0000 + 0.5000) + 2(0.9412 + 0.8000 + 0.6400) \right]$$

$$\tan^{-1} x \Big|_0^1 = 0.7828$$

$$\tan^{-1} 1 - \cancel{\tan^{-1} 0} = 0.7828$$

$$\Rightarrow \frac{\pi}{4} = 0.7828$$

$$\Rightarrow \pi = 3.1312 \quad \text{we know } \pi = 3.1415$$

difference occurred at the place.

Simpson  $\frac{1}{3}$ rd Rule —

$$\int_0^1 \frac{1}{1+x^2} dx = \frac{0.25}{3} \left[ (1.0000 + 0.5000) + 4(0.9412 + 0.6400) + 2(0.8000) \right]$$

$$\tan^{-1} \Big|_0^1 = 0.7854$$

$$\Rightarrow \tan^{-1} \Big|_{\cancel{\tan 0}^0} = 0.7854$$

$$\Rightarrow \frac{\pi}{4} = 0.7854$$

$$\Rightarrow \pi = 0.7854 \times 4$$

$$\pi = 3.1416 \quad \text{we know } \pi = 3.1415$$

diff. occur at 4<sup>th</sup> place.

By Shortcut  
methods.

Q.  $\int_0^6 \frac{1}{1+x^2} dx$  by Trapezoidal rule by dividing the interval into 6 equal parts.

a) 1.4132    b) 1.3756    c) 1.2326    d) 1.0986.

→

$$\int_0^6 \frac{1}{1+x^2} dx$$

$$= \tan^{-1} 6 - \cancel{\tan^{-1} 0} \rightarrow 0$$

$$= 1.40 \dots$$

Compare with option → diff<sup>n</sup> will be at 2<sup>nd</sup> place.

∴ option (a) is correct.

29)  $\int_1^3 \log_e x \, dx$ ,  $n=2 = \underline{\hspace{2cm}}$  By Simpson's  $\frac{1}{4}$ th rule.

- a) 0.50
- b) 0.80
- c) 1.00
- d) 1.29

→ Directly integrate and Compare ~~the~~ value at 4<sup>th</sup> place will differ.

3)  $\int_0^{\pi/4} \frac{\sin x}{\cos^3 x} \, dx$ ,  $n=4$ , By Simpson's  $\frac{3}{8}$  rule.

- a) 0.3883
- b) 0.4344
- c) 0.5026
- d) 0.5589

→  $\int_0^{\pi/4} \frac{\sin x}{\cos^3 x} \, dx = \int_0^{\pi/4} \frac{\sin x}{\cos x} \cdot \frac{1}{\cos^2 x} \, dx = \int_0^{\pi/4} \tan x \cdot \sec^2 x \, dx$   
 $= \left( \frac{\tan^2 x}{2} \right)_0^{\pi/4}$   
 $= 0.5000$

4)  $\int_1^3 \frac{1}{x} \, dx$ ,  $n=2$ , By Simpson's  $\frac{1}{3}$ rd rule.

- a) 1.000
- b) 1.098
- c) 1.111
- d) 1.120

→ Comparison ⇒ Use only Numeric method  
 Do not ever integrate or shortcut method

→  $y = \frac{1}{x}$

x	1	2	3
y	1	0.50	0.333

⇒  $\int_1^3 \frac{1}{x} \, dx = \frac{1}{3} [1.333 + 4(0.5)]$   
 $= \frac{3.333}{3} = 1.111$

Q. Torque Exerted on a flywheel over a cycle is listed in the table —

Angle ( $^{\circ}$ )	$0^{\circ}$	$60^{\circ}$	$120^{\circ}$	$180^{\circ}$	$240^{\circ}$	$300^{\circ}$	$360^{\circ}$
Torque (Nm)	0	1066	323	0	-323	-355	0
	$\gamma_0$	$\gamma_1$	$\gamma_2$	$\gamma_3$	$\gamma_4$	$\gamma_5$	$\gamma_6$

Estimate the value of the fn by Simpson  $\frac{1}{3}$  rule.

- (A) 542    
 (B) 993    
 (C) 1444    
 (D) 1986

$$\rightarrow \int_0^{2\pi} f(x) dx$$

$$= \frac{60^{\circ}}{3} \left[ (0+0) + 4(1066 + 0 - 355) + 2(323 - 323) \right]$$

$$= \frac{\pi}{3} \times 4(711)$$

$$= \frac{\pi}{9} \times 2844$$

$$= 992.7$$

$$\approx 993.$$

# # Numerical Sol<sup>n</sup> of Ordinary Differential Eq<sup>n</sup>:-

$$\frac{dy}{dx} = f(x, y), \quad x = x_0, \quad y = y_0, \quad y(x_1) = \underline{\hspace{2cm}}$$

## Methods -

- ① Taylor's series method
- ② Picard's method
- ③ Runge - Kutta method.

## Taylor Series -

$$y = y_0 + (x-x_0)(y')_0 + \frac{(x-x_0)^2}{2!} (y'')_0 + \frac{(x-x_0)^3}{3!} (y''')_0 + \dots \infty$$

Q.  $\frac{dy}{dx} = -xy$ ,  $y(0) = 1$ ,  $y(0.1) = \underline{\hspace{2cm}}$

$$\begin{aligned} x+y &= 1 \\ y &= 1-x \\ &= 1-0.1 \end{aligned}$$

→

$$y' = -xy$$

$$y'' = -xy' - y \cdot 1$$

$$y''' = -xy'' - 2y'$$

$$y^{IV} = -xy^{III} - 3y''$$

$$y = 1 + x(0) + \frac{x^2}{2!} (-1) + \frac{x^3}{3!} (0) + \frac{x^4}{4!} (3) + \dots$$

$$= 1 - \frac{x^2}{2} + \frac{x^4}{8} \xrightarrow{\text{Consider only upto 4th order rest can be neglected}}$$

$$y(0.1) = 1 - \frac{0.01}{2} + \frac{0.0001}{8} \text{ neglected}$$

$$= 1 - 0.005$$

$$= 0.995 \quad \underline{\underline{\text{Ans}}}$$

Q. Picard's method:-

$$\frac{dy}{dx} = f(x, y), \quad x = x_0, \quad y = y_0.$$

$$\Rightarrow \int_{y_0}^y dy = \int_{x_0}^x f(x, y) dx.$$

$$\Rightarrow y - y_0 = \int_{x_0}^x f(x, y) dx$$

$$\Rightarrow y = y_0 + \int_{x_0}^x f(x, y) dx.$$

$$y_1 = y_0 + \int_{x_0}^x f(x, y_0) dx$$

$$y_2 = y_0 + \int_{x_0}^x f(x, y_1) dx$$

$$y_3 = y_0 + \int_{x_0}^x f(x, y_2) dx.$$

-----  
-----

$$y = ( \quad ) + \dots$$

Q.  $\frac{dy}{dx} = -xy$ ,  $y(0) = 1$ ,  $y(0.1) = \underline{\quad}$

$$y_1 = 1 + \int_0^x -x(1) dx$$

$$y_1 = (1 - \frac{x^2}{2})$$

$$y_2 = 1 + \int_0^x -x(1 - \frac{x^2}{2}) dx$$

$$y_2 = 1 - \frac{x^2}{2} + \frac{x^4}{8}$$

$$\therefore y = 1 - \frac{x^2}{2} + \frac{x^4}{8} + \dots$$

Q.  $\frac{dy}{dx} = \frac{x^2}{1+y^2}$ ,  $y(0) = 0$ ,  $y(0.25) = \underline{\hspace{2cm}}$

$$\rightarrow y_1 = 0 + \int \frac{x^2}{1+0^2} = \frac{x^3}{3}$$

$$y_2 = 0 + \int \frac{x^2}{1+(\frac{x^3}{3})^2} dx$$

$$= \tan^{-1}\left(\frac{x^3}{3}\right)$$

$$y_3 = 0 + \int \frac{x^2}{1+(\tan^{-1}\frac{x^3}{3})^2} dx \rightarrow \int \text{becomes difficult}$$

$\therefore$  Take prev. one as approx soln.

$$\triangle y = \tan^{-1} \frac{x^3}{3}$$

$$y(0.25) = \tan^{-1} \frac{(0.25)^3}{3}$$

When derivatives are convenient  
Taylor Series

When Taylor Ser.  $\int$  is convenient  
Picard's method

Q.  $\frac{dy}{dx} = -xy$ ,  $y(0) = 1$ , &  $y(0.1) = \underline{\hspace{2cm}}$

$$\frac{dy}{dx} + xy = 0$$

$$\text{I.F.} = e^{\int p dx} = e^{\int x dx} = e^{\frac{x^2}{2}}$$

$$y \cdot e^{\frac{x^2}{2}} = c$$

$$1 \cdot e^0 = c \Rightarrow c = 1$$

$$y e^{\frac{x^2}{2}} = 1$$

$$y = \frac{1}{e^{\frac{x^2}{2}}} = e^{-\frac{x^2}{2}} = e^{-\frac{(0.1)^2}{2}} = e^{-\frac{0.01}{2}} = e^{-0.005}$$

$$= \underline{\underline{0.995}}$$

## # Runge-Kutta Method:-

$$y = y_0 + \frac{1}{6}(k_1 + 2k_2 + 2k_3 + k_4)$$

Where  $k_1 = h \cdot f(x_0, y_0)$   $\longrightarrow$  Where  $h = x_1 - x_0$ .

$$k_2 = h f\left(x_0 + \frac{h}{2}, y_0 + \frac{k_1}{2}\right)$$

$$k_3 = h f\left(x_0 + \frac{h}{2}, y_0 + \frac{k_2}{2}\right)$$

$$k_4 = h f(x_0 + h, y_0 + k_3)$$

Q. ①  $\frac{dy}{dx} = \frac{f(x,y)}{x+y}$   $y(x_0) = y_0$ ,  $y(x_1) = \underline{\hspace{2cm}}$

$$h = x_1 - x_0 = 0.1 - 0 = 0.1$$

$$k_1 = h f(x_0, y_0) = 0.1(0+1) = 0.1$$

$$k_2 = h f\left(x_0 + \frac{h}{2}, y_0 + \frac{k_1}{2}\right) = 0.1[0.05 + 1.05] = 0.11$$

$$k_3 = h f\left(x_0 + \frac{h}{2}, y_0 + \frac{k_2}{2}\right) = 0.1[0.05 + 1.055] = 0.1105$$

$$k_4 = h f(x_0 + h, y_0 + k_3) = 0.1[0.1 + 1.1105] = 0.12105$$

Q Match the following —

- |                       |                   |   |
|-----------------------|-------------------|---|
| E: Newton Raphson     | $\longrightarrow$ | 1) Sol <sup>n</sup> of non linear eq <sup>n</sup>                               |
| F: Runge Kutta        | $\longrightarrow$ | 2) Sol <sup>n</sup> of linear algebraic   |
| G: Simpson rule       | $\longrightarrow$ | 3) Sol <sup>n</sup> of O.D. eq <sup>n</sup> .                                   |
| H: Gauss Elimination. | $\longrightarrow$ | 4) Sol <sup>n</sup> of linear algebraic eq <sup>n</sup><br>Numeric Integration. |

## Complex Variable

$$\textcircled{1} \begin{cases} z = x + iy \\ \bar{z} = x - iy \\ z\bar{z} = |z|^2 = x^2 + y^2 \end{cases}$$

$\textcircled{2} \quad x + iy = r(\cos\theta + i\sin\theta) = re^{i\theta}$  is called the modulus-Amplitude form  
or  
Polar form

$$r = |z| = \sqrt{x^2 + y^2}$$

$$\theta = \tan^{-1}\left(\frac{y}{x}\right)$$


$$\textcircled{3} \quad \cos\theta + i\sin\theta = e^{i\theta}$$

$$(\cos\theta + i\sin\theta)^n = \cos n\theta + i\sin n\theta$$

$$1 = \cos 0^\circ + i\sin 0^\circ = e^{i0}$$

$$i = \cos \frac{\pi}{2} + i\sin \frac{\pi}{2} = e^{i\pi/2}$$

$$1 + i = \sqrt{2} \left[ \cos \frac{\pi}{4} + i\sin \frac{\pi}{4} \right] = \sqrt{2} e^{i\pi/4}$$



Q.  $(i)^i = \underline{\hspace{2cm}}$  where  $i = \sqrt{-1}$

$(z)^z = \underline{\hspace{2cm}}$  where  $z = \sqrt{-1}$

$$(i)^i = (e^{i\pi/2})^i = \underline{\underline{e^{-\pi/2}}}$$

Q.  $\textcircled{1} \quad (1+i)(2-5i) = \underline{\hspace{2cm}}$

$\textcircled{2} \quad \left| \frac{3+4i}{1-2i} \right| = \underline{\hspace{2cm}}$

$\textcircled{3} \quad \frac{i+3}{i+1} = \underline{\hspace{2cm}}$

$\textcircled{4} \quad \frac{1+2i}{i-2} = \underline{\hspace{2cm}}$

$\textcircled{5} \quad \frac{-5+i10}{3+i4} = \underline{\hspace{2cm}}$



Q. The value of  $(1+i)^8 =$  \_\_\_\_\_

Ⓐ 2 Ⓑ 4 Ⓒ 8 Ⓓ = 16.

$$\begin{aligned} &\rightarrow (1+i)^8 \\ &= [(1+i)^2]^4 \\ &= [\cancel{1+1} + 2i]^4 \\ &= 16i^4 \\ &= 16 \underline{\text{Ans}} \end{aligned}$$

Q. The Polar form of  $2+2i$  is \_\_\_\_\_

$$r = |z| = \sqrt{4+4} = 2\sqrt{2}.$$

$$\theta = \tan^{-1} 1 = \frac{\pi}{4}$$

$$\therefore 2+2i = 2\sqrt{2} e^{i\frac{\pi}{4}}$$

Q.

## # Analytic function:-

A single valued fn which is defined and differentiable at each point of domain  $D$  is said to be an analytic fn in that domain.

Note:-

The necessary and sufficient cond<sup>n</sup> for a function

$f(z) = u(x, y) + i\{v(x, y)\}$  to be analytic is it should

satisfy the Cauchy-Riemann eq<sup>n</sup>.

Cond<sup>n</sup> to be  
Analytic

$$\begin{aligned} \frac{\partial u}{\partial x} &= \frac{\partial v}{\partial y} \\ \frac{\partial u}{\partial y} &= -\frac{\partial v}{\partial x} \end{aligned}$$

$$\begin{aligned} u_x &= v_y \\ u_y &= -v_x \end{aligned}$$

Q.

$$f(x) = (x^2 - y^2) + i(2xy)$$

$$u = x^2 - y^2, \quad v = 2xy.$$

$$\frac{\partial u}{\partial x} = 2x \quad , \quad \frac{\partial v}{\partial y} = 2x$$

$$\frac{\partial u}{\partial y} = -2y \quad , \quad \frac{\partial v}{\partial x} = 2y$$

Satisfying C.R eq<sup>n</sup>

∴ Analytic in nature

## # Harmonic function:-

— Any function of  $x$  and  $y$  satisfies the Laplace's eq<sup>n</sup> is said to be an harmonic function.

↓ represented by

$$\boxed{\nabla^2 u = 0}$$

ex: ①  $u = x^2 - y^2$

$$\left. \begin{aligned} \frac{\partial u}{\partial x} &= 2x \\ \frac{\partial^2 u}{\partial x^2} &= 2 \end{aligned} \right| \begin{aligned} \frac{\partial u}{\partial y} &= -2y \\ \frac{\partial^2 u}{\partial y^2} &= -2 \end{aligned}$$

$$\nabla^2 u = \frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} = 2 - 2 = 0 \therefore \text{Harmonic } \checkmark$$

Note:-

Real and Imj part of an Analytic eq<sup>n</sup> satisfies the Laplacian Equation.

i.e.,  $\frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} = 0$

# If  $f(z) = u(x, y) + iv(x, y)$  ~~then~~ If  $u(x, y)$  <sup>Real Part</sup> given

then  $f(z) = 2u\left(\frac{z}{2}, \frac{z}{2i}\right) - u(0, 0) + ci$

If Real part Given then asked directly  $f(z)$  then how to get

Q.  $u = x^2 - y^2$

$$\begin{aligned} u\left(\frac{z}{2}, \frac{z}{2i}\right) &= \left(\frac{z}{2}\right)^2 - \left(\frac{z}{2i}\right)^2 \\ &= \frac{z^2}{4} + \frac{z^2}{4} \\ &= \frac{z^2}{2} \end{aligned}$$

$$u(0, 0) = 0 - 0 = 0.$$

$$f(z) = 2 \times \frac{z^2}{2} - 0 + ci = z^2 + ci$$

$$u = x^3 + 3xy^2 + 3x + 1.$$

$$u\left(\frac{z}{2}, \frac{z}{2i}\right) = \frac{z^3}{8} - \frac{3z}{2} \left(\frac{z^2}{4i^2}\right) + \frac{3z}{2} + 1$$

$$= \frac{z^3}{8} + \frac{3z^3}{8} + \frac{3z}{2} + 1$$

$$= \cancel{\frac{z^3}{8}} + \frac{3z^3}{8} + \frac{3z}{2} + 1$$

$$u(0,0) = 0 - 0 + 0 + 1 = 1.$$

$$f(z) = 2\left(\frac{z^3}{8} + \frac{3z}{2} + 1\right) - 1 + Ci$$

$$f(z) = z^3 + 3z + 1 + Ci$$

→

$$Q. f(z) = z^2$$

$$= (x+iy)^2$$

$$= x^2 + i^2y^2 + 2ixy$$

$$= x^2 - y^2 + 2ixy$$

If real part given, how to get Imj part.

$$dv = \frac{\partial v}{\partial x} dx + \frac{\partial v}{\partial y} dy$$

$$dv = -\frac{\partial u}{\partial y} dx + \frac{\partial u}{\partial x} dy$$

which is exact.

$$\text{Let } u = x^2 - y^2$$

$$\frac{\partial u}{\partial x} = 2x, \quad \frac{\partial u}{\partial y} = -2y$$

$$dv = 2y dx + 2x dy$$

↳ exact  
no need to  
check

Gen. soln —

$$v = \int 2y dx + \int 0 dy$$

$$v = 2xy + C$$

$$a. \quad u = x^3 - 3xy^2 + 3x + 1$$

$$\frac{\partial u}{\partial x} = 3x^2 - 3y^2 + 3$$

$$\frac{\partial u}{\partial y} = -6xy$$

$$dv = \int 6xy \, dx + \int (\cancel{3x^2 - 3y^2 + 3}) \, dy$$
$$=$$

$$b. \quad u = e^x \cos y$$

$$\frac{\partial u}{\partial x} = e^x \cos y$$

$$\frac{\partial u}{\partial y} = -e^x \sin y$$

$$dv = -\frac{\partial u}{\partial y} \, dx + \frac{\partial u}{\partial x} \, dy$$

$$\int dv = \int e^x \sin y \, dx + \int \cancel{e^x} \cos y \, dy$$

$$\boxed{v = e^x \sin y}$$

\* If imag part given and asked real part then how to get?

$$du = \frac{\partial u}{\partial x} dx + \frac{\partial u}{\partial y} dy.$$

or

$$\boxed{du = \frac{\partial v}{\partial y} dx - \frac{\partial v}{\partial x} dy}$$

which is exact.

ex:-

①  $v = -xy$

$$\frac{\partial v}{\partial x} = -y, \quad \frac{\partial v}{\partial y} = -x.$$

$$\int du = \int -x dx + \int y dy.$$

$$u = -\frac{x^2}{2} + \frac{y^2}{2} = \frac{1}{2} (y^2 - x^2).$$

② Given  $v = y^3 - 3x^2y$ , find  $u$ . where  $f(z)$  is harmonic

$$\frac{\partial v}{\partial x} = -6xy$$

$$\frac{\partial v}{\partial y} = 3y^2 - 3x^2$$

$$\int du = \int (3y^2 - 3x^2) dx + \int ~~6xy~~ dy$$

$$\boxed{u = 3xy^2 - x^3}$$

## # Singular Points! → or Pole

Singular points are those at which the given function  $f(z)$  is not analytic.

These points are also called Singularities or poles.

find Singular Points.

$$Q. ① \quad f(z) = \frac{1-2z}{z(z-1)(z-2)}$$

$$z = 0, 1, 2.$$

$$Q. \quad f(z) = \frac{z}{z^2 - z - 2}$$

$$f(z) = \frac{z}{(z-2)(z+1)}$$

$$\Rightarrow z = 2, -1.$$

$$Q. \quad f(z) = \frac{z}{z^4 - 1}$$

$$z = \pm 1, \pm i$$

## # Cauchy's Theorem:-


If  $f(z)$  is an analytic fn and its derivative  $f'(z)$  is continuous at all points inside and on a simple and closed curve  $C$ , then-

$$\oint_C f(z) dz = 0$$

means

if pole outside the limit, then  $\int = 0$


Q.  $\oint_C \frac{3z^2 + 7z + 1}{z+1} dz$ , where  $C$  is  $|z| = \frac{1}{2}$



$\therefore$  pole is outside curve.

$\therefore$  Ans is 0.

Q.  $\oint_C \frac{z+4}{z^2+2z+5} dz$  where  $C$  is  $|z+1| = 1$




Pole  $\Rightarrow z^2 + 2z + 5 = 0$

$$\Rightarrow \frac{-2 \pm \sqrt{4-20}}{2} = \frac{-2 \pm 4i}{2} = -1 \pm 2i,$$

Pole =  $-1 + 2i, -1 - 2i \Rightarrow$  outside the curve  $\Rightarrow$  Ans = 0

Q. #  $\oint_C \frac{-3z+4}{z^2+4z+5} dz$  Where  $C$  is  $|z|=1$



$$z^2 + 4z + 5 = 0$$

$$\frac{-4 \pm \sqrt{16 - 20}}{2}$$

$$= \frac{-4 \pm 2i}{2}$$

$$= -2 \pm i$$

↪ Pole outside the curve.  
 $\therefore$  Answer is 0.

### # Cauchy's Integral formula

V.V. Imp  
2 marks

If  $f(z)$  is analytic for inside and on a simple and closed curve 'C' and if  $a$  is any point inside C then,

$$\oint_C \frac{f(z)}{z-a} dz = 2\pi i f(a)$$

$$\oint_C \frac{f(z)}{(z-a)^2} dz = 2\pi i f'(a)$$

$$\oint_C \frac{f(z)}{(z-a)^3} dz = \frac{2\pi i}{2!} f''(a)$$


$$\oint_C \frac{f(z)}{(z-a)^{n+1}} dz = \frac{2\pi i}{n!} f^{(n)}(a)$$

Q. ① The value of  $\oint_C \frac{\cos z}{z} dz$  where  $C$  is an ellipse

$$9x^2 + 4y^2 = 1$$

$$\rightarrow 9x^2 + 4y^2 = 1$$


$$\frac{x^2}{(\frac{1}{3})^2} + \frac{y^2}{(\frac{1}{2})^2} = 1$$



∴ Pole inside  
 ∴ Cauchy Integral formula applicable

$$\begin{aligned} \therefore \oint \frac{\cos z}{z} dz &= 2\pi i \cos z \Big|_{z=0} \\ &= 2\pi i \underline{\underline{\text{Ans}}} \end{aligned}$$

Q.  $\oint_C \frac{1}{z^2-1} dz$  where  $C$  is the circle  $x^2+y^2=4$ .



Method - 1

Partial fraction -

$$\begin{aligned}\oint_C \frac{1}{(z-1)(z+1)} dz &= \frac{1}{2} \oint_C \left[ \frac{1}{z-1} - \frac{1}{z+1} \right] dz \\ &= \frac{1}{2} \left( \cancel{2\pi i(1)} - \cancel{2\pi i(1)} \right) \\ &= \underline{\underline{0}}\end{aligned}$$

Observation:-

Sometimes even also pole inside then  $\int = 0$

Method - 2

Shortcut


$$\begin{aligned}\oint_C \frac{1}{z^2-1} dz &= \oint_C \frac{1}{(z-1)(z+1)} dz \\ &= \oint_C \frac{\left(\frac{1}{z+1}\right)}{(z-1)} + \frac{\left(\frac{1}{z-1}\right)}{(z+1)} dz \\ &= 2\pi i \left(\frac{1}{z+1}\right)_{z=1} + 2\pi i \left(\frac{1}{z-1}\right)_{z=-1} \\ &= \cancel{2\pi i\left(\frac{1}{2}\right)} + \cancel{2\pi i\left(-\frac{1}{2}\right)} \\ &= \underline{\underline{0}}\end{aligned}$$

Q.  
Repeated  
2 times


$$\oint_C \frac{z-1}{(z+1)^2(z-2)} dz$$

where C is  $|z-i|=2$

If one pole inside  
another outside.  
then?



यहाँ बाहर है उसको बाहर ले जाना है  
और inside में उसको नीचे ले जाना है



$$\oint_C \frac{z-1}{(z+1)^2(z-2)} dz$$

$$= \oint_C \frac{f(z)}{(z+1)^2} dz$$


$$f'(z) = \frac{-1}{(z-2)^2}$$

$$= 2\pi i \left[ \frac{-1}{(z-2)^2} \right]_{z=-1}$$

$$= \frac{-2\pi i}{(-1-2)^2}$$

$$= \frac{-2\pi i}{9}$$

10.  $\oint_C \frac{z^2 - 4}{z^2 + 4} dz$  where  $C$  is the circle  $|z - i| = 2$



$$\int \frac{z^2 - 4}{z^2 + 4} dz = \int \frac{z^2 - 4}{(z - 2i)(z + 2i)} dz$$

$$= 2\pi i \left[ \frac{z^2 - 4}{z + 2i} \right]_{z = 2i}$$

$$\oint \frac{z^2 - 4}{(z - 2i)(z + 2i)} dz$$


$$= \oint \frac{\frac{z^2 - 4}{z + 2i}}{z - 2i} dz$$

$$= 2\pi i \left( \frac{z^2 - 4}{z + 2i} \right)_{z = 2i}$$

$$= 2\pi i \left( \frac{-4 - 4}{2i + 2i} \right) = 2\pi i \left( \frac{-8}{4i} \right)$$

$$= -4\pi$$

Q.  $\oint_c \frac{z^2}{z^2+1} dz$      $c$  is  $|z+1|=1$



$$\oint \frac{z^2}{(z^2-1)(z^2+1)} = \oint \frac{z^2}{\underbrace{(z-1)(z+1)(z-i)(z+i)}_{\text{outside pole}}}$$

inside की संख्या  
outside की संख्या


$$\oint \frac{z^2}{(z-1)(z^2+1)} dz$$

$$= 2\pi i \left( \frac{z^2}{(z-1)(z^2+1)} \right)_{z=-1}$$

$$= 2\pi i \left( \frac{1}{-2 \times 2} \right)$$

$$= -\frac{\pi i}{2}$$

Q.  $\oint \frac{5z^2 + 7z + 3}{(z-1)^3}$  where  $C$  is  $|z| = 1$



$$\oint \frac{5z^2 + 7z + 3}{(z-1)^3}$$

$$f(z) = 5z^2 + 7z + 3.$$

$$f'(z) = 10z + 7.$$

$$f''(z) = 10.$$

$$\therefore \oint \frac{5z^2 + 7z + 3}{(z-1)^3} = \frac{2\pi i}{2!} (10)$$

$$= 10\pi i$$

\*\*\*

# Residue at  $z=a$  where  $a$  is the pole of order  $n$ .

$$\lim_{z \rightarrow a} \frac{1}{(n-1)!} \frac{d^{n-1}}{dz^{n-1}} \left[ (z-a)^n \cdot f(z) \right]$$

Q. → find residue of  $f(z) = \frac{1-2z}{z(z-1)(z-2)}$

at  $z=0$ ,  $\frac{1-0}{(0-1)(0-2)} = \frac{1}{2} =$

at  $z=1$ ,  $\frac{1-2}{1 \times 1} = \underline{\underline{1}}$

at  $z=2$ ,  $\frac{1-4}{2 \times 1} = \underline{\underline{-\frac{3}{2}}}$

Q. ① find residue of fn.  $f(z) = \frac{z^2}{(z-1)(z+1)(z-2)}$

$$\text{at } z=1 \quad \frac{1}{2 \times -1} = -\frac{1}{2}$$

$$z=-1 \quad \frac{1}{-2 \times -3} = \frac{1}{6}$$

$$z=2, \quad \frac{4}{1 \times 3} = \frac{4}{3}$$

Q. ②  $f(z) = \frac{1}{z(z+2)^3}$

$$\frac{1}{z(z+2)^3} = \frac{A}{z} + \frac{B}{(z+2)^2} + \frac{C}{(z+2)^3}$$

Ans  $(z+2)^3 \rightarrow$   $\lim_{z \rightarrow -2} \frac{1}{2!} \frac{d^2}{dz^2} \left[ (z+2)^3 \cdot \frac{1}{z(z+2)^3} \right]$

$$= \frac{1}{2} \frac{d^2}{dz^2} \left( \frac{1}{z} \right)$$

$$= \frac{1}{2} \cdot \frac{2}{z^3}$$

$$= \frac{1}{(-2)^3} = \frac{1}{8}$$

Workbook

$$\textcircled{1} \quad A = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \quad \Delta = 1 \quad \text{or} \quad \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} \quad \Delta = -1$$

$$\Delta = \underline{\underline{\pm 1}}$$

$$\textcircled{2} \quad \begin{aligned} XY &= Y \\ YX &= X \\ X^2 + Y^2 &= ? \end{aligned}$$

$$= XX + YY$$

$$= \cancel{(YX)X} + \cancel{(XY)Y}$$

$$= (YX)(YX) + (XY)(XY)$$

$$= Y(XY) + X(XY)$$

$$= YX + XY$$

$$= X + Y \quad \underline{\underline{Ans}}$$

$$\textcircled{7} \quad \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \\ \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} \begin{bmatrix} 1 & 1 \\ 1 & 0 \end{bmatrix} \begin{bmatrix} 0 & 1 \\ 1 & 1 \end{bmatrix} \begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix}$$

Total 8

$$\textcircled{8} \quad A = [a_{ij}]_{m \times n}$$

$$a_{ij} = i + j$$

$$A_{2 \times 2} = \begin{bmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{bmatrix} = \begin{bmatrix} 2 & 3 \\ 3 & 4 \end{bmatrix}$$

$\rightarrow$  12 Check with options

9)  $4^4$   
 $3 \times 4^4$

10) wrong 0

11) 10

$$\begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & 1 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} 0 & 0 \\ 1 & 1 \end{bmatrix}$$

~~12)~~

$$\begin{bmatrix} 1 & 0 \\ 1 & 0 \end{bmatrix} \begin{bmatrix} 0 & 1 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix} \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} 0 & 0 \\ 1 & 0 \end{bmatrix}$$

13) 14)  $5! = \underline{\underline{120}}$

16)

$$a_{ij} \begin{bmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{bmatrix}$$

$$= \begin{bmatrix} 0 & -3 & -8 \\ 3 & 0 & -5 \\ 8 & 5 & 0 \end{bmatrix}$$

$\therefore \Delta = 0$

$\therefore$  inverse not possible.

~~17)~~

18)

19)

Correct  $2x + 3y + z = \lambda z \rightarrow 3^{\text{rd}}$

38

$$\left[ \frac{1}{1 \cdot 2} \quad 0 \quad \dots \right]$$

$$\frac{1}{1 \cdot 2} + \frac{1}{2 \cdot 3} + \frac{1}{3 \cdot 4} + \dots + \frac{1}{n(n+1)} \dots$$

$$\begin{aligned} S_n &= \sum T_n \\ &= \sum \frac{1}{n(n+1)} \\ &= \sum \frac{1}{n} - \frac{1}{n+1} \\ &= \frac{1}{1} - \frac{1}{2} + \frac{1}{2} - \frac{1}{3} + \frac{1}{3} - \frac{1}{4} + \dots + \frac{1}{n} - \frac{1}{n+1} \\ &= 1 - \frac{1}{n+1} \quad \underline{\underline{A}} \end{aligned}$$

39

$$1 \cdot \frac{1}{2} \cdot \frac{1}{3} \cdot \frac{1}{4} \dots \frac{1}{n}$$

$$= \frac{1}{n!} \quad \underline{\underline{A}}$$

40

$$A = \begin{bmatrix} 40 & -29 & -11 \\ -18 & 30 & -12 \\ 26 & 24 & -50 \end{bmatrix}$$

$$\lambda_1, \lambda_2 = ?$$

$$\lambda_1 + \lambda_2 + \lambda_3 = 20$$

$$\lambda_2 = 20 - \lambda_1 \quad \underline{\underline{A}}$$

$\Delta = 0$

45

3, 2, -1

$$\begin{aligned}
 B &= A^2 - A \\
 &= 9 - 3 = 6 \\
 &= 2^2 - 2 = 2 \\
 &= (-1)^2 - (-1) = 2
 \end{aligned}$$

$$|B| = 6 \times 2 \times 2 = 24.$$

46

$$A = \begin{bmatrix} 1 & 0 & 0 \\ i & \frac{-1+i\sqrt{3}}{2} & 0 \\ 0 & 1+2i & \frac{-1-i\sqrt{3}}{2} \end{bmatrix}$$

Trace ?

Trace is  $\sum$  principal diagonals

$$\begin{aligned}
 (A)^{102} &= (1)^{102}, \left(\frac{-1+i\sqrt{3}}{2}\right)^{102}, \left(\frac{-1-i\sqrt{3}}{2}\right)^{102} \\
 &= 1, \left(e^{-i\frac{\pi}{3}}\right)^{102}, \left(e^{i\frac{\pi}{3}}\right)^{102} \\
 &= 1, e^{-34\pi i}, e^{34\pi i} \\
 &= 1 + 1 + 1 = 3
 \end{aligned}$$

48

$\lambda = -1, 1, 0$

$$|A^{100} + I|$$

2, 2, 1

$$|A^{100} + I| = 2 \times 2 \times 1 = \underline{\underline{4}}$$

(49)  $\lambda_1 = 1, \lambda_2 = 4$   
 $X_1 = \begin{bmatrix} 1 \\ -1 \end{bmatrix}, X_2 = \begin{bmatrix} 2 \\ 1 \end{bmatrix}$

(a)  $\begin{bmatrix} 4 & 8 \\ 5 & 9 \end{bmatrix}$

(b)  $\begin{bmatrix} 9 & 8 \\ 5 & -4 \end{bmatrix}$

(c)  $\begin{bmatrix} 9 & 2 \\ 1 & 2 \end{bmatrix}$

(d)  $\begin{bmatrix} 2 & 2 \\ 1 & 3 \end{bmatrix}$

(51)  $A = 1, -1, \lambda$   
 $A + 3I = 1+3, -1+3, \lambda+3$   
 $A + 3I = 4, 2, \lambda+3$   
 $A^2 + 3I = 1(4), -1(2), \lambda(\lambda+3)$   
 $= 4, -2, \lambda^2 + 3\lambda$

$\lambda^2 + 3\lambda = 18$

$\hookrightarrow \lambda \neq 0$

$\Delta \neq 0$

$\therefore$  Inverse exists.

Similarly  
 Inverse  
 exists for A

Correction  $\rightarrow$  2x2 not 3x3

(62)  $A^{-1} = I - 2A$

$I = A - 2A^2$

$I = \lambda - 2\lambda^2$

$2\lambda^2 - \lambda + 1 = 0$

$\Delta = \lambda_1 \lambda_2$

$= \frac{1}{2} \checkmark$

63

$$A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$$

$$a+d = ad-bc = 1$$

Then  $A^3 = ?$

$$\rightarrow \lambda_1 + \lambda_2 = 1$$

$$\lambda_1 \lambda_2 = 1$$

$$\lambda_1 + \frac{1}{\lambda_1} = 1$$

~~$$\begin{aligned} \lambda_1^2 - \lambda_1 + 1 &= 0 \\ \lambda_1^2 + \lambda_1 + 1 &= 0 \\ \lambda_1 &= \end{aligned}$$~~

$$\lambda_1^2 + 1 = \lambda_1$$

$$A^2 = A - I$$

$$A^3 = A^2 - A$$

$$= (A - I)A$$

$$= A^2 - A$$

$$= A - I - A$$

$$= -I$$

□

## Laplace Transform:

→ Let  $f(t)$  be any fn. of  $t$  defined  $\forall$  positive values of  $T$ , then the Laplace Transform of  $f(t)$  is denoted by  $\mathcal{L}[f(t)]$  and is defined as —

$$\mathcal{L}[f(t)] = \int_0^{\infty} e^{-st} \cdot f(t) dt = F(s)$$

$$\Rightarrow f(t) = \mathcal{L}^{-1}[F(s)]$$

is called the Inverse Laplace Transform of  $F(s)$ .

$$\textcircled{1} \quad \mathcal{L}(t^n) = \int_0^{\infty} t^n e^{-st} dt$$

$$\text{let } x = st \Rightarrow t = \frac{x}{s} \Rightarrow dt = \frac{dx}{s}$$

$$t \rightarrow \infty \Rightarrow x \rightarrow \infty$$

$$t \rightarrow 0 \Rightarrow x \rightarrow 0$$

$$= \int_0^{\infty} e^{-x} \left(\frac{x}{s}\right)^n \cdot \frac{dx}{s}$$

$$= \int_0^{\infty} e^{-x} \frac{x^n}{s^n} \cdot \frac{dx}{s}$$

$$= \frac{1}{s^{n+1}} \int_0^{\infty} e^{-x} x^n dx$$

$$= \frac{1}{s^{n+1}} \int_0^{\infty} e^{-x} x^{n-1} dx$$

$$= \frac{\Gamma(n+1)}{s^{n+1}}$$

$$\therefore \mathcal{L}[t^n] = \frac{n!}{s^{n+1}}, \text{ if } n \in \mathbb{N}$$

$$\mathcal{L}[t^n] = \frac{\Gamma(n)}{s^{n+1}}, \text{ if } n > 0$$

$$\therefore \begin{array}{l} \mathcal{L}(1) = \frac{1}{s} \\ \mathcal{L}(t) = \frac{1}{s^2} \\ \mathcal{L}(t^2) = \frac{2!}{s^3} \\ \vdots \end{array}$$

Improper Integral.

$$\Gamma_n = \int_0^{\infty} e^{-x} x^{n-1} dx$$

$$\Gamma_{n+1} = n\Gamma_n, \text{ if } n > 0$$

$$\Gamma_{n+1} = n!, \text{ if } n \in \mathbb{N}$$

$$\Gamma_1 = 1$$

$$\Gamma_{\frac{1}{2}} = \sqrt{\pi}$$

$$L(1) = \frac{1}{s}$$

$$L(t) = \frac{1}{s^2}$$

$$L(t^2) = \frac{2}{s^3}$$

$$L(e^{at}) = \frac{1}{s-a}$$

$$L(e^{-at}) = \frac{1}{s+a}$$

$$L(\sin at) = \frac{a}{s^2+a^2}$$

$$L(\cos at) = \frac{s}{s^2+a^2}$$

$$L(\sinh at) = \frac{a}{s^2-a^2}$$

$$L(\cosh at) = \frac{s}{s^2-a^2}$$

Q. find  $L(t^2+6t+8)$ ,  $L(\sqrt{t})$ ,  $L(\frac{1}{\sqrt{t}})$ ,  $L(\sin^2 t)$ ,  $L(\sin^3 t)$ ,  $L(\sin 2t \cdot \cos t)$

$$\& f(t) = \begin{cases} 0, & 0 < t < 2 \\ 3, & t \geq 2 \end{cases}$$

find  $L[f(t)]$ .

$$\rightarrow \textcircled{1} L(t^2+6t+8) = \frac{2}{s^3} + \frac{6}{s^2} + \frac{8}{s}$$

$$\textcircled{2} L(t^{1/2}) = \frac{\sqrt{\frac{1}{2}} + 1}{s^{1/2+1}} = \frac{\frac{1}{2}\sqrt{\frac{1}{2}}}{s^{3/2}} = \frac{\sqrt{\pi}}{2s^{3/2}}$$

$$\textcircled{3} L(t^{-1/2}) = \frac{\sqrt{-\frac{1}{2}+1}}{s^{-1/2+1}} = \frac{\sqrt{\frac{1}{2}}}{s^{1/2}} = \frac{\sqrt{\pi}}{\sqrt{s}} = \sqrt{\frac{\pi}{s}}$$

$$\textcircled{4} L(\sin^2 t) = L\left(\frac{1-\cos 2t}{2}\right) = \frac{1}{2s} - \frac{1}{2} \frac{s}{s^2+4} = \frac{1}{2s} - \frac{s}{2(s^2+4)}$$

$\sin 3t = 3\sin t - 4\sin^3 t$

$$\textcircled{5} L(\sin^3 t) = L\left(\frac{3\sin t - \sin 3t}{4}\right) = \frac{3}{4} \left[ \frac{1}{s^2+1} - \frac{3}{s^2+9} \right] = \frac{3}{4} \left[ \frac{8^2}{(s^2+1)(s^2+9)} \right] = \frac{6}{(s^2+1)(s^2+9)}$$

$$\textcircled{6} \quad L(\sin 2t \cdot \cos t) = \frac{1}{2} L[\sin 3t + \sin t]$$

$$= \frac{1}{2} \left[ \frac{3}{s^2+9} + \frac{1}{s^2+1} \right]$$

$$\textcircled{7} \quad f(t) = \begin{cases} 0, & 0 < t < 2 \\ 3, & t \geq 2 \end{cases}$$

$$L[f(t)] = \int_0^{\infty} f(t) e^{-st} dt$$

$$= \int_0^2 0 dt + \int_2^{\infty} 3 e^{-st} dt$$

$$= 3 \left. \frac{e^{-st}}{-s} \right|_2^{\infty}$$

$$= -\frac{3}{s} [0 - e^{-2s}]$$

$$= \frac{3e^{-2s}}{s}$$

# Shifting Theorem:-

$$\text{If } L[f(t)] = \int_0^{\infty} e^{-st} f(t) dt = F(s)$$

$$\text{then } L[e^{at} f(t)] = F(s-a)$$

$$\text{L.H.S} = L[e^{at} f(t)] = \int_0^{\infty} f(t) e^{-t(s-a)} dt$$

$$= \underline{F(s-a)}$$

Formulae:-

$$L(t^n) = \frac{n!}{s^{n+1}}$$

$$L(e^{at} t^n) = \frac{n!}{(s-a)^{n+1}}$$

$$L[e^{-at} t^n] = \frac{n!}{(s+a)^{n+1}}$$

$$L[e^{at} \sin bt] = \frac{b}{(s-a)^2 + b^2}$$

$$L[e^{at} \cos bt] = \frac{(s-a)}{(s-a)^2 + b^2}$$

$$L[e^{at} \sin bt] = \frac{b}{(s-a)^2 + b^2}$$

$$L[e^{-at} \cos bt] = \frac{s+a}{(s+a)^2 + b^2}$$

$$L[e^{at} \sinh bt] = \frac{b}{(s-a)^2 - b^2}$$

$$L[e^{-at} \sinh bt] = \frac{b}{(s+a)^2 - b^2}$$

$$L(t \sin at)$$

$$= \text{I.P of } L(t e^{ait}) \quad L(t) = \frac{1}{s^2}$$

$$= \text{I.P of } \left[ \frac{1}{(s-ai)^2} \times \frac{(s+ai)^2}{(s+ai)^2} \right]$$

$$= \text{I.P of } \left[ \frac{s^2 - a^2 + 2ias}{(s^2 + a^2)^2} \right]$$

$$L(t \sin at) = \frac{2as}{(s^2 + a^2)^2}$$

$$L(t \sin t) = \frac{2s}{(s^2 + 1)^2}$$

$$L(t \cos at) = \frac{s^2 - a^2}{(s^2 + a^2)^2}$$

$$L(t \cos t) = \frac{s^2 - 1}{(s^2 + 1)^2}$$

✓ marks Q.

### # Inverse Laplace Transform:-

$$L(t^n) = \frac{n!}{s^{n+1}}$$

$$L\left(\frac{t^n}{n!}\right) = \frac{1}{s^{n+1}}$$

$$L^{-1}\left(\frac{1}{s^{n+1}}\right) = \frac{t^n}{n!}$$

$$\therefore L^{-1}\left[\frac{1}{s}\right] = 1$$

$$L^{-1}\left[\frac{1}{s^2}\right] = t$$

$$L^{-1}\left[\frac{1}{s^3}\right] = \frac{t^2}{2}$$

$$L^{-1}\left(\frac{1}{s-a}\right) = e^{at}$$

$$L^{-1}\left(\frac{1}{s+a}\right) = e^{-at}$$

$$L^{-1}\left(\frac{1}{s^2 + a^2}\right) = \frac{1}{a} \sin at$$

$$L^{-1}\left(\frac{s}{s^2 + a^2}\right) = \cos at$$

$$L^{-1}\left(\frac{1}{s^2 - a^2}\right) = \frac{1}{a} \sinh at$$

$$L^{-1}\left(\frac{s}{s^2 - a^2}\right) = \cosh at$$

Q. Find  $L^{-1}\left[\frac{1}{2s-5}\right]$ ,  $L^{-1}\left[\frac{1}{s(s+1)}\right]$ ,  $L^{-1}\left[\frac{1}{(s+1)^2}\right]$

$L^{-1}\left[\frac{s}{(s+a)^2}\right]$ ,  $L^{-1}\left[\frac{s+23}{s^2-4s+13}\right]$ ,  $L^{-1}\left[\frac{s-2}{s^2+2s+2}\right]$

→ ①  $L^{-1}\left[\frac{1}{2s-5}\right] = \frac{1}{2} L^{-1}\left[\frac{1}{s-\frac{5}{2}}\right] = \frac{1}{2} e^{\frac{5}{2}t}$

②  $L^{-1}\left[\frac{1}{s(s+1)}\right] = L^{-1}\left[\frac{1}{s} - \frac{1}{s+1}\right] = 1 - e^{-t}$

③  $L^{-1}\left[\frac{1}{(s+1)^2}\right] = t e^{-t}$

④  $L^{-1}\left[\frac{s}{(s+a)^2}\right] = L^{-1}\left[\frac{s+a-a}{(s+a)^2}\right] = e^{-at} - a t e^{-at}$   
 $= e^{-at}(1-at)$

⑤  $L^{-1}\left[\frac{s+23}{s^2-4s+13}\right]$

$= L^{-1}\left[\frac{s+23}{s^2-2 \times 2s+4+13-4}\right]$

$= L^{-1}\left[\frac{s-2+25}{(s-2)^2+3^2}\right]$

$= e^{2t} \cos 3t + \frac{25}{3} e^{2t} \sin 3t$


⑥  $L^{-1}\left[\frac{s-2}{s^2+2s+1+2-1}\right]$

$= L^{-1}\left[\frac{s+1-3}{(s+1)^2+1}\right]$

$= e^{-t} \cos t - 3 e^{-t} \sin t$

$= e^{-t} [\cos t - 3 \sin t]$

## # Unit Step function:-



The unit step function


$u(t-a)$  is defined by —

$$u(t-a) = \begin{cases} 0, & 0 < t < a \\ 1, & t \geq a \end{cases}$$

$$\begin{aligned} \text{then } \mathcal{L}[u(t-a)] &= \int_0^{\infty} e^{-st} u(t-a) dt \\ &= \int_0^a 0 dt + \int_a^{\infty} 1 e^{-st} dt \\ &= \frac{1}{s} [e^{-as} - e^{-\infty s}] \end{aligned}$$

$$\boxed{\mathcal{L}[u(t-a)] = \frac{e^{-as}}{s}}$$

①



$$= u(t-a) - u(t-b)$$

$$= \frac{e^{-as}}{s} - \frac{e^{-bs}}{s} = \frac{1}{s} [e^{-as} - e^{-bs}]$$

## # Periodic function:-

→ A fn.  $f(t)$  is said to be periodic with the period ' $T$ ', if  $f(t+T) = f(t)$ , where  $T$  is the least time number.

ex:-  $f(t) = \sin t$   
 $f(t+2\pi) = \sin(t+2\pi)$   
 $= \sin t$   
 $f(t+2\pi) = f(t)$

## # Laplace Transform of Periodic function:-

Let  $f(t)$  be a periodic fn with period  $T$ , then the Laplace Transform of the periodic fn is defined by —

$$L[f(t)] = \frac{1}{1-e^{-sT}} \int_0^T e^{-st} \cdot f(t) dt$$

ex:-  $f(t) = \begin{cases} 3, & 0 < t < 2 \\ 0, & 2 < t < 4 \end{cases}$

and  $f(t+4) = f(t)$ , then find  $L[f(t)]$ .

$$\begin{aligned} \rightarrow L[f(t)] &= \frac{1}{1-e^{-4s}} \left[ \int_0^2 e^{-st} \times 3 dt + \int_2^4 0 \right] \\ &= \frac{1}{1-e^{-4s}} \left[ \frac{3}{s} [e^{-2t} - 1] \right] \\ &= \frac{3(1/e^{-2s})}{s(1-e^{-4s})} \rightarrow (1/e^{-2s})(1+e^{-2s}) \\ &= \frac{3}{s(1+e^{-2s})} \end{aligned}$$

Q.  $f(t) = \begin{cases} 1, & 0 < t < b. \\ -1, & b < t < 2b \end{cases}$  ,  $f(t+2b) = f(t)$  , find  $L[f(t)]$

→  $L[f(t)] = \frac{1}{1 - e^{-2bs}} \left[ \int_0^b e^{-st} dt + \int_b^{2b} e^{-st} dt \right]$

Imp Result


$$= \frac{1}{1 - e^{-2bs}} \left[ \frac{1 - e^{-bs}}{s} + \frac{e^{-2bs} - e^{-bs}}{s} \right]$$

$$= \frac{1}{(1 - e^{-2bs})} \cdot \frac{1 - e^{-bs} + e^{-2bs} - e^{-bs}}{s}$$


$$= \frac{1}{(1 - e^{-2bs})} \left[ \frac{1 - 2e^{-bs} + e^{-2bs}}{s} \right] (1 - e^{-bs})$$

$$= \frac{1 - e^{-bs}}{s(1 + e^{-bs})}$$

Remember :  $e^{-bs}$   
Imp Result



ex:



$$L[f(t)] = ?$$

Here  $b=1$

$$L[f(t)] = \frac{1 - e^{-s}}{s(1 + e^{-s})}$$

Q. ~~Find the Laplace transform of the function~~

$$L[f(t)] = F(s) = \frac{5s^2 + 23s + 6}{s(s^2 + 2s + 2)}$$

then  $\lim_{t \rightarrow \infty} f(t) = \underline{\hspace{2cm}}$

- (a) 3 (b) 5 (c)  $\frac{17}{2}$  (d)  $\infty$

Q.  $f(t) = L^{-1} \left[ \frac{3s+1}{s^3 + 4s^2 + (k-3)s} \right]$

$\lim_{t \rightarrow \infty} f(t) = 1$

then  $k = \underline{\hspace{2cm}}$

→ A/c to F.V.T -  $\frac{1}{k-3} = 1 \Rightarrow k = \underline{\underline{4}}$

Q. The Laplace Transform -

$$L[f(t)] = \frac{1}{s^2(s+1)}$$

$f(t) = ?$

$$\rightarrow \frac{1}{s^2(s+1)} = \frac{A}{s^2} + \frac{B}{s} + \frac{C}{s+1}$$

$$\frac{1}{2} = 1 + a_1 + \frac{1}{2}$$

$$a_1 = -1$$

$$= t^2 - 1 + e^{-t}$$

# Probability

→ If an experiment is conducted under essentially given conditions upto  $n$  times, let  $m$  cases are favourable to an event  $E$  then the Probability of  $E$  is denoted by  $P(E)$  and is defined by -

$$P(E) = \frac{m}{n}$$

$$P(\bar{E}) = \frac{n-m}{n} = 1 - \frac{m}{n} = 1 - P(E)$$

$$\therefore P(E) + P(\bar{E}) = 1$$

Sum of success & failure is always 1.

## # Axioms of Probability:-

① Axiom of Positivity :-  $0 \leq P(E) \leq 1$ .

② Certainty :-  $P(S) = 1$ .

③ Union :-  $P(E_1 \cup E_2) = P(E_1) + P(E_2) - P(E_1 \cap E_2)$

Note:- If  $E_1$  and  $E_2$  are mutually Exclusive or Disjoint events  
i.e.,  $E_1 \cap E_2 = \phi$

then  $P(E_1 \cup E_2) = P(E_1) + P(E_2)$

## # Sample Space:-


The set of all possible outcomes in an experiment is said to be the sample space.

2 coins tossed  
 $S = \{HH, HT, TH, TT\}$

1 dice thrown  
 $S = \{1, 2, 3, 4, 5, 6\}$

2 dice thrown  
 $S = \{(1,1), (1,2), (1,3), \dots, (6,6)\} \rightarrow$  Total 36

Spade    club    diamond    Heart



A  
2  
3  
⋮  
10  
J  
Q  
K

13

### # Conditional Probability:-

Let  $S$  be the sample space, and Event  $E_1, E_2 \subset S$  then the conditional Probability of  $E_1$  after the occurrence of  $E_2$  is denoted by -

$$P\left(\frac{E_1}{E_2}\right) = \frac{P(E_1 \cap E_2)}{P(E_2)}, \quad (P(E_2) \neq 0)$$

$$P\left(\frac{E_2}{E_1}\right) = \frac{P(E_1 \cap E_2)}{P(E_1)}, \quad (P(E_1) \neq 0).$$

### # Random Variable:-

Mean:-  $\sum x_i P(x=x_i)$  if exists, it is called the mean of R.V.

$$E = \mu = \sum x_i P(x=x_i).$$

Variance of R.V  $\sum (x_i - \mu)^2 P(x=x_i)$  if exists, it is called the variance of R.V

$$\sigma^2 = \sum (x_i - \mu)^2 P(x=x_i)$$

$$= \sum (x_i^2 + \mu^2 - 2x_i\mu) P(x=x_i)$$

$$= \sum x_i^2 P(x=x_i) + \sum \mu^2 \cdot P(x=x_i) - \sum 2x_i\mu P(x=x_i)$$

$$= \sum x_i^2 P(x=x_i) + \mu^2 \sum P(x=x_i) - 2\mu \sum x_i P(x=x_i)$$

$$= \sum x_i^2 [P(x=x_i)] + \mu^2 - 2\mu^2$$

$$\boxed{\sigma^2 = \sum x_i^2 P(x=x_i) - \mu^2}$$

### ③ Standard Deviation:-

The standard deviation is the sq. root of variance and is denoted by  $\sigma$ .

ABOVE WERE FOR GENERAL DISTRIBUTION:

### # Binomial Distribution:-

$n$  = no. of trials


$p$  = prob. of success.

$q$  = prob. of failures.

$$p + q = 1$$

$$P(X = k) = {}^n C_k q^{n-k} p^k$$

$$\begin{aligned} \text{Mean} &= n \cdot p \\ \text{Variance} &= npq \\ \text{S.D.} &= \sqrt{npq} \end{aligned}$$



If  $p = q$


$$P(X = k) = {}^n C_k p^k$$

### # Poisson Distribution:-

$$P(X = k) = \frac{\lambda^k}{k!} e^{-\lambda}, \text{ where } \lambda \text{ is a parameter.}$$

$$\begin{aligned} \text{Mean} &= \lambda \\ \text{Variance} &= \lambda \\ \text{S.D.} &= \sqrt{\lambda} \end{aligned}$$

### # Normal Distribution:-



$$z = \frac{x - \mu}{\sigma}$$


$\mu$  = mean.

$\sigma$  = S.D.

Q. What is the chance that a Leap year will have 53 Sundays.

→  $52 \times 7 = 364$

for rest 2 days —



for Non leap year —  $P(53 \text{ Sunday}) = \frac{1}{7} =$

Q. From a deck of playing Cards - 2 Cards are drawn at Random. What will be the probab. that both Cards will be Kings? if —

①. the first Card is not replaced

②. Replacement is allowed.

→ ①  $\frac{{}^4C_1}{{}^{52}C_1} \times \frac{3}{51}$

$= \frac{4}{52} \times \frac{3}{51} = \frac{1}{221}$

②  $\frac{4}{52} \times \frac{4}{52} = \frac{1}{169}$

Q. A Bag Contains 3 red & 6 white and 7 black balls, what is the Prob that 2 balls drawn are white and Black.

→  $\frac{{}^6C_1 \times {}^7C_1}{{}^{16}C_2} = \frac{6 \times 7}{\frac{16 \times 15}{2}} = \frac{7}{20}$

Q. A problem in electronics is given to 3 students A, B, C whose chances of solving it are  $\frac{1}{2}, \frac{1}{3}, \frac{1}{4}$  respectively. What is the prob. that the prob. will be solved.

$$\begin{aligned} \rightarrow & \left. \begin{array}{l} P(A) = \frac{1}{2} \\ P(B) = \frac{1}{3} \\ P(C) = \frac{1}{4} \end{array} \right\} \text{individual} \\ & \swarrow \text{then} \\ & P(A) \cdot P(B) \cdot P(C) \\ & 1 - P(\bar{A} \cap \bar{B} \cap \bar{C}) \\ & = 1 - P(\bar{A}) P(\bar{B}) P(\bar{C}) \\ & = 1 - \frac{1}{2} \times \frac{2}{3} \times \frac{3}{4} \\ & = \frac{3}{4} \\ & = 75\% \text{ chances.} \end{aligned}$$

Q. A husband & wife appear in interview for 2 vacancies for the same post. The prob. of husband selection is  $\frac{1}{7}$  & that of wife selection is  $\frac{1}{5}$ . What is the Prob. that —

- ① both are selected.
- ② only one of them is selected.
- ③ None of them is selected.

$$\begin{aligned} \rightarrow & P(A) = \frac{1}{7} \\ & P(B) = \frac{1}{5} \end{aligned}$$

$$\textcircled{1} \quad P(A \cap B) = P(A) \cdot P(B) = \frac{1}{7} \times \frac{1}{5} = \frac{1}{35}$$

~~②  $P(A \cap B) + P(\bar{A} \cap \bar{B})$~~

$$\begin{aligned} & P(A \cap \bar{B}) + P(\bar{A} \cap B) \\ & = \frac{1}{7} \times \frac{4}{5} + \frac{6}{7} \times \frac{1}{5} \\ & = \frac{10}{35} = \frac{2}{7} \end{aligned}$$

$$\begin{aligned} \textcircled{3} \quad P(\bar{A} \cap \bar{B}) & = P(\bar{A}) \cdot P(\bar{B}) \\ & = \frac{6}{7} \times \frac{4}{5} = \frac{24}{35} \end{aligned}$$

Don't use  $1 - \frac{1}{35}$  by individual selection diff.

Q. An integer is selected at random from the first 200 digits. What is the prob. that the integer chosen is divisible by either 6 or 8?

→ 1, 2, 3, ..., 200

$\div 6 \rightarrow 6, 12, 18, \dots, 198$

$\div 8 \rightarrow 8, 16, 24, \dots, 200$

$\div 6 \& \div 8 \rightarrow 24, 48, \dots, 192$

$$\text{no. of integers } \div 6 = \frac{198}{6} = 33 = n(E_1)$$

$$\text{no. of integer } \div 8 = \frac{200}{8} = 25 = n(E_2)$$

$$\text{no. of integers } \div 6 \& \div 8 = \frac{192}{24} = 8 = n(E_1 \cap E_2)$$

$$\begin{aligned} n(E_1 \cup E_2) &= n(E_1) + n(E_2) - n(E_1 \cap E_2) \\ &= 33 + 25 - 8 \\ &= 50 \end{aligned}$$

$$\therefore P(E_1 \cup E_2) = \frac{50}{200} = \frac{1}{4}$$

Q. A hits a target 4 times in 5 <sup>shots</sup> ~~shots~~ → B → 3 times in 4 <sup>shots</sup> ~~shots~~

C → 2 in 3 shots.

The fire a volley. What is Prob. that 2 shots at least hit the

target?   
 ↓ means A, B, C then again ABC... continuous

→ A → 4 in 5

B → 3 in 4

C → 2 in 3

for at least 2 shots —

$$P(A \cap B \cap \bar{C}) + P(A \cap \bar{B} \cap C) + P(\bar{A} \cap B \cap C) + P(A \cap B \cap C)$$

$$= \frac{4}{5} \times \frac{3}{4} \times \frac{1}{3} + \frac{4}{5} \times \frac{1}{4} \times \frac{2}{3} + \frac{1}{5} \times \frac{3}{4} \times \frac{2}{3} + \frac{4}{5} \times \frac{3}{4} \times \frac{2}{3}$$

$$= \frac{12}{60} + \frac{8}{60} + \frac{6}{60} + \frac{24}{60} = \frac{50}{60} = \frac{5}{6}$$

For at least 1 shot  $\rightarrow 1 - P(\bar{A} \cap \bar{B} \cap \bar{C})$

Q In a college, 25% of student fail in Math, 15% failed in Chemistry, 10% failed in both Math and chemistry if a student is selected at random —

① - If he failed in chemo then what is the probability that he fails in Math.

② vice-versa

③ What is the prob. that he failed neither Mathematics, nor chemistry?  
 $\rightarrow$

$$\rightarrow P(M) = 0.25$$

$$P(C) = 0.15$$

$$P(M \cap C) = 0.10$$

$$\textcircled{1} P\left(\frac{M}{C}\right) = \frac{P(M \cap C)}{P(C)} = \frac{0.10}{0.15} = \frac{2}{3}$$

$$\textcircled{2} P\left(\frac{C}{M}\right) = \frac{P(C \cap M)}{P(M)} = \frac{0.10}{0.25} = \frac{2}{5}$$

$$\begin{aligned} \textcircled{3} P(\overline{M \cup C}) &= 1 - P(M \cup C) \\ &= 1 - [P(M) + P(C) - P(M \cap C)] \\ &= 1 - \left[ \cancel{\frac{1}{4}} \cdot 0.25 + 0.15 - 0.10 \right] \\ &= 1 - 0.3 \\ &= 0.7 \end{aligned}$$

Q. A single dice is thrown twice, what is the probability that their sum is neither 8 nor 9.

→  $n(S) = 36$

$$n(E_1) = \{(2,6), (3,5), (4,4), (5,3), (6,2)\}$$

$$\therefore P(E_1) = \frac{5}{36}$$

$$n(E_2) = \{(3,6), (6,3), (5,4), (4,5)\}$$

$$P(E_2) = \frac{4}{36}$$

∴ These are mutually Exclusive → *leg.* 8 आने से 9 नहीं आता  
9 आने से 8 नहीं आता

$$\therefore P(\overline{E_1 \cup E_2}) = 1 - P(E_1 \cup E_2)$$

$$= 1 - [P(E_1) + P(E_2)]$$

$$= 1 - \left[ \frac{5}{36} + \frac{4}{36} \right]$$

$$= \frac{27}{36} = \frac{3}{4}$$

Q. A pair of fair dice is thrown, if the two numbers appearing are different, find the Prob that -

- ① The sum is 6
- ② an ace appears
- ③ The sum is 4 or less

→  $n(S) = 36 - 6 = 30$

*Same no appearing.*

for Sum is 6 -

$$n(E_1) = \{(1,5), (2,4), (5,1), (4,2)\} \Rightarrow P(E_1) = \frac{4}{30} = \frac{2}{15}$$

for Ace Appears —  $P(E) = 0$  if option available

But Here Ace means 1

$$n(E_2) = \{(1,2), (1,3), (1,4), (1,5), (1,6), \dots \times 2\} = 10$$

$$\therefore P(E_2) = \frac{10}{30} = \frac{1}{3}$$

③ The sum is 4 or less

$$h(E) = \{(1,2), (1,3), (2,1), (3,1)\} = 4$$

$$P(E) = \frac{4}{20} = \frac{2}{15}$$

Binomial  
Type-2

Q. Team A has probability  $\frac{2}{3}$  of winning whenever it plays.  
Suppose A plays 4 games. find the prob. that A wins more than half of its games.

$$\rightarrow \text{Team A} \rightarrow \frac{2}{3}$$

$$n = 4$$

$$p = \frac{2}{3}$$

$$q = \frac{1}{3}$$

$$P(X > 2) = P(X=3) + P(X=4)$$

$$= {}^4C_3 \left(\frac{1}{3}\right)^1 \left(\frac{2}{3}\right)^3 + {}^4C_4 \left(\frac{2}{3}\right)^4$$

$$= {}^4C_3 \left(\frac{1}{3}\right)^1 \left(\frac{2}{3}\right)^3 + {}^4C_4 \left(\frac{2}{3}\right)^4$$

$$= \frac{4 \times 8}{81} + \frac{1 \times 16}{81} = \frac{48}{81} = \frac{16}{27}$$

How to know Binomial?

n choose k  
k

Given not imp  
asked imp, a/c to that  
p is selected

Q. In a bank 60% of all customers applied for a loan are rejected. If 4 new loans applications received. find the prob that 3 of them are accepted.

$$\rightarrow \begin{aligned} n &= 4 \\ p &= 0.4 \\ q &= 0.6 \end{aligned}$$

$$P(X=3) = {}^4C_3 (0.6)^1 (0.4)^3$$

$$= 4 (0.6)(0.4)^3$$

$$= 0.1536$$

Q. A player tosses 2 fair coins. He wins \$2 if 2 heads occurs and wins \$1 if 1 head occurs. On the other hand ~~even~~ he loses \$3, if no head occurs. find the expected value 'E' of the game.

→ \$2 → 2H  
 \$1 → 1H  
 - \$3 → 0H

fair game means —  
 no loss no gain //

fairness of game  $\propto$  mean value of game.

if mean = +ve → favourable to player  
 = -ve → Unfavourable to player  
 mean = 0 → Fair Game

$S = \{(H,H), (H,T), (T,H), (TT)\}$

$x_i$	\$2	\$1	-\$3
$P_i$	$\frac{1}{4}$	$\frac{2}{4}$	$\frac{1}{4}$

$$E(x) = \sum x_i P(x = x_i)$$

$$= 2\left(\frac{1}{4}\right) + 1 \times \left(\frac{2}{4}\right) - 3\left(\frac{1}{4}\right)$$

$$= \frac{1}{2} + \frac{1}{2} - \frac{3}{4}$$

$$= \frac{1}{4} > 0 \rightarrow \text{favourable to Player.}$$

Q. 4 fair coins are tossed simultaneously. find the prob. that at least one head and one tail turn up.

→  $n = 4$   
 $P = 2/2$

$$P(X=1) + P(X=2) + P(X=3) \quad \left| \quad = 1 - [P(X=4) + P(X=0)]\right.$$

$$= \left( {}^4C_1 + {}^4C_2 + {}^4C_3 \right) \left( \frac{1}{2} \right) \quad \left| \quad = 1 - \left[ \frac{1+1}{16} \right]\right.$$

$$= \frac{4+6+4}{16} = \frac{14}{16} = \frac{7}{8} \quad \left| \quad = 1 - \frac{1}{8} = \frac{7}{8}\right.$$

Q. Five fair coins are tossed simultaneously. Find the prob. that at least one head turn up.

$$\begin{aligned} &\rightarrow 1 - P(X=0) \\ &= 1 - {}^5C_0 \left(\frac{1}{2}\right)^5 \\ &= 1 - \frac{1}{32} = \frac{31}{32} \end{aligned}$$

Q. 4 fair coins are tossed simultaneously. Find the Prob. of event the no. of times heads show up is more than the no. of times Tails show up.

$$\begin{aligned} &\rightarrow P(X=3) + P(X=4) \\ &= {}^4C_3 + {}^4C_4 + \left(\frac{1}{2}\right)^4 \\ &= \frac{4+1}{16} = \frac{5}{16} \end{aligned}$$

Q. There are 5 duplicate and 10 original items in an automobile shop. 3 items are bought up by customer at random. Find the Prob that none of the item is duplicate.

$$\begin{aligned} &\rightarrow \text{means all original.} \\ &= \frac{{}^{10}C_3}{{}^{15}C_3} = \frac{10 \times 9 \times 8}{15 \times 14 \times 13} = \frac{24}{91} \end{aligned}$$

Q. Out of 800 families with 4 children each how many families would be expected to have —

- ① 2 boys and 2 girls.
- ② At least one boy
- ③ At most 2 girls.

for 1 family —  
 $n=4$   
 $p=q=\frac{1}{2}$

$$\therefore P(X=2) = {}^4C_2 \left(\frac{1}{2}\right)^4 = \frac{6 \times \frac{1}{16} \times 800}{16} = \underline{\underline{300}}$$

Poisson distribution  
 $n$  is large (very large)  
 Binomial distribution  
 individual ref. &  
 Not applicable

Now  
 for 800 family

$$\begin{aligned}
 \textcircled{2} \quad P(X \geq 1) &= 1 - P(X=0) \\
 &= 1 - {}^4C_0 \left(\frac{1}{2}\right)^4 \\
 &= 1 - \frac{1}{16} \\
 &= \frac{15}{16} \times 800 \\
 &= 750
 \end{aligned}$$

$$\begin{aligned}
 \textcircled{3} \quad P(X \leq 2) &= P(X=0) + P(X=1) + P(X=2) \\
 &= \left({}^4C_0 + {}^4C_1 + {}^4C_2\right) \left(\frac{1}{2}\right)^4 \\
 &= \frac{1+4+6}{16} \\
 &= \frac{11}{16} \times 800 \\
 &= \underline{\underline{550}}
 \end{aligned}$$

Q. A book of 300 pages containing 30 printing mistakes. Assuming that these errors are randomly distributed throughout the book and  $X$  is the no. of errors per page has a poisson's distribution. What is the prob. that 20 pages selected at random will be free of errors.

→ error in page  $P = \frac{30}{300} = \frac{1}{10}$

Given  $n = 20$

$$\begin{aligned}
 \lambda &= np \\
 &= 20 \times \frac{1}{10}
 \end{aligned}$$

$$\boxed{\lambda = 2}$$

$$P(X=0) = \frac{\lambda^0}{0!} e^{-\lambda} = e^{-2} = \underline{\underline{0.135}}$$

Prob  
 When in Poisson distribution  $\lambda$  is not mentioned, we will find it by Binomial  
 i.e.  $\lambda = np$   
 $= 20 \times \frac{1}{10} = 2$   
 $\lambda = 2$

Q. 1000 Students had written an examination with the mean of test = 35  $\longrightarrow$  and S.D = 5. Assuming the distribution to be normal. find how many students marks —

- (a) lie between 25 & 40.
- (b) How many got more than 40.
- (c) How many got below 20.

Given  $P(0 \leq z \leq 1) = 0.3415$

$P(0 \leq z \leq 2) = 0.4772$


$P(0 \leq z \leq 3) = 0.449$

$\rightarrow \mu = 35$

$\sigma = 5$

$z = \frac{x - \mu}{\sigma}$

$P(\underset{x_1}{25} \leq x \leq \underset{x_2}{40})$



for  $x_1 = 25$   $\rightarrow z_1 = \frac{25 - 35}{5} = -2$

for  $x_2 = 40$   $\rightarrow z_2 = \frac{40 - 35}{5} = 1$

$\therefore P(25 \leq x \leq 40) = P(-2 \leq z \leq 1)$

$= P(-2 \leq z \leq 0) + P(0 \leq z \leq 1)$

$= P(0 \leq z \leq 2) + P(0 \leq z \leq 1)$

$= 0.4772 + 0.3415$

$= 0.8187$

$= 0.819 \times 1000 \rightarrow$  Total Students

$= 819$

$P(x \geq 40) = P(z \geq 1)$

$= 0.5 - P(0 \leq z \leq 1)$

$= 0.5 - 0.3415$

$= 0.1585$

$= 0.159 \times 1000$


$= \underline{159}$

①.  $P(X \leq 20)$

$$Z = \frac{20 - 35}{5} = \frac{-15}{5} = -3.$$

$$\begin{aligned} P(Z \leq -3) &= P(X < 20) = 0.5 - P(0 \leq Z \leq 3) \\ &= 0.5 - 0.499 \\ &= 0.001 \times 1000 \\ &= \underline{1 \text{ student}}. \end{aligned}$$

②. for a Random Variable —  
 $(-\infty < x < \infty)$  following a normal distribution with the ~~mean~~  
 Mean is 100. If the prob. of ~~the~~  $P(X \geq 110) = \alpha$   
 then  $P(90 \leq x \leq 110) = \underline{\hspace{2cm}}$




$$P(X \geq 110) = \alpha$$

$$P(90 \leq x \leq 110) = 1 - 2\alpha.$$

③. Let  $X$  be the normal variant with Mean = 1 &  $\sigma^2 = 4$   
 then the Prob. of  $P(X < 0)$  —

- (a) 0.5    (b)  $> 0$  and  $< 0.5$     (c)  $> 0.5$  and  $< 1$     (d) 1.



$$\mu = 1$$


$$\sigma^2 = 4$$

$$\sigma = 2$$

$$Z = \frac{x - \mu}{\sigma} = \frac{-1}{2} = -0.5$$

$$\begin{aligned} P(X < 0) &= P(Z < -0.5) \\ &= 0.5 - P(0 \leq Z \leq 0.5) \end{aligned}$$

Q. A box contain 3 blue and 4 red ball. Another identical Box contain 2 blue ball and 5 red ball. One ball is selected at Random from one of the two boxes and it is red. The Prob. that it came from the first box —



$$P(E_1) = P(E_2) = \frac{1}{2}$$

$$P\left(\frac{A}{E_1}\right) = \frac{4}{7}, \quad P\left(\frac{A}{E_2}\right) = \frac{5}{7}$$

$$P\left(\frac{E_1}{A}\right) = \frac{P(E_1) \cdot P\left(\frac{A}{E_1}\right)}{P(E_1) \cdot P\left(\frac{A}{E_1}\right) + P(E_2) \cdot P\left(\frac{A}{E_2}\right)}$$

$$= \frac{\frac{1}{2} \left(\frac{4}{7}\right)}{\frac{1}{2} \left(\frac{4}{7}\right) + \frac{1}{2} \left(\frac{5}{7}\right)}$$

$$= \frac{\frac{4}{7}}{\frac{9}{7}} = \frac{4}{9}$$

Q. Box P has 2 Red & 3 blue balls. Box Q has 3 Red & 1 Blue ball. The Prob. of selecting boxes P and Q are  $\frac{1}{3}$  and  $\frac{2}{3}$  respectively. One ball is selected from one of two boxes and it is red. the Prob that it came from box P.

$$\rightarrow P(P) = \frac{1}{3}, \quad P(Q) = \frac{2}{3}, \quad P\left(\frac{A}{P}\right) = \frac{2}{5}, \quad P\left(\frac{A}{Q}\right) = \frac{3}{4}$$

$$P\left(\frac{P}{A}\right) = \frac{P(P) \cdot P\left(\frac{A}{P}\right)}{P(P) \cdot P\left(\frac{A}{P}\right) + P(Q) \cdot P\left(\frac{A}{Q}\right)} = \frac{\frac{1}{3} \cdot \frac{2}{5}}{\frac{1}{3} \cdot \frac{2}{5} + \frac{2}{3} \cdot \frac{3}{4}} = \frac{\frac{2}{15}}{\frac{2}{15} + \frac{2}{2}} = \frac{\frac{2}{15}}{\frac{2}{15} + \frac{20}{15}} = \frac{2}{22} = \frac{1}{11}$$

Q. The Probability density fn of the form  $f(x) = k e^{-\alpha|x|}$ ,  $x \in (-\infty, \infty)$ , then the value of  $k$  —

$$\rightarrow \int_{-\infty}^{\infty} f(x) dx = 1 \Rightarrow \int_{-\infty}^{\infty} k e^{-\alpha|x|} dx = 1 \Rightarrow 2 \int_0^{\infty} k e^{-\alpha x} dx = 1$$

$$\Rightarrow \frac{2k}{\alpha} [e^{-\infty} - e^{-0}] = 1$$

$$\Rightarrow \frac{-2k}{\alpha} (0 - 1) = 1$$

$$\Rightarrow k = \frac{\alpha}{2} = 0.5\alpha$$

Q. Two Bag Contain 10 lotos each and the colors in each bag are numbered from 1 to 10. One coin is drawn at random from each bag. What is the prob that one of the coin has value (1, 2, 3, 4) while the other has value (7, 8, 9 or 10)...

## Differential Eq<sup>n</sup>:-

— Formation of Diff<sup>n</sup> eq<sup>n</sup> → diff. upto no. of constants.

— Solution of Diff. eq<sup>n</sup>

— Variable - Seperable — normal  
— reducible

